



Polymorphisms of COMT Val158Met and the Risk of Hepatocellular Carcinoma Development among Egyptian HCV Infected Patients

Eman Labib^{1&2}, Hany M. Ibrahim^{2*}, Mai El-Daly^{1&3}, Heba S. Ghanem¹, Ibrahim A. El-Elaimy², Mohamed Abdel-Hamid⁴

1- Clinical Pathology Department, National Liver Institute, Menoufia University, Shebin El-Kom, Egypt.

2- Department of Zoology, Faculty of Science, Menoufia University, Shebin El-Kom, Egypt.

3-Special Infectious Agents Unit, King Fahd Medical Research Center, King Abdulaziz University, Jeddah, Saudi Arabia.

4-Microbiology Department, Faculty of Medicine, Minia University, Minia, Egypt.

E-Mail : hany.mohamed@science.menofia.edu.eg or hanyibrahimeg@gmail.com

ARTICLE INFO

Article History

Received: 15/8/2018

Accepted: 20/9/2018

Keywords:

Hepatocellular Carcinoma;
Hepatitis C Virus; COMT;
Egypt

ABSTRACT

Chronic viral hepatitis is the most important risk factor for progression to hepatocellular carcinoma (HCC) in Egypt. Catechol-O-methyltransferase (COMT) plays a central role in DNA repair and estrogen-induced carcinogenesis. Many recent epidemiologic studies have investigated the association between the COMT Val158Met polymorphism and cancer risk.

In the current study, we investigated the association between COMT Val158Met A/G variations and the risk of HCC development among Egyptian HCV infected patients. This study was conducted on two groups; HCC patients group and healthy control group each included 100 subjects. Single nucleotide polymorphisms (SNPs) have been studied for COMT using real-time PCR.

Our results suggest that COMT is not associated with HCC risk among HCV infected patients. A significant increase was detected regarding the different COMT SNPs in the aged category more than 50 years compared to younger patients. No significant difference was detected on the level of gender, liver cirrhosis, biochemical and hematological parameters.

In conclusion, COMT Val158Met polymorphism could not be considered as a risk factor for HCC risk among hepatitis C virus patients in Egypt.

INTRODUCTION

Hepatocellular carcinoma (HCC) is the most widely recognized primary liver tumor (Balogh et al., 2016; Abdel-Hamid et al., 2018). Incidence varies widely between geographical areas, probably because of variations in the exposure to hepatitis virus and other environmental pathogens (El-Serag, 2001). The clinical risk factors include cirrhosis, chronic

hepatitis C virus (HCV) infection, chronic hepatitis B virus (HBV) infection, co-infection of HBV and HCV viruses, schistosomiasis, aflatoxin B1 (AFB1), and alcoholism. The environmental risk factors include pesticides and cigarette smoking (Anwar et al., 2008). HCV infection causes liver inflammation and a variable grade of damage to the organ that over decades can lead to cirrhosis (Ikeda et al., 1993; Tsukuma et al., 1993; Lauer and Walker, 2001). Egypt has one of the highest prevalence rates of HCV infection. Disease progression is influenced by additional factors such as duration of infection, age at infection, gender, co-infection with HBV, and the level of HCV viremia and its genotype (Anwar et al., 2008).

As with many cancers, variants of genes involved in multistage carcinogenesis may determine an individual's susceptibility to developing HCC. Single nucleotide polymorphisms (SNPs) are the most common type of genomic sequence variation and are thought to be associated with population diversity, susceptibility to disease, and individual response to drug treatment (Shastry, 2002).

Epidemiological and endocrinological studies indicate that sex hormones control normal growth and development of target organs, but can also influence susceptibility to disease and to neoplastic transformation. Researchers implicate sex hormones in the pathogenesis of several cancer types, such as ovary, prostate and breast (Henderson and Feigelson, 2000).

Catechol-O-methyltransferase (COMT) is the main enzyme sequentially participating in the pathways of estrogen and androgen biosynthesis and inactivation. COMT is expressed at high levels in liver, kidney, endometrium and breast (Guldberg and Marsden, 1975). A G-to-A transition at codon 158 leads to substitution of valine to methionine causes a 3–4-fold reduction of enzyme activity (Lachman et al., 1996).

COMT gene is localized on chromosome 22q11.2 and contains six exons

(Grossman et al., 1992). It is a vital enzyme participates in the inactivation of endogenous catecholamine and catechol estrogens which have the capability to destroy DNA and carcinogenetic agents (Cavalieri et al., 1997). So, the damage, the loss of or changes in COMT is hypothetically contributed to genomic instability and tumor formation.

Several evidence studies indicate that COMT polymorphisms can affect individual susceptibility to developing cancers (Garner et al., 2002; Sasak et al., 2003; Shen et al., 2006; Tanaka et al., 2006; Zhang et al., 2008). Functional polymorphisms of these genes have been studied in association with hormone-induced cancers (Reichardt et al., 1995; Febbo et al., 1999; Lunn et al., 1999; Makridakis et al., 1999; Jaffe et al., 2000; Haiman et al., 2001; Yamada et al., 2001), but they have been poorly investigated in the setting of chronic liver disease (Yu et al., 2001). The aim of the current study was to investigate the association between COMT Val158Met (rs4680) and the HCC risk among HCV patients in Egypt.

MATERIALS & METHODS

This study was a case-control study. It was conducted on a total number of 200 individuals divided into two groups, group I included 100 HCC cases that were recruited from National Cancer Institute, Cairo University and group II included 100 healthy subjects negative for anti-HCV and HBsAg were recruited from National Liver Institute, Menoufia University (age and gender were considered). Any healthy subject has a history of tumor, liver disease, renal disease, coronary artery disease or other metabolic disorders was excluded (Hao et al., 2014). A written agreement was gained from all contributors and the institutional ethical committee permitted the study.

Ten ml of blood was withdrawn from all patients and healthy controls under a complete aseptic condition in vacutainer tubes containing EDTA. All samples were centrifuged at 2000 rpm for 10 min, plasma and buffy coat were separated, aliquoted and stored at -80°C. Plasma samples were used

for the determination of HCV Ab, HBsAg, and AFP using ADVIA Centaur CP (Switzerland) according to the manufacturer instructions. Assessment of aspartate aminotransferase (AST), alanine aminotransferase (ALT), total bilirubin (TBIL), and Albumin (ALB) was measured using Beckman CX4 chemistry analyzer (USA). Hemoglobin (HB) and platelets were measured using Sysmex XT-1800i (Japan). Also, plasma was used to perform in-house RT HCV PCR for the detection of HCV (Abdel-Hamid et al., 1997). In patients, compensated cirrhosis was determined by Fibro-scan™ > 12.5 kPa (Ibrahim et al., 2018).

Genomic DNA was isolated from the buffy coat of cases and healthy control samples using a DNA isolation kit (QiAamp DNA mini kit, Qiagen, Hilden, Germany). Amplification of genomic DNA samples was used to detect the genetic polymorphisms COMT Val158Met (rs4680) (González-Castro et al., 2013). One of the allelic probes of the gene was labeled with FAM dye and the other with the fluorescent VIC dye using real-time PCR System (Applied Biosystems: Foster City, CA, USA). The PCR reaction was carried out using a TaqMan universal master mix (Applied Biosystems: Foster City, CA, USA) at a probe concentration of 20X. The reaction was performed in a 96-well format in a total reaction volume of 25 µl using 20 ng of genomic DNA. The reaction was heated for 2 min at 50°C, then 10 min at 95°C, followed by 40 cycles of 95°C for 15 sec and 60°C for 1.5 min. Then the fluorescence intensity of each well in the TaqMan assay plate was read.

Data were collected and entered to the computer using SPSS 20 (Chicago, Inc, Illinois). Quantitative data were shown as mean and standard deviation (SD) or expressed as frequency and percent. Chi-

square test and Fisher exact test were used to measure the association between qualitative variables as appropriate. Mann Whitney and independent sample t-tests were done to measure the association between two quantitative variables as appropriate. The Kruskal-Wallis and ANOVA tests were done to measure the association between more than two quantitative variables as appropriate.

RESULTS

Table 1 showed that there is no significant ($P = 0.607$) difference was detected on the level of COMT polymorphisms where (AA as wild genotype), (AG as heterozygous genotype) and (GG as mutant genotype) were detected in HCC group as 45(45.0%), 27(27.0%) and 28(28.0%), respectively, while the indicated genotypes were found in healthy control as 48(48.0%), 21(21.0%) and 31(31.0%), respectively.

Based on gender, no significant percentage ($P = 0.463$) was demonstrated on the levels of AA, AG, and GG polymorphisms in male patients 34(75.6%), 21(77.8%) and 18(64.3%), and in female patients 11(24.4%), 6(22.2%) and 10(35.7%), respectively. Interestingly, the higher percentage was detected with male patients than female (Table 2). On the level of age, a significant increase ($P = 0.018$) was detected regarding the different COMT SNPs in the aged category more than 50 years compared to younger patients. Moreover, No significant difference ($P = 0.138$) was detected in the percentage of liver cirrhosis among the different COMT genetic variants (Table 2).

Studying COMT SNP variants with laboratory findings among patients group showed no significant association between the COMT genotypes and biochemical and hematological studied parameters (Table 3).

Table (1): Comparison of genotype and allele distributions of COMT between the studied groups.

		HCC N = 100	Control N = 100	P-value	
COMT	Genotypes	AA AG GG	45 (45.0%) 27 (27.0%) 28 (28.0%)	48 (48.0%) 21 (21.0%) 31 (31.0%)	0.607
			Data are presented in terms of numbers of patients (%).		

Table (2): The gender, age and liver cirrhosis comparison among the COMT genotypes in the HCC group.

		COMT			P-value
		AA N = 45	AG N = 27	GG N = 28	
Gender	Male	34 (75.6%)	21 (77.8%)	18 (64.3%)	0.463
	Female	11 (24.4%)	6 (22.2%)	10 (35.7%)	
Age	≤50 years	4 (8.9%)	7 (25.9%)	10 (35.7%)	0.018*
	>50 years	41 (91.1%)	20 (74.1%)	18 (64.3%)	
Liver cirrhosis	36/100	14 (31.1%)	13 (48.1%)	9 (32.1%)	0.138

Data are presented in terms of numbers of patients (%).

*P-value < 0.05 was considered significant.

Table (3): The laboratory parameters among the COMT genotypes in HCC group.

	COMT			P-value
	AA N = 45	AG N = 27	GG N = 28	
AFP (ng/ml)	461.3 ± 495.4	599.8 ± 492.6	467.3 ± 467.8	0.400
ALT (IU/L)	52.5 ± 26.70	60.0 ± 38.4	40.6 ± 34.3	0.160
AST (IU/L)	46.5 ± 18.8	70.8 ± 58.8	53.3 ± 36.5	0.234
TBIL (mg/dL)	2.4 ± 1.7	2.6 ± 2.7	2.4 ± 2.2	0.921
ALB (g/dL)	2.9 ± 0.2	3.0 ± 0.4	3.1 ± 0.5	0.376
HB (g/dL)	11.3 ± 1.2	10.6 ± 1.1	11.0 ± 1.3	0.164
Platelets (10 ³ /µL)	117.1 ± 30.6	114.4 ± 35.2	121.9 ± 50.4	0.783

Data are presented in terms of Mean ± SD.

DISCUSSION

COMT is an enzyme belongs to estrogen metabolism enzymes that are the main enzymes involved in disrupting catechol estrogens, which are the most metabolites of estrogens and can cause oxidative DNA damage. It has been shown that Val158Met at codon 158 in exon 4 makes a severe decrease in the enzyme activity (Lachman et al., 1996; Lee et al., 2005).

Some studies have established that estrogen metabolites can bind to DNA and initiate damage, predicting that the estrogens might be endogenous genotoxic agents that can cause genetic alteration and begin tumor development (Cavalieri et al., 2000; Newbold et al., 2000). This concept is supported by the result that women with fewer quantities of the enzymes responsible for removing reactive estrogen metabolites have a higher risk of breast cancer development (Lavigne et al., 1997). Based on the obvious relevance of estrogens in human hepatocarcinogenesis (Neuberger et al., 1986), it is sensible to guess that the same mechanism may occur in HCC.

In our study, there is no association between COMT gene polymorphisms and HCC in Egyptian patients infected with HCV. Consequently, these results are indicating that COMT polymorphism is not associated with HCC risk related to HCV infection. In agreement with our findings, the previous report recorded that COMT Val158Met is not related to the risk of HCV-induced HCC (Rossi et al., 2003). Furthermore, another study found no association between the functional polymorphism of COMT and the risk of HCC caused by HBV infection (Yuan et al., 2008). Other records found to support the theory that estrogen levels are not involved in the modulation of liver disease outcome, which comes from observation that no effect was demonstrated with COMT polymorphism in the studies of other diseases, such as breast and ovarian cancer, where estrogen plays an essential role

(Lavigne et al., 1997; Thompson et al., 1998).

In disagreement with our findings, Yin et al. (2004) had reported that women who have low COMT Val158Met allele activity are strongly associated with the risk of developing HCC in Taiwanese females. The conflicting results could be attributable to differences in demography, ethnicity, lifestyles, type of viral infections, and clinical settings. In addition, other methodological factors in studies, such as small sample size, inadequate adjustment for confounding factors, or failure to correct multiple tests, may lead to inconsistent results.

In the current study, a higher percentage of COMT SNP variants were detected in aged patients than younger ones. Several studies found an obvious association between age and HCC or HCV infection (Abdel-Wahab et al., 2007; Do Carmo et al., 2012; Abdel-Hamid et al., 2018). The intimate relation between HCC, HCV and age might be reflected on COMT SNP studied in this current report.

Lastly, cancer is an illness as a result of a multifactor reaction between different genetic and environmental risk factors. So, the gene only or an environmental factor only is not enough to explain the risk to cancer. In conclusion, our findings do not support associations between the polymorphisms of COMT Val158Met gene and the risk for hepatocellular carcinoma among HCV infected patients in Egypt.

Conflict of Interest:

The author declares no conflict of interest.

REFERENCES

- Abdel-Hamid M, Edelman DC, Highsmith WE, Constantine NT. (1997). Optimization, assessment, and proposed use of a direct nested reverse transcription-polymerase chain reaction protocol for the detection of hepatitis C virus. *Journal of Human Virology* 1: 58–65.

- Abdel-Hamid M, Nada OH, Ellakwa DE, Ahmed LK. (2018). Role of Myeloperoxidase in hepatitis C virus related hepatocellular Carcinoma. *Meta Gene* 18: 1–8.
- Abdel-Wahab M, El-Ghawalby N, Mostafa M, Sultan A, Ezzat F. (2007). Epidemiology of hepatocellular carcinoma in Lower Egypt, Mansoura Gastroenterology center. *Hepato-Gastroenterology* 54 (73): 157–162.
- Anwar WA, Khaled HM, Amra HA, El-Nezami H, Loffredo CA. (2008). Changing pattern of hepatocellular carcinoma (HCC) and its risk factors in Egypt: possibilities for prevention. *Mutation Research* 659: 176–184.
- Balogh J, Victor D, Asham EH, Burroughs SG, Boktour M, Saharia A, Li X, Ghobrial RM, Monsour HP. (2016). Hepatocellular carcinoma: A Review. *Journal of Hepatocellular Carcinoma* 3: 41–53.
- Cavalieri E., Frenkel K, Liehr JG, Rogan E, Roy D. (2000). Estrogens as endogenous genotoxic agents—DNA adducts and mutations. *Journal of National Cancer Institute Monographs* 27: 75–93.
- Cavalieri EL, Stack DE, Todorovic R, Dwivedy I, Higginbotham S, Johansson SL, Patil D, Gross ML, Gooden JK, Ramanathan R, Cerny RL, Rogan EG. (1997). Molecular origin of cancer: catechol estrogen-3, 4-quinones as endogenous tumor initiators. *Proceedings of National Academy of Science of the USA* 94(20): 10937–10942.
- Do Carmo RF, de Almeida D, Aroucha DB, Vasconcelos LR, de Moraes AC. (2012). Plasma myeloperoxidase levels correlate with in chronic hepatitis C. *Human Immunology* 73 (11) 1127–1131.
- El-Serag HB. (2001). Epidemiology of hepatocellular carcinoma. *Clinical Liver Disease* 5: 87–107.
- Febbo PG, Kantoff PW, Platz EA, Casey D, Batter S, Giovannucci E, Hennekens CH, Stampfer MJ. (1999). The V89L polymorphism in the 5a-reductase type 2 gene and risk of prostate cancer. *Cancer Research* 59: 5878–5881.
- Garner EI, Stokes EE, Berkowitz RS, Mok SC, Cramer DW. (2002). Polymorphisms of the estrogen-metabolizing genes CYP17 and catechol-O-methyltransferase and risk of epithelial ovarian cancer. *Cancer Research* 62: 3058–3062.
- González-Castro TB, Tovilla-Zárate C, Juárez-Rojop I, Pool García SP, Genis A, Nicolini H, Narváez LL. (2013). Distribution of the Val108/158Met polymorphism of the COMT gene in healthy Mexican population. *Gene* 526(2): 454–458.
- Grossman MH, Emanuel BS, Budarf ML. (1992). Chromosomal mapping of the human catechol-O-methyltransferase gene to 22q11.1–q11.2. *Genomics* 12(4): 822–825.
- Guldberg HC, Marsden CA. (1975). Catechol-O-methyl transferase: pharmacological aspects and physiological role. *Pharmacological Reviews* 27: 135–206.
- Haiman CA, Stampfer MJ, Giovannucci E, Ma J, Decalo NE, Kantoff PW, Hunter DJ. (2001). The relationship between a polymorphism in CYP17 with plasma hormone levels and prostate cancer. *Cancer Epidemiology, Biomarkers & Prevention* 10: 743–748.
- Hao YX, Wang JP, Zhao LF. (2014). Associations between three common MicroRNA polymorphisms and hepatocellular carcinoma risk in Chinese. *Asian Pacific Journal of Cancer Prevention* 14: 6601–6604.
- Henderson BE, Feigelson HS. (2000). Hormonal carcinogenesis. *Carcinogenesis* 21: 427–433.
- Ibrahim HM, Abdel Ghaffar FR, El Shaer RE, Madian MA. (2018). Prevalence

- of Epstein-Barr Virus and Hepatitis B Virus Infections among Chronic HCV Patients Attending Kafer El Sheikh Liver and Heart Institute, Egypt. *Journal of Advances in Medicine and Medical Research* 26(8): 1-8.
- Ikeda K, Saitoh S, Koida I, Arase Y, Tsubota A, Chayama K, Kumada H, Kawanishi M. (1993). A multivariate analysis of risk factors for hepatocellular carcinogenesis: a prospective observation of 795 patients with viral and alcoholic cirrhosis. *Hepatology* 18: 47-53.
- Jaffe JM, Malkowicz SB, Walker AH, MacBride S, Peschel R, Tomaszewski J, Arsdalen KV, Wein AJ, Rebbeck TR. (2000). Association of SRD5A2 genotype and pathological characteristics of prostate tumors. *Cancer Research* 60: 1626-1630.
- Lachman HM, Papolos DF, Saito T, Yu YM, Szumlanski CL, Weinshilboum RM. (1996). Human catechol-O-methyltransferase pharmacogenetics: description of a functional polymorphism and its potential application to neuropsychiatric disorders. *Pharmacogenetics* 6: 243-250.
- Lauer GM, Walker BD. (2001). Hepatitis C virus infection. *The New England Journal of Medicine* 345: 41-52.
- Lavigne JA, Helzlsouer KJ, Huang HY, Strickland PT, Bell DA, Selmin O, Watson MA, Hoffman S, Comstock GW, Yager JD. (1997). An association between the allele coding for a low activity variant of catechol-O-methyltransferase and the risk for breast cancer. *Cancer Research* 57: 5493-5497.
- Lee SG, Joo Y, Kim B, Chung S, Kim HL, Lee I, Choi B, Kim C, Song K. (2005). Association of Ala72Ser polymorphism with COMT enzyme activity and the risk of schizophrenia in Koreans. *Human Genetics* 116: 319-328.
- Lunn RM, Bell DA, Mohler JL, Taylor JA. (1999). Prostate cancer risk and polymorphism in 17-hydroxylase (CYP17) and steroid reductase (SRD5A2). *Carcinogenesis* 20: 1727-1731.
- Makridakis NM, Ross RK, Pike MC, Crocitto LE, Kolonel LN, Pearce CL, Henderson BE, Reichardt JK. (1999). Association of mis-sense substitution in SRD5A2 gene with prostate cancer in African-American and Hispanic men in Los Angeles, USA. *Lancet* 354: 975-978.
- Neuberger J, Forman D, Doll R, Williams R. (1986). Oral contraceptives and hepatocellular carcinoma. *British Medical Journal* 292: 1355-1357.
- Newbold RR, Hanson RB, Jefferson WN, Bullock BC, Haseman J, McLachlan JA. (2000). Proliferative lesions and reproductive tract tumors in male descendants of mice exposed developmentally to diethylstilbestrol. *Carcinogenesis* 21: 1355-1363.
- Reichardt JK, Makridakis N, Henderson BE, Yu MC, Pike MC, Ross RK. (1995). Genetic variability of the human SRD5A2 gene: implications for prostate cancer risk. *Cancer Research* 55: 3973-3975.
- Rossi L, Leveri M, Gritti C, De Silvestri A, Zavaglia C, Sonzogni L, Silvestri L, Civardi E, Mondelli MU, Silini EM. (2003). Genetic polymorphisms of steroid hormone metabolizing enzymes and risk of liver cancer in hepatitis C- infected patients. *Journal of Hepatology* 39: 564-570.
- Sasaki M, Tanaka Y, Kaneuchi M, Sakuragi N, Dahiya R. (2003). CYP1B1 gene polymorphisms have higher risk for endometrial cancer, and positive correlations with estrogen receptor alpha and estrogen receptor beta expressions. *Cancer Research* 63: 3913-3918.

- Shastry BS. (2002). SNP alleles in human disease and evolution. *Journal of Human Genetics* 47: 561–566.
- Shen Y, Li DK, Wu J, Zhang Z, Gao E. (2006). Joint effects of the CYP1A1 MspI, ERalpha PvuII, and ERalpha XbaI polymorphisms on the risk of breast cancer: results from a population-based case-control study in Shanghai, China. *Cancer Epidemiology, Biomarkers & Prevention* 15: 342–347.
- Tanaka Y, Sasaki M, Shiina H, Tokizane T, Deguchi M, Hirata H, Hinoda Y, Okayama N, Suehiro Y, Urakami S, Kawakami T, Kaneuchi M, Pookot D, Igawa M, Okuyama A, Ishii N, Dahiya R. (2006). Catechol-O-methyltransferase gene polymorphisms in benign prostatic hyperplasia and sporadic prostate cancer. *Cancer Epidemiology, Biomarkers & Prevention* 15: 238–244.
- Thompson PA, Shields PG, Freudenheim JL, Stone A, Vena JE, Marshall JR, Graham S, Laughlin R, Nemoto T, Kadlubar FF, Ambrosone CB. (1998). Genetic polymorphism in catechol-O-methyltransferase, menopausal status, and breast cancer risk. *Cancer Research* 58: 2107–2110.
- Tsukuma H, Hiyama T, Tanaka S, Nakao M, Yabuuchi T, Kitamura T, Nakanishi K, Fujimoto I, Inoue A, Yamazaki H. (1993). Risk factors for Hepatocellular carcinoma among patients with chronic liver disease. *The New England Journal of Medicine* 328: 1797-1801.
- Yamada Y, Watanabe M, Murata M, Yamanaka M, Kubota Y, Ito H, Katoh T, Kawamura J, Yatani R, Shiraishi T. (2001). Impact of genetic polymorphisms of 17-hydroxylase cytochrome P-450 (CYP17) and steroid 5a-reductase type II (SRD5A2) genes on prostate-cancer risk among the Japanese population. *International Journal of Cancer* 92: 683–686.
- Yin PH, Lee HC, Chau GY, Liu TY, Liu HC, Lui WY, Chi CW. (2004). Polymorphisms of estrogen metabolizing genes and risk of hepatocellular carcinoma in Taiwan females. *Cancer Letter* 212: 195–201.
- Yu MW, Yang YC, Yang SY, Cheng SW, Liaw YF, Lin SM, Chen CJ. (2001). Hormonal markers and hepatitis B virus-related hepatocellular carcinoma risk: a nested case-control study among men. *Journal of the National Cancer Institute* 93: 1644–1651.
- Yuan X, Zhou G, Zhai Y, Xie W, Cui Y, Cao J, Zhi L, Zhang H, Yang H, Qiu W, Peng Y, Zhang X, Yu L, Xia X, He F. (2008). Lack of association between the functional polymorphisms in the estrogen-metabolizing genes and the risk for hepatocellular carcinoma. *Cancer Epidemiology, Biomarkers & Prevention* 17(12): 3621-3627.
- Zhang L, Gu L, Qian B, Hao X, Zhang W, Wei Q, Chen K. (2008). Association of genetic polymorphisms of ER-alpha and the estradiol-synthesizing enzyme genes CYP17 and CYP19 with breast cancer risk in Chinese women. *Breast Cancer Research and Treatment* 114(2): 327-338.

ARABIC SUMMERY

العلاقة بين تعدد الأشكال الجينية ل COMT Val158Met وخطر تطور سرطان الخلايا الكبدية في المرضى المصريين المصابين بفيروس الالتهاب الكبدي الوبائي (سي)

إيمان لبيب^{١,٢*}، هانى محمد ابراهيم^٣، مي الدالى^{١,٢}، هبة سامى غانم^١، ابراهيم عبد الحميد العليمي^٤، محمد عبد الحميد^٤

١-قسم الباثولوجيا الاكلينيكية- معهد الكبد القومى - جامعة المنوفية- شبين الكوم- مصر

٢-قسم الحيوان- كلية العلوم - جامعة المنوفية- شبين الكوم- مصر

٣-وحدة الكائنات المعدية الخاصة- مركز الملك فهد للبحوث الطبية- جامعة الملك عبد العزيز- جدة- المملكة العربية السعودية

٤- قسم الميكروبيولوجي - كلية الطب - جامعة المنيا

بعد التهاب الكبد الفيروسي المزمن من أهم عوامل الخطير التي تؤدى إلى حدوث سرطان الخلايا الكبدية في مصر. يلعب جين الـ COMT دوراً أساسياً في إصلاح الحمض النووي والتسرطن الناجم عن الاستروجين. وقد حفقت العديد من الدراسات الوباينية الحديثة العلاقة بين تعدد الأشكال لجين COMT Val158Met وخطر السرطان.

في هذه الدراسة ، قمنا بالتحري عن الارتباط بين تعدد الأشكال لجين COMT Val158Met A/G وخطر تطور سرطان الخلايا الكبدية بين المرضى المصريين المصابين بفيروس الالتهاب الكبدي الوبائي (سي). أجريت هذه الدراسة على مجموعتين. وشملت مجموعة المرضى المصابين بسرطان الخلايا الكبدية ومجموعة الأصحاء كل مجموعة تشمل ١٠٠ فرد. وقد تمت دراسة تعدد الأشكال الفردية (SNPs) لجين COMT باستخدام تقاعل البلمرة التسلسلي Real-time PCR.

نتائجنا تشير إلى أن جين الـ COMT لا يرتبط مع خطر الإصابة بسرطان الخلايا الكبدية بين المرضى المصريين بفيروس الالتهاب الكبدي الوبائي (سي). تم الكشف عن وجود علاقة بين جين الـ COMT في الفتنة العمرية أكثر من ٥٠ سنة مقارنة مع المرضى الأصغر سنا. تم الكشف عن عدم وجود علاقة بين جين الـ COMT على مستوى الجنس ، حالة الكبد (نليف الكبد) ، و تحاليل البيوكيميائية و الدموية المختلفة محل الدراسة.

في الختام لا يوجد علاقة بين تعدد الأشكال الجينية ل COMT Val158Met وخطر تطور سرطان الخلايا الكبدية في المرضى المصريين المصابين بفيروس الالتهاب الكبدي الوبائي (سي).