



Interaction of Red Cell Indices and Blood Group in Diabetic Patients above 50 Years of Age

Moore-Igwe Beatrice W. ^{a*}, Ken-Ezihuo Stella U. ^a
and Baribefe Banavule Daniel Koate ^a

^a Department of Medical Laboratory Science, Faculty of Science, Rivers State University, Nkpolu-Oroworukwo, P.M.B. 5080, Port-Harcourt, Rivers State, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IBRR/2023/v14i4322

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/105460>

Original Research Article

Received: 01/07/2023

Accepted: 03/09/2023

Published: 29/09/2023

ABSTRACT

Introduction: A group of metabolic diseases with a hyperglycemic phenotype known as diabetes mellitus impact both industrialized and developing nations. It is brought on by deviations in insulin synthesis, action, or both.

Research Objective: This study seeks to associate the interaction of red cell indices and blood group in diabetic patients above 50 years of age. This is a case control study in which a convenient sample size of 100 individuals were recruited from a consenting population at a private hospital in Port Harcourt, Rivers State, 50 of whom were diabetic, and the remaining 50 were not.

Methods: Using the automated 3-part differential hematology analyzer Sysmex KX-21N, a Full blood count (FBC) and differential was obtained using (Sysmex Corporation Kobe, Japan) Standardization, instrument calibration, and analysis of sample was achieved in line with accordance with the manufacturer's manual to determine the red cell indices and grouping of cells using the tile approach to determine the ABO Blood Groups. Blood samples were taken from the

*Corresponding author: E-mail: beatrice.moore-igwe1@ust.edu.ng;

veins using a vacutainer tube and needle according to Chesbrough's instructions, and then put in EDTA anticoagulated bottles with clear labelling. These samples were transported right away using the triple packing procedure to the lab and stored until testing at 4°C. Inferential statistics utilizing the students' statistical t-test were employed to compare the results.

Results: Blood group O, with a total of 25 (50%), had the highest prevalence of diabetes mellitus, followed by blood group A with a total of 12 subjects (24%) and blood group B with a total of 8 subjects (16%) and blood group AB came in last with a total of 5 participants (10%), respectively. In this study, blood group O predominated over blood groups A and B (47% against 25% versus 17%, respectively). The least common blood type in this study was AB (11%). When the Packed Cell Volume, Haemoglobin, Red Blood Cell, Mean Cell volume, Mean Cell Haemoglobin, Mean Cell Haemoglobin Concentration, and Red cell Distribution Width were examined between diabetic and non-diabetic participants, no statistically significant ($p < 0.05$) difference was found.

Conclusion: The findings of this study suggest that blood types O and AB are more prevalent in diabetics. We can deduce that a person's blood type may be a risk factor for developing this condition and may be useful in diagnosing it.

Keywords: Interaction; diabetes; red cell indices; ABO blood groups.

1. INTRODUCTION

Both industrialized and developing nations experience diabetes mellitus [1], a hyperglycemic phenotypic group of metabolic diseases, brought on by abnormalities in insulin synthesis, action, or both. The complex interaction between inherited and environmental factors causes these errors (Sarwar, 2014). In 2013, there were 382 million cases of diabetes globally, up from an estimated 30 million cases in 1985, according to the World Health Organization (WHO).

The World Health Organization (WHO) describes diabetes mellitus as a chronic metabolic disorder characterized by elevated blood glucose levels that, over time, have an impact on the heart, blood vessels, kidneys, eyes, and nerves [1]. 1.7% of the people in Nigeria and the rest of the world suffer from diabetes mellitus, which is a serious concern [2]. According to estimates made by Wild et al. [3], there would be 366,212 million diabetics worldwide by the year 2030. In Rivers State, the prevalence of diabetes mellitus has climbed to roughly 42.7%. However, according to the International Diabetes Federation (IDF), by 2045, adults aged 40 and older are expected to have a diabetes rate of about 10.9%, up roughly 9.3% from 2019. [4].

Families pass down blood type antigens, which are crucial for comprehending genetics, inheritance patterns, disease susceptibility, and the security of blood transfusions. Blood group antigens have been connected to a number of disorders, both in their absence and presence. Due to the fact that diabetes mellitus and blood type share the same genetic and immunological foundations [5]. The relationship between the

ABO blood group and disease risk has recently become the subject of extensive research.

Thankfully, new research from authors like Anifowoshe et al. [6] and Oladele et al. [7] indicates that ABO blood types affect disease susceptibility, which can be used as a management strategy. Studies by Qureshi and Bhatti [8] and Bener and Yousafzai [9], among others, also demonstrate that ABO blood groups affect diabetes susceptibility. ABO blood groups are hereditary and are controlled by a gene on chromosome nine known as the ABO gene or immunoglobulin [10]. Thus, some cases of diabetes may be genetic [11,12].

Additionally, there is evidence supporting the notion that demographic and environmental factors have an impact on illness susceptibility [13]. Shittu et al. [14] advised that high prevalence of diabetes and pre-diabetes among women and those younger than 61 in Oyo State, Nigeria, is noteworthy. In all, these revealed that the association of ABO blood groups and red cell indices can offer basic information for controlling a person's health as well as figure out how diseases are distributed among diabetics.

Consequently, it is necessary to ascertain the distribution of the ABO blood group and among diabetics in a given population so as to effectively employ ABO blood types as well as red cell indices to anticipate and manage a disease. In Port Harcourt, Rivers State, Nigeria, as in other underdeveloped countries, there are few such studies among the diabetic community. Hence, this study aims at ascertaining the relationship between ABO blood groups and red cell indices in diabetics visiting a private hospital in Port Harcourt Rivers State, Nigeria.

RBC parameters are part of the Complete Blood Cell Count (CBC), a panel of analytical assays typically used to distinguish between various kinds of anemia [15]. Red Blood Cell Count (RBC), Hematocrit (HCT), Hemoglobin (HGB), Mean Corpuscular Volume (MCV), Mean Corpuscular Hemoglobin (MCH), Mean Corpuscular Hemoglobin Concentration (MCHC), and Red cell Distribution Width (RDW) are RBC parameters [16]. Continuously exposing RBCs to hyperglycemia causes hemoglobin protein to be permanently glycated, which alters the hemoglobin molecule's structural and functional properties [17]. In addition to changing the mechanical properties and internal viscosity of RBCs, hyperglycemia also promotes aggregation and osmotic fragility. It also causes proteins to become glycated. The structure and hemodynamic properties of the erythrocyte are affected by these processes [18,19]. These changes could be reflected in any or all of the red blood cell analytical measures, such as RBC count, HGB, HCT, MCV, MCH, and RDW.

A measurement of the size diversity of circulating erythrocytes is the red cell distribution width (RDW). In the differential diagnosis of macrocytic anemia, it is mostly utilized as an auxiliary index. A high RDW is linked to anemia caused by iron deficiency. According to a study, whether their RDW is higher or even within the normal reference range, middle-aged and older persons have a noticeably elevated risk of cardiovascular disease (CVD) events. [20,21,22,23]

2. MATERIALS AND METHODS

2.1 Study Design

Patients who sought treatment at a private hospital participated in this cross-sectional study.

2.2 Study Area

At Port Harcourt, the state capital of Rivers State, a private hospital hosted this facility-based cross-sectional study. With a population of 1,865,000 and a location along the Bonny River, Port Harcourt is a metropolis in Nigeria's South-South geopolitical zone that is sometimes frequently referred to as the Niger Delta.

2.3 Study Population

In total, 100 male and female participants who were 50 years of age or older were included in this study. The other 50 test subjects, who

functioned as controls and did not have diabetes mellitus, whereas the 50 others were diabetic patients.

2.4 Collection of Blood Samples, Storage and Transportation

Following the directions of Chesbrough's [24] blood samples were drawn from the veins using a vacutainer tube and needle and placed in EDTA anticoagulated bottles with clear labels. Following their immediate transfer to the lab, these samples were kept at 4°C until testing using a triple packing method.

2.5 Methodology

2.5.1 ABO blood grouping

2.5.1.1 Methods: grouping cells using tiles

A white, flawless tile was divided into sections. In compartments 1, 2, and 3, one drop of anti-A serum, one drop each of anti-B and anti-AB serums, and one drop of the patient's serum was added (auto control). A single drop of 20% patient red blood cells that had been properly blended with an applicator stick was injected into each of the 4 compartments. After gently moving the tile, it was permitted to stand for three to five minutes. Within five minutes, the agglutination on the tile was studied macroscopically and microscopically, with the results being read and recorded.

3. RESULTS

3.1 Demographic Details of Study Participants

The study included 100 individuals in total. Those with diabetes were 50, subjects without diabetes were 50. Males without diabetes were 25, while those with diabetes were 27. Females without diabetes were 28, while those with diabetes were 23. Details are shown in Table.1.

3.2 Frequency Distribution of on ABO Blood Group in Diabetic and Non Diabetic Subjects

Table.2 revealed that blood group O had the greatest incidence of diabetes mellitus 25(50), followed in descending order by blood groups A 12(24), B 8(16), and AB 5 (10).

Table 1. Demographic details of study participants

Parameters	Number
Total Number of Subjects	100
Total Number of Diabetic Subjects	50
Total Number of Non-Diabetic Subjects	50
Total Number of Diabetic Males	27
Total Number of Diabetic Females	23
Total Number of Diabetics with Blood Group A	25
Total Number of Diabetics with Blood Group B	17
Total Number of Diabetics with Blood Group AB	11
Total Number of Diabetics with Blood Group O	47

Table 2. Frequency distribution of on ABO blood group in diabetic and non diabetic subjects

Blood Group	Diabetic		Non-Diabetic		Total Frequency	Total Percent (%)
	Frequency	Percentage (%)	Frequency	Percentage (%)		
A	12	12	13	13	25	25
B	8	8	9	9	17	17
AB	5	5	6	6	11	11
O	25	25	22	22	47	47

4. DISCUSSION

Numerous researchers have attempted to find evidence of a connection between diabetics and specific ABO groupings. Variable, inconsistent, and regionally distinct outcomes have been obtained. Although some researchers have shown a link between blood types and diabetes, other studies have not been able to confirm this [25,9,26,27,28].

This study's findings showed that blood group O, of which there was a total of 25 (50%), had the highest prevalence of diabetes mellitus. Blood group AB came in last with a total of 5 participants (10%), followed by blood group A with a total of 12 subjects (24%) and blood group B with a total of 8 subjects (16%) respectively. In this study, blood group O predominated over blood groups A and B (47% against 25% versus 17%, respectively). The least common blood type in this study was AB (11%). The findings of this study are at odds with a study by Waseem AG et al. [29] that discovered diabetics had a higher likelihood of having blood type AB than non-diabetics. Blood types B (32%) and O (18%) showed the strongest and weakest links, respectively, with diabetes mellitus, according to a study by Zabida et al. (2021). There is a lot of evidence connecting certain blood types, particularly the A, AB, and Rh-positive blood groups, to the condition of diabetes mellitus, according to Sidhu et al. inquiry on the subject in 2018.

In their investigation on the relationship between ABO blood groups and diabetes mellitus, Muhammad et al. (2010) discovered a negative correlation between ABO blood groups A and O and diabetes, with A and O group having decreased risks of diabetes.

When ABO blood groups are compared to diabetic populations around the world, different blood groups are distributed in different ways.

This study found that blood types O and A were the most prevalent among diabetics, and these results were in line with those of studies conducted in Japan and other Sub-Saharan African nations. Other nations including Nigeria, Algeria, Qatar, and Iraq have the highest prevalence of blood group O. While the B blood group was the most prevalent blood group in nations like Malaysia, Italy, and Trinidad [30].

Racial and geographic differences may have an impact on how the disease is expressed genetically could be one explanation for the contradictory findings about the link between ABO blood types and diabetes.

5. CONCLUSION

According to the results of our study, diabetics are more likely to have the blood types O and AB. We can infer that blood type might be a risk factor and might be helpful in determining the condition.

CONSENT AND ETHICAL APPROVAL

After receiving approval from the Department of Medical Laboratory Science at Rivers State University, Port Harcourt, informed consent was obtained from patients who appeared to be in good health prior to enrollment.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. WHO. Expert Committee on the Selection, Use of Essential Medicines, World Health Organization. The Selection and Use of Essential Medicines: Report of the WHO Expert Committee, 2013 (including the 18th WHO Model List of Essential Medicines and the 4th WHO Model List of Essential Medicines for Children). World Health Organization; 2014.
2. Wokoma FS. Diabetes and hypertension in Africa, an overview. *Diabetes international*. 2002;12(12):36-40.
3. Wild S, Roglic G, Green A, Sicree R, King H. Global prevalence of diabetes: estimates for the year 2000 and projections for 2030. *Diabetes care*. 2004 May 1;27(5):1047-53.
4. Saeedi P, Petersohn I, Salpea P, Malanda B, Karuranga S, Unwin N, Colagiuri S, Guariguata L, Motala AA, Ogurtsova K, Shaw JE. Global and regional diabetes prevalence estimates for 2019 and projections for 2030 and 2045: Results from the International Diabetes Federation Diabetes Atlas. *Diabetes research and clinical practice*. 2019 Nov 1;157:107843.
5. Desisto CL, Kim SY, Sharma AJ. Peer reviewed: Prevalence estimates of gestational diabetes mellitus in the United States, pregnancy risk assessment monitoring system (prams), 2007–2010. *The Lancet*. 2014;11(2):541–549.
6. Anifowoshe AT, Owolodun OB, Akinseye KM, Iyioloa OA, Oyeyemi BF. Gene frequencies of ABO and Rh blood groups in Nigeria: a review. *Egypt J Med Human Genetics*. 2017;18(3):205–210. DOI:<https://doi.org/10.1016/j.ejmhg.2016.10.004>
7. Oladele EO, Yahaya TO, Adewunmi OO, David B, Oladipo AJ. Distribution of ABO/rhesus blood groups among hepatitis B virus (HBV) positive patients in Lagos, South-Western, Nigeria. *FUDMA J Sci*. 2020;4(3):196–200.
8. Qureshi MA, Bhatti R. Frequency of ABO blood groups among the diabetes mellitus type 2 patients. *J Coll Phys Surg Pak*. 2003;13(8):453–455.
9. Bener A, Yousafzai M. “The distribution of the ABO blood groups among the diabetes mellitus patients,” *Nigerian journal of clinical practice*. 2014;17(5):565–568.
10. Zahid H, Yahyaoui A, Uwingabiye J, El Khazraji A, El Hadeif R, Messaoudi N. Phenotypic and allelic distribution of ABO and Rh-D blood Groups in Blood Transfusion Center of Avicenna Military Hospital, Marrakech, Morocco. *Int J Med Health Res*. 2016;2(1):1–9.
11. Abdel Ghafar MT, Shalaby KH, Okda HI, Rizk, FH. Association of ABCA1 (C69T) gene polymorphism with dyslipidemia and type 2 diabetes among the Egyptian population. *Meta Gene*. 2020;25:100714. DOI:<https://doi.org/10.1016/j.mgene.2020.100714>
12. Yahaya T, Oladele E, Fatodu I, Abdulazeez A, Yeldu Y. Concentrations and health risk evaluation of heavy metals and microorganisms in groundwater in Lagos, Southwest Nigeria. *Journal of Advances in Environmental Health Research*. 2020;8(3): 234–242. DOI:<https://doi.org/10.22102/jaehr.2020.245629.1183>
13. Yahaya T, Obaroh IO, Oladele EO. The roles of environmental pollutants in the pathogenesis and prevalence of diabetes: a review. *J Appl Sci Environ Manag*. 2017; 21(1):5–8. DOI:<https://doi.org/10.4314/jasem.v21i1.1>
14. Shittu RO, Kasali FO, Biliaminu SA, Odeigah LO, Sule AG, Musah Y. Prevalence of diabetes and pre-diabetes in Oke-Ogun Region of Oyo State, Nigeria. *J Med Rese Health Educ*. 2017;1:1.
15. Doig K, Zhang B. A Methodical Approach to Interpreting the Red Blood Cell Parameters of the Complete Blood Count. *Am Soc Clin Lab Sci*. 2017;30(3):173–85.
16. Vinupritha P, Hariharan M, Kathirvelu D, Chinnadurai S. Estimation of HbA1C using CBC in the diagnosis of diabetes. *Asian J Pharm Clin Res*. 2017;10(9):214–8.
17. Turpin C, Catan A, Guerin-Dubourg A, Debussche X, Bravo S, Álvarez EI. Enhanced oxidative stress and damage in glycated erythrocytes. *PLoS One*. 2020.

18. Juhi A. Comparative study of erythrocyte fragility in diabetes mellitus and nondiabetes Mellitus. *Int J Med Res Heal Sci.* 2015;4(1):183–5.
19. Li Q, Yang LZ. Hemoglobin A1c level higher than 9.05% causes a significant impairment of erythrocyte deformability in DM. *Acta Endocrinol.* 2018;14(1):66–75.
20. Anderson JL, Ronnow BS, Horne BD, Carlquist JF, May HT. Usefulness of a complete blood count-derived risk score to predict incident mortality in patients with suspected cardiovascular disease. *Am J Cardiol.* 2007;99:169-174.
21. Ani C, Ovbiagele B. Elevated red blood cell distribution width predicts mortality in persons with known stroke. *J Neurol Sci.* 2009;277:103-108.
22. Patel KV, Ferrucci L, Ershler WB, Longo DL, Guralnik JM. Red blood cell distribution width and the risk of death in middle-aged and older adults. *Arch Intern Med.* 2009;169:515-523.
23. Perlstein TS, Weuve J, Pfeffer MA, Beckman JA. Red blood cell distribution width and mortality risk in a community-based prospective cohort. *Arch Intern Med.* 2007;169: 588-594.
24. Cheesbrough M. *District Laboratory Practice in Tropical Countries.* 2nd Edition, Cambridge University Press, Cambridge, United Kingdom; 2010.
25. Sukalingam K, Ganesan K. “Rh blood groups associated with risk to obesity and diabetes mellitus: a report on Punjabi population in Selangor, Malaysia,” *International Journal of Integrative Medical Sciences.* 2015;2(4): 105–109.
26. Dali SM, Aour M, Belmokhtar, AF, Belmokhtar R, Boazza F. “The relationship between ABO/Rh blood groups and type 2 diabetes mellitus in Maghnia, western Algeria,” *South African Family Practice.* 2014;53(6):568–572.
27. Moinzadeh F, Najafabady GM, Toghiani A. “Type 2 diabetes mellitus and ABO/Rh blood groups,” *Journal of research in medical sciences: the official journal of Isfahan University of Medical Sciences.* 2014;19(4):382.
28. Saeedi P, Petersohn I, Salpea P, Malanda B, Karuranga S, Unwin N, Colagiuri S, Guariguata L, Motala AA, Ogurtsova K, Shaw JE, Bright D, Williams R, & IDF Diabetes Atlas Committee. Global and regional diabetes prevalence estimates for 2019 and projections for 2030 and 2045: Results from the International Diabetes Federation Diabetes Atlas, 9th edition. *Diabetes research and clinical practice.* 2019;157:107843.
29. Waseem AG, Iqbal M, Khan OA, Tahir M. Association of diabetes mellitus with ABO and Rh blood groups. *Ann Pak Inst Med Sci.* 2012;8(2):134-6.
30. Jayawardena R, Ranasinghe P, Galappatthy P, Malkanthi RL, Constantine GR, Katulanda P. Effects of zinc supplementation on diabetes mellitus: a systematic review and meta-analysis. *Diabetology & Metabolic Syndrome.* 2012 Dec;4(1):1-2.

© 2023 Beatrice et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/105460>