



Quality of Tea Soil Induced by Cultivation Period

**Sayma Muzib^a, M. Hasinur Rahman^a, Hafiz Ashraful Haque^b,
Ferdouse Zaman Tanu^{a*} and Azizul Hakim^c**

^a Department of Soil and Environmental Sciences, University of Barishal, Barishal – 8254, Bangladesh.

^b Department of Coastal Studies and Disaster Management, University of Barishal, Barishal – 8254, Bangladesh.

^c Department of Soil Science, University of Chittagong, Chattogram - 4331, Bangladesh.

Authors' contributions

This work was carried out in collaboration among all authors. Author SM completed the sampling and laboratory works, managed the literature searches and wrote the protocol. Authors MHR and HAH designed the study, managed all the analyses and guided the overall study. Author FZT wrote the draft of the manuscript, presented data through tables and graphs, and performed all the statistical analysis and author AH guided the manuscript preparation, edited and corrected the draft. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/ASRJ/2023/v7i3134

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/105494>

Original Research Article

Received: 25/06/2023

Accepted: 30/08/2023

Published: 09/09/2023

ABSTRACT

Tea is Bangladesh's second-highest agricultural export earner, and the nation is rated 15th among all tea-exporting nations. To develop and support the tea business in Bangladesh, it is crucial to comprehend the current nutrient status of tea soils. The purpose of the study was to examine the inherent physical and chemical characteristics of tea soils in relation to the effects of soil depth, topography and cultivation periods. The results showed the significant effects of soil depths, topography, and cultivation periods on the measured physical and chemical properties, including texture, bulk density, organic matter (OM), organic carbon (OC), pH, and cation exchange capacity

*Corresponding author: E-mail: fztanu@bu.ac.bd;

(CEC). The available and total contents of major nutrients such as nitrogen (N), phosphorus (P), and potassium (K) were also significant ($P < 0.001$). The contents of P and K were observed to be lower than the critical values ideal for tea cultivation. However, the results indicated that the soils kept for nutrient restoration for years showed a nutrient status close to the ideal value of tea cultivation.

Keywords: Nutrient status; tea soils; soil depth; topography; cultivation period.

1. INTRODUCTION

As a traditional beverage, tea (*Camellia sinensis* L.) is the second most widely consumed health drink worldwide after water [1,2]. It primarily originated in south-eastern China but is now being cultivated in many countries across tropical and subtropical regions. Among the 50 tea-producing countries, some of the major countries are China, India, Kenya, Sri Lanka, Vietnam, Turkey, Indonesia, and Iran [3]. In Bangladesh, tea is cultivated in three ecological zones: Surma Valley in greater Sylhet, Halda Valley in Chattogram, and Karatoa Valley in Panchagarh [4]. Tea is the second-highest agricultural export-earning product in Bangladesh. Bangladesh ranks 12th in the list of the highest tea producing countries and 15th among the highest tea-exporting countries [5]. However, a knowledge gap in proper tea soil management still limits the success of the tea industry in Bangladesh.

Long-term cultivation of tea crops leads to a decline in soil quality [6]. Maintaining nutrient status of soils through proper management plays a vital role in tea cultivation. The harvestable portion of tea is the succulent shoot, which contains the largest percentage of nutrients in the plant [7]. Therefore, tea plant requires a high nutrient supply for commercial production. The traditional approach of tea cultivation using chemical fertilizers and pesticides causes a significant soil degradation [8] by the reduction of soil organic matter (OM), nitrogen (N) and (P) loss due to erosion, fixation of P in acidic pH, and reduction of nutrients by soil microorganisms. A combination of several physiochemical properties of tea soils, management practices, and amendments can affect tea-yield. Tea-yield per hectare in Bangladesh is low compared to other tea growing countries of the world. The causes of low tea production are determined by low soil pH, low nutrient status and anthropogenic disturbances [9]. To ensure sustainable tea production [10], nutrient management of tea soils is important particularly for improving soil health. In general, N, P and K contents and other nutrient status of 90% tea cultivated areas are in

medium-low [11]. Tea-farmers in Bangladesh use NPK fertilizer without any proper knowledge about existing nutrient status of the soils. Tea production may decrease with unplanned and imperfect fertilizing program. In Bangladesh, tea industry expansion is therefore threatened by knowledge gap and inadequate management practices.

Currently in Bangladesh, ecological and/or sustainable (integrated and natural resource-based) approaches through the usage of various organic additives (compost, vermicompost, biochar, etc.) in combination with inorganic fertilizers are increasingly important to replace traditional (chemical fertilizer-based) approaches in tea cultivation [12,13]. Still, nutrient loss due to erosion and leaching in tea soils is significant. The determination of the current nutrient status for the adaptation of the effective fertilizing program and other tea soil management practice to increase tea production and expansion of the tea industry in Bangladesh is essential.

Therefore, the objectives of the current study were to determine the nutrient status and the variation in properties of tea soils in Bangladesh in relation to soil depth, topography, and cultivation period. The findings will provide a scientific groundwork for tea soil management and proper fertilization approaches to improving Bangladesh's tea soils. The outcomes will ultimately assist tea farmers and promote the development of the tea industry in Bangladesh.

2. MATERIALS AND METHODS

2.1 Soil Sampling Sites

Three hills in a sub-station of Bangladesh Tea Research Institute at Udulia of Fatikchari upazila in Chattagram district of Bangladesh were selected as the soil sampling sites. The hills were selected based on different tea cultivation periods. They were categorized as: i) Hill 1 (10 years cultivated hill): under cultivation from 1972 to 1982 and tea leaf collection has already been stopped for nutrient restoration; ii) Hill 2 (2 years

cultivated hill): under cultivation from 2008 to 2010 and tea leaf collection is ongoing; iii) Hill 3 (1 year cultivated hill): cultivation started at 2018 and tea leaf collection has not been started until 2019.

The average yield of made tea in this sub-station is near about 1500 Kg/ha. In Fatikchari, the wet season is hot, oppressive, and overcast and the dry season is warm, humid, and mostly clear. The most common form of precipitation throughout the year is rain. Fatikchari experiences extreme seasonal variation in monthly rainfall and humidity. The topography within 2 miles of Fatikchari is essentially flat with a maximum elevation change of 49 feet and an average elevation above sea level is 41 feet, and within 10 miles is essentially flat (1,033 feet).

2.2 Collection of Soils

A total of 27 representative soil samples with 3 replications were collected from each of 3 hills at 3 different topographic positions (top, slope, and base) at 3 depths including top-soil (0-15 cm), sub-soil (16-30 cm) and the substratum-soil (31-45 cm) in January 2019 [14]. The location of sampling sites is presented in Fig. 1 and the geographic coordinates of sampling points are shown in Table 1. Both disturbed and undisturbed soils were sampled by using auger and core. Samples were taken to the laboratory of the department of Soil and Environmental Sciences in University of Barishal, Bangladesh.

2.3 Preparation and Analysis of Soil Samples

Soil samples were air-dried, ground and sieved through a 2 mm sieve. Soil samples were analyzed for physical, physico-chemical and chemical parameters. Soil pH was measured using the glass electrode method (HACH instruments, PHC10101). Wet oxidation method of Walkley and Black [15] was followed to determine organic carbon (OC %). Soils were analyzed for both total and available form of N, P and K. Digestion of soil samples were done by a wet-oxidation method using concentrated sulphuric acid. The total N was determined by micro Kjeldahl's method [16]. For total P and total K analysis, wet digestion of soils was done by using nitric acid and perchloric acid [17]. The available N was extracted by 1M KCl and determined by steam distillation method with the Devada's alloy [18]. The available P was extracted by using the method described in Bray

and Kurtz [19] and determined at 880 nm when soil pH was less than 7. The available K was extracted by using neutral 1M Ammonium acetate (NH₄OAc) solution [20] and determined by a flame photometer (Jenway 500701 PFP7 Industrial Flame Photometer). The textural class and CEC were measured by using the hydrometer [21] and ammonium acetate extraction method [22], respectively. Bulk density of the soil sample was determined by Core method [23].

2.4 Statistical Analysis

Values of different soil parameters are reported as mean \pm standard deviation. Measured data were tested for normality of distributions and homogeneity of variances prior to further statistical analysis. Effects of three soil depths (0-15cm, 16-30 cm, and 31-45cm), the topographical variation (three different slope positions), and the three different cultivation periods on soil parameters were compared separately by using the one-way analysis of variance (one-way ANOVA). Statistical analysis was performed by using SPSS V. 23 at the level of $P < 0.05$.

3. RESULTS AND DISCUSSION

3.1 Physical and Physico-Chemical Properties of Tea Soils

Textural classes of the studied soils were sandy loam and sandy clay loam (Table 2), suitable for tea cultivation [24,25]. The textural classes of the soils might affect many other properties of soils including moisture availability, soil temperature, nutrient supply, and accessibility of soil organic matter to microbial decomposition [26]. Bulk densities of studied soils ranged from 1.25 to 1.71 gcm⁻³ (Table 2) with a mean of 1.45 gcm⁻³ in Hill 1, 1.49 gcm⁻³ in Hill 2 and 1.40 gcm⁻³ in Hill 3. The results of the bulk densities showed an increasing trend across the soil depths as well as the topography from top to base position of the hills except some exceptions (Table 2). The results were congruent with the observation by Lee et al. [27]. The increasing trend of bulk density with soil depth might be attributed to the lower clay and organic matter content [28] and compaction of the lower layer soils. Soil quality typically decreases with increasing bulk density [29].

The pH of the studied soils ranged from 4.7 to 5.5, 3.4 –to 4.4 and 4.3 to 5.0 in Hill 1, Hill 2, and



Fig. 1. Location of sampling sites

Hill 3, respectively, whereas the optimum pH range for tea cultivation is 4.5-5.5 [30]. The results showed that Hill 1 was in the best condition regarding pH for tea cultivation among the hills. Being cultivated for 10 years from 1972 to 1982, the Hill 1 is not currently under tea collection and left for quality restoration for several years. In contrast, Hill 2 soils were out of

the favorable pH range for tea cultivation, whereas some of the Hill 3 soils have the favorable pH (Table 2). Hill 2 had been cultivated for 2 years and was under production during soil sampling, whereas Hill 3 had been cultivated for only 1 year and tea collection was yet to be started at the sampling time. It is quite clear that the Hill 2 soil needs proper management for

brining pH in favorable range for tea cultivation. Lime and organic fertilizer amendments can maintain the optimum soil pH for tea cultivation. The conventional tea cultivation approach usually uses chemical fertilizers including ammonium nitrate, ammonium sulphate, urea, calcium-ammonium nitrate, and ammonium chloride. These fertilizers eventually increase soil acidity [31] through the enhancement of microbial decomposition of organic matter resulting in organic acids production in soils [32]. This might be the reason why the pH of the upper soil was slightly lower than that of the lower depth soil.

The CEC ranged from 9.26 to 15.6 meq/100 g in Hill 1, 8.29 to 14.0 meq/100 g in Hill 2 and 6.19 to 12.6 meq/100 g in Hill 3 (Table 2). Average values of CEC were found 12.14, 11.75, and 9.77 meq/100g, respectively in Hill 1, Hill 2, and Hill 3. Hossain et al. [33] reported on a range of CEC of 5.15 to 33.25 meq/100g soil at Satgoan, Baraora and Kurmah tea estates, in Sylhet, Bangladesh. Additionally, CEC ranged from

11.42 to 24.86 meq/100 g soil of Moulvibazar district, Bangladesh [34]. This variation of CEC was accompanied with the amount of clay content, pH, and percentage of organic matter in the investigated soils [35].

3.2 Chemical Properties of Tea Soils

The relative variation of organic matter (OM) and organic carbon (OC) of collected soils of three hills are presented in Fig. 2. The observed values of OC ranged from 0.565 to 1.093%, 0.565 to 1.068%, and 0.641 to 1.858% with average values of 0.861%, 0.761%, and 0.995% in Hill 1, Hill 2, and Hill 3, respectively. Additionally, the OM ranged from 0.97 to 2.01%, 0.71 to 2.07%, and 1.1 to 3.14% with average values of 1.48%, 1.31%, and 1.71% in Hill 1, Hill 2, and Hill 3, respectively. In top positions of Hill 1 and Hill 3, OC contents of upper soils (0 to 15 cm) were around the critical value (1%) for tea cultivation [25], whereas OC contents in other sampling points were much lower.

Table 1. Geographic coordinates of the sampling points in three hills in BTRI sub-station at Fatikchhari upazila, in Chattagram district, Bangladesh

Topography	Sampling point	Coordinates
10 years cultivated hill		
Top	1	22°36'59"N 91°45'31"E
	2	22°36'58"N 91°45'32"E
	3	22°36'57"N 91°45'34"E
Slope	1	22°36'57"N 91°45'33"E
	2	22°36'57"N 91°45'32"E
	3	22°36'57"N 91°45'31"E
Base	1	22°36'56"N 91°45'32"E
	2	22°36'56"N 91°45'31"E
	3	22°36'56"N 91°45'33"E
2 years cultivated hill		
Top	1	22°36'50"N 91°45'37"E
	2	22°36'49"N 91°45'36"E
	3	22°36'49"N 91°45'35"E
Slope	1	22°36'50"N 91°45'38"E
	2	22°36'51"N 91°45'37"E
	3	22°36'50"N 91°45'36"E
Base	1	22°36'50"N 91°45'34"E
	2	22°36'50"N 91°45'35"E
	3	22°36'50"N 91°45'36"E
1 year cultivated hill		
Top	1	22°37'05"N 91°45'29"E
	2	22°37'00"N 91°45'29"E
	3	22°37'06"N 91°45'28"E
Slope	1	22°37'03"N 91°45'29"E
	2	22°37'04"N 91°45'29"E
	3	22°37'06"N 91°45'29"E
Base	1	22°37'05"N 91°45'30"E
	2	22°37'06"N 91°45'30"E
	3	22°37'04"N 91°45'30"E




Fig. 2. Relative variation in organic carbon (OC) and organic matter (OM) in soils of three different depths and topography in three hills under different cultivation periods in a sub-station of Bangladesh Tea Research Institute at Uduia of Fatikchari upazila in Chattagram district of Bangladesh

Additionally, OC contents were higher in upper layer of top, slope and base soils in Hill 3 than those in Hill 1. The amount of plant residues returned to the soil are observed lower in very young and very old tea fields than in the more productive tea gardens [36]. The variation in OM regarding quantity and quality in tea soils depends on the age of plantation, plant residue inputs, soil loss, conditions governing residue decomposition, and related management practices [37]. Both of OC and OM contents were observed to be decreased subsequently with the depth (Fig. 2) as concurrently found in Haorongbam et al. [38]. The higher OM contents in upper soils were presumably due to the accumulation of fallen leaves and microbial residues [39]. Soil conditions necessary for attaining optimum tea yields in the uplands of Vietnam include OM content higher than 2% with a range of pH from 4.0 to 5.5 [40]. Therefore, organic fertilizers can be recommended to maintain the balance of OC and OM in the studied soils [36].

The relative variation in total and available N, P and K contents of collected tea soils are presented in Fig. 3. In Hill 1, the available N varied between 520 and 830 mgKg⁻¹ with an average of 660 mgKg⁻¹, whereas the total N ranged between 963 and 2888 mgKg⁻¹ with an average of 1536 mgKg⁻¹. In Hill 2, the available N ranged from 370 to 830 mgKg⁻¹ with an average of 600 mgKg⁻¹, whereas the total N ranged from 709 to 2835 mgKg⁻¹ with an average of 1523 mgKg⁻¹. Additionally, in Hill 3, the available N ranged from 370 to 1000 mgKg⁻¹ with an average of 620 mgKg⁻¹ and the total N ranged from 709 to 4253 mgKg⁻¹ with an average of 1864 mgKg⁻¹. The available N contents were much higher in all the hill soils

than the minimum level (149 mgKg⁻¹) for tea soils [41]. All the studied soils also have sufficient total N content for tea cultivation. However, the variation in total N was not significantly different and showed no regular pattern in distribution across the depth, topography and cultivation period. According to Ranganathan and Natesan [42], N is required in large quantities for tea cultivation because it is accounted for approximately 4 to 5% of the dry weight of the harvested shoots.

The available P in Hill 1, Hill 2 and Hill 3 ranged from 0.13 to 9.42 mgKg⁻¹, 1.01 to 12.1 mgKg⁻¹, and 0.26 to 4.44 mgKg⁻¹, respectively. The average available P contents in Hill 1, Hill 2, and Hill 3 were 3.11, 4.30 and 1.51 mgKg⁻¹, respectively. Those values are comparable with the results ranging from 1.00 to 4.33 mgKg⁻¹ in Islam et al. [43]. On the other hand, Additionally, the average available K contents were 25.4 mgKg⁻¹ in Hill 1, 79.6 mgKg⁻¹ in Hill 2, and 62 mgKg⁻¹ in Hill 3 ranging from 13.3 to 35.8 mgKg⁻¹, 17.3 to 202.5 mgKg⁻¹, and 37.6 to 118.1 mgKg⁻¹, respectively. The available K of only Hill 1 is comparable to the result ranging from 31.2 to 58.5 mgKg⁻¹ for soils of 0-91 cm depth in Islam et al. [43]. The minimum levels of available P and K were determined as 32 and 110 mgKg⁻¹, respectively in Zhang et al. [41]. Another study showed the critical values of available P and K for tea cultivation were 10 mgKg⁻¹ and 80 mgKg⁻¹, respectively [44]. Therefore, the current study sites are considered P and K deficit regarding available P and K contents [39]. The low contents of nutrients might be accounted for the high rate of assimilation of available forms by microorganisms and vegetations, and/or high rate of fixation by Aluminium (with P) in the prevailing acidic conditions.

Table 2. Physical and physico-chemical properties of tea soils collected from three soil depths and three topographic positions of three hills under different cultivation periods in a sub-station of Bangladesh Tea Research Institute at Udulia, Fatikchari upazila, Chattagram district, Bangladesh

Hill	Topography	Depth (cm)	Sand (%)	Silt (%)	Clay (%)	Textural classes	Bulk density (gcm ⁻¹)	pH	CEC (meq/100g soil)
			Mean ± stdv	Mean ± stdv	Mean ± stdv		Mean ± stdv	Mean ± stdv	Mean ± stdv
10 years cultivated hill	Top	0-15	71.2 ± 3.14	14.3 ± 3.89	14.5 ± 0.81	Sandy Loam	1.30 ± 0.05	4.93 ± 0.08	10.4 ± 1.19
		16-30	67.3 ± 6.80	15.0 ± 5.77	17.8 ± 2.00	Sandy Clay Loam	1.34 ± 0.02	4.68 ± 0.03	15.6 ± 0.79
		31-45	67.4 ± 6.13	9.36 ± 2.64	23.3 ± 3.70	Sandy Clay Loam	1.43 ± 0.05	4.66 ± 0.08	15.1 ± 1.38
	Slope	0-15	73.9 ± 2.04	13.8 ± 0.63	12.4 ± 1.45	Sandy Loam	1.27 ± 0.01	5.00 ± 0.13	9.64 ± 1.38
		16-30	71.7 ± 2.00	13.5 ± 1.48	14.8 ± 1.98	Sandy Loam	1.46 ± 0.05	5.01 ± 0.13	12.1 ± 0.35
		31-45	72.1 ± 2.47	10.7 ± 2.46	17.2 ± 4.08	Sandy Clay Loam	1.51 ± 0.04	4.97 ± 0.12	12.4 ± 0.00
	Base	0-15	65.5 ± 2.41	22.9 ± 2.50	11.6 ± 1.95	Sandy Loam	1.31 ± 0.06	5.49 ± 0.14	9.26 ± 2.11
		16-30	62.6 ± 0.63	22.6 ± 0.85	14.8 ± 1.35	Sandy Loam	1.47 ± 0.02	4.99 ± 0.03	11.0 ± 1.19
		31-45	61.9 ± 1.35	21.2 ± 0.44	16.9 ± 1.92	Sandy Loam	1.56 ± 0.09	4.67 ± 0.12	13.8 ± 0.60
2 years cultivated hill	Top	0-15	66.6 ± 2.60	18.8 ± 1.25	14.7 ± 1.44	Sandy Loam	1.32 ± 0.08	3.76 ± 0.03	10.6 ± 0.68
		16-30	57.2 ± 1.38	20.3 ± 1.81	22.5 ± 0.50	Sandy Loam	1.57 ± 0.11	3.42 ± 0.02	11.2 ± 0.48
		31-45	59.1 ± 4.39	16.3 ± 2.50	24.7 ± 1.91	Sandy Clay Loam	1.48 ± 0.04	3.38 ± 0.04	12.4 ± 0.76
	Slope	0-15	63.2 ± 4.08	19.6 ± 3.11	17.3 ± 1.15	Sandy Loam	1.33 ± 0.03	3.98 ± 0.17	11.2 ± 0.56
		16-30	56.6 ± 4.39	19.8 ± 2.54	21.3 ± 1.88	Sandy Clay Loam	1.35 ± 0.06	4.06 ± 0.24	13.3 ± 0.40
		31-45	58.6 ± 3.15	19.7 ± 2.91	21.8 ± 2.14	Sandy Clay Loam	1.55 ± 0.06	4.14 ± 0.16	14.0 ± 0.40
	Base	0-15	68.1 ± 5.64	15.8 ± 4.39	16.1 ± 1.44	Sandy Loam	1.51 ± 0.04	4.37 ± 0.42	12.8 ± 0.44
		16-30	66.8 ± 1.91	15.0 ± 1.25	18.2 ± 0.81	Sandy Loam	1.64 ± 0.04	4.25 ± 0.03	8.29 ± 0.37
		31-45	69.0 ± 1.80	13.3 ± 2.89	17.7 ± 1.26	Sandy Loam	1.71 ± 0.02	4.28 ± 0.06	12.0 ± 0.40
1 year cultivated hill	Top	0-15	66.9 ± 1.91	24.5 ± 1.25	8.58 ± 0.72	Sandy Loam	1.33 ± 0.03	4.96 ± 0.19	9.86 ± 1.04
		16-30	58.3 ± 3.00	25.6 ± 1.88	16.1 ± 1.91	Sandy Loam	1.30 ± 0.03	4.69 ± 0.05	7.56 ± 0.69
		31-45	57.5 ± 2.14	22.3 ± 1.35	20.2 ± 1.13	Sandy Clay Loam	1.46 ± 0.10	4.50 ± 0.10	10.9 ± 1.19

Hill	Topography	Depth (cm)	Sand (%) Mean ± stdv	Silt (%) Mean ± stdv	Clay (%) Mean ± stdv	Textural classes	Bulk density (gcm ⁻¹) Mean ± stdv	pH Mean ± stdv	CEC (meq/100g soil) Mean ± stdv
	Slope	0-15	61.5 ± 1.32	25.1 ± 1.18	13.4 ± 1.33	Sandy Loam	1.33 ± 0.01	4.62 ± 0.11	8.25 ± 0.00
		16-30	57.3 ± 2.81	25.7 ± 2.70	17.0 ± 1.25	Sandy Loam	1.37 ± 0.04	4.42 ± 0.11	6.19 ± 0.69
		31-45	54.3 ± 2.50	22.3 ± 2.32	23.4 ± 1.01	Sandy Clay Loam	1.51 ± 0.03	4.27 ± 0.04	12.6 ± 1.01
	Base	0-15	64.1 ± 4.58	24.6 ± 4.02	11.3 ± 1.70	Sandy Loam	1.47 ± 0.06	4.58 ± 0.16	13.8 ± 1.38
		16-30	66.7 ± 2.60	18.8 ± 1.25	14.6 ± 3.15	Sandy Loam	1.59 ± 0.06	4.40 ± 0.01	7.13 ± 0.43
		31-45	63.4 ± 3.61	18.3 ± 2.89	18.3 ± 3.31	Sandy Loam	1.69 ± 0.04	4.56 ± 0.13	11.7 ± 0.69




Fig. 3. Relative variation in total NPK (solid colored bars) and available NPK (no-colored bars) in soils of three different depths (0-15, 16-30, and 31-45 cm) and topography (top, slope, and base) in three hills under different cultivation periods (10 years, 2 years and 1 year) in a sub-station of Bangladesh Tea Research Institute at Udulia of Fatikchari upazila in Chattagram district of Bangladesh

Furthermore, the total P in Hill 1 ranged from 52.63 to 210.85 mgKg⁻¹ with an average of 107.05 mgKg⁻¹. In Hill 2, that ranged from 78.95 to 289.45 mgKg⁻¹ with an average of 167.66 mgKg⁻¹. In Hill 3, that ranged from 52 to 157.9 mgKg⁻¹ with an average of 94.64 mgKg⁻¹. Whereas, the total K in Hill 1, Hill 2, Hill 3 ranged from 1801.07 to 3141.55 mgKg⁻¹, 2069.17 to 3945.84 mgKg⁻¹, and 1801 to 3677.75 mgKg⁻¹, respectively. The average total K contents of soils in Hill 1, Hill 2, and Hill 3 were 2416.7, 3161.41, and 2615.29, respectively. Thus, the current study sites are considered much rich in total nutrient contents, however having minimal

available nutrients. Moreover, both the total and the available contents of both P and K significantly varied ($P < 0.05$) across the depth, topography and cultivation periods.

3.3 Tea Soil Quality under the Impacts of Soil Depths, Topography and Cultivation Periods

The results of one-way ANOVA regarding the three factors of soil depth, topography and cultivation periods are presented in Table 3. The results manifested that both the soil depth and the cultivation period had the most significant

($P < 0.05$) influences on all the soil parameters except two exceptions (Table 3). However, the topography showed significant effects only on the soil bulk density and total and available contents of K (Table 3). Though there were no significant effects on total and available N contents of soil by the cultivation period, the soil of Hill 1 kept fallow for years had a higher N content. Moreover, one of the most important soil

parameters the soil pH, which controls the availability of other nutrients, did not varied significantly with depth but varied significantly with the cultivation period. Thus, the tea soil quality in the current study is mostly induced by the cultivation period, which is considered as a substantial factor because the fallow period in Hill 1 soil has helped to restore nutrients and enhanced soil quality for tea cultivation.

Table 3. Results of ANOVA regarding the three factors (soil depth, topography and cultivation period) influencing the soil parameters

Factors influencing soil parameters:		Soil depth			Topography		Cultivation period	
Soil parameters	Comparison	df	F	Sig.	F	Sig.	F	Sig.
Sand (%)	Between Groups	2	4.538	.014*	.905	.409	13.591	.000*
	Within Groups	78						
	Total	80						
Silt (%)	Between Groups	2	2.677	.075	.119	.888	21.867	.000*
	Within Groups	78						
	Total	80						
Clay (%)	Between Groups	2	33.719	.000*	2.833	.065	6.619	.002*
	Within Groups	78						
	Total	80						
Bulk Density (gcm ⁻¹)	Between Groups	2	21.779	.000*	16.204	.000*	3.365	.040*
	Within Groups	78						
	Total	80						
pH	Between Groups	2	1.939	.151	2.386	.099	74.783	.000*
	Within Groups	78						
	Total	80						
CEC (meq/100g soil)	Between Groups	2	9.716	.000*	.268	.766	8.341	.001*
	Within Groups	78						
	Total	80						
OC (%)	Between Groups	2	43.528	.000*	1.543	.220	5.298	.007*
	Within Groups	78						
	Total	80						
OC (%)	Between Groups	2	43.604	.000*	1.541	.221	5.307	.007*
	Within Groups	78						
	Total	80						
Available N (mgKg ⁻¹)	Between Groups	2	3.658	.030*	1.895	.157	1.192	.309
	Within Groups	78						
	Total	80						
Available P (mgKg ⁻¹)	Between Groups	2	7.146	.001*	1.965	.147	8.039	.001*
	Within Groups	78						
	Total	80						
Available K (mgKg ⁻¹)	Between Groups	2	44.072	.000*	5.009	.009*	18.494	.000*
	Within Groups	78						
	Total	80						
Total N (mgKg ⁻¹)	Between Groups	2	6.088	.003*	1.025	.364	1.236	.296
	Within Groups	78						
	Total	80						
Total P (mgKg ⁻¹)	Between Groups	2	9.553	.000*	1.940	.151	13.806	.000*
	Within Groups	78						
	Total	80						
Total K (mgKg ⁻¹)	Between Groups	2	25.763	.000*	16.779	.000*	15.762	.000*
	Within Groups	78						
	Total	80						

The bold values of Sig. with asterisk (*) in the table represent the significant P values

4. CONCLUSION

The soils of three different hills were investigated for the effects of soil depth, topography, and cultivation period on tea soil quality. The investigation showed that the soil nutrient status varied widely across the hills under the effects of the concerned factors, mostly the cultivation period. All the three hills were P and K deficit. Comparatively higher available N content in Hill 1, which was kept fallow, indicated that the restoration period of this hill helped to enhance the soil quality. But the highest content of total N in the surface soil of the base position of Hill 3, where the cultivation process is ongoing, demonstrates the potential use of nitrogen fertilizers. The statistical analysis further manifested the importance of the cultivation period in controlling the soil quality. Differences in soil properties between tea plantation age classes would primarily reflect the impact of cultivation history. Therefore, it is recommended that the use of the hills for tea cultivation be more beneficial after a fallow period. This reuse could increase tea production without the use of commercial fertilizer and will be sustainable for the soil and water environment.

ACKNOWLEDGEMENTS

Authors acknowledge the Ministry of Science and Technology, Bangladesh for allowing NST (National Science and Technology) Fellowship for this work. Authors also acknowledge the department of Soil and Environmental Sciences, and the department of Coastal Studies and Disaster Management of the University of Barishal for supporting this work. Authors are grateful to Ashim Kumar Saha, Senior Scientific Officer (Soil Science), BTRI Sub Station, Fatikchari, Chattagram, and Professor Dr. Shaikh Motasim Billah, Soil, Water & Environment Discipline, Khulna University, for their valuable cooperation.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Zhen YS. Tea: Bioactivity and therapeutic potential. ed. CRC Press, Taylor & Francis, New York, USA; 2002.
2. Ho CT, Lin JK, Shahidi F. Tea and tea products: chemistry and health-promoting properties. CRC press, Taylor & Francis group, Boca Raton, FL, USA; 2008.
3. FAO (Food and Agriculture Organization of The United Nations), FAOSTAT: Available:<http://www.fao.org/faostat/en/#data/QC>; 2014.
4. Mamun MSA. Development of tea science & tea industry in Bangladesh and advances of plant extracts in tea pest management. International Journal of Sustainable Agriculture and Technology. 2011;7(5):40-46.
5. BTB. Annual Report. Bangladesh Tea Board, Chittagong; 2017.
6. Dang VM, Anderson DW. Application of soil taxonomy classification into Northern Vietnam. Vietnamese Soil Sci. 2000;13:15-23.
7. Do NQ, Vo TTN, Vi KT. Results of experiment: Chemical fertilizers (NPK) on tea crop in Phu Ho. In Agricultural research. Agricultural Publishing House, Hanoi. (in Vietnamese). 1980;127-146.
8. Ahsan Q. Maximizing yield of tea in some selected problematic acidic soils through improved management practices. Bangladesh Tea Research Institute (BTRI), Bulletin No. 17, 2011, Center for Agri-Research and Sustainable Environment and Entrepreneurship Development (CASEED), Dhanmondi, Dhaka; 2011.
9. Saha AK, Rahman SH, Hoque S. Requirement of lime in tea growing areas to improve growth and yield of tea in Bangladesh. Tea J. Bangladesh. 2015; 44:10-18.
10. Hamid FS. Yield and quality of tea under varying conditions of soils and nitrogen availability, Pakistan Research Repository. Higher Education Commission Pakistan; 2006.
11. Zicheng Z, Xiaoling HE, Tingxuan LL. Status and evaluation of the soil nutrients in tea plantation. Procedia Environmental Sciences. 2012;12:45–51.
12. BTB. Bangladesh Tea Board. Chittagong. Annual Report; 2009.
13. RTRS. Regional Tea Research Station. Annual Report, 2011, Panchagarh. 2012;7-19.
14. USDA (United States Department of Agriculture). Soil survey manual. soil survey staff, bureau of plant industry, soil and agricultural engineering, United States Department of Agriculture, Washington. 1951;18:205.

15. Walkley A, Black IA. An examination of degtjareff method for determining soil organic matter and a proposed modification of the chromic acid titration method. *Soil Science*. 1934;37:29-38.
16. Jackson ML. *Soil chemical analysis*. 1st ed. Prentice-Hall Inc. Englewood Cliffs. NJ; 1973.
17. Jackson ML. *Soil chemical analysis*. Prentice Hall Inc., Englewood Cliffs N. J., USA. 1962;498.
18. Bremner JM. Inorganic forms of nitrogen. In: Black CA, Evans DD, White JL, Ensminger E, Clark FE, editors. *Methods of Soils Analysis. Part 2. Agronomy No. 9*. American Society of Agronomy, Madison, WI. 1965;1179–1237.
19. Bray RH, Kurtz LT. Determination of total, organic, and available forms of Phosphorus in soils. *Soil Science*. 1945; 59:39-46.
Available:<https://doi.org/10.1097/00010694-194501000-00006>
20. Helmke PA, Sparks DL. Neutron activation analysis. In: Sparks DL, editor, *Methods of Soil Analysis*. Madison, Wisconsin. 1996;551–575.
21. Day PR. Particle fraction and particle size analysis. In: Black CA, editor, *Methods of Soil Analysis*, ASA, Madison. 1965;545-567.
22. Sumner ME, Miller WP. Cation exchange capacity, and exchange coefficients. In: Sparks DL, editor, *Methods of soil analysis. Part 2: Chemical properties*. 3rd ed. ASA, SSSA, CSSA, Madison, WI; 1996.
23. Blake GR, Hartge KH. Bulk density. In: A. Klute, (ed.). *Method of soil analysis. Part 1. Agronomy 9*. ASA, Madison, WI. 1986;363-375
24. Kibria AKMG. Soil management with special reference to young tea, BTRI, Srimangal; 1984.
25. Chowdhury SH, Hassan KA, Shome KM, Rahman MA. Nutrient status of the tea soils of balisera circle, sylhet, pamphlet no.2. notes on soil survey, Bangladesh Tea Research Institute, Srimangal, Sylhet; 1974.
26. Schimel DS, Braswell BH, McKeown R, Ojima DS, Parton WJ, Pulliam W. Climate and nitrogen controls on the geography and timescales of terrestrial biogeochemical cycling. *Global Biogeochemical Cycles*. 1996;10(4):677-692.
27. Lee YL, Osumanu HA, Nik Muhamad Ab M, Jalloh MB. Organic Matter, Carbon and Humic Acids in Rehabilitated and Secondary Forest Soils. *American Journal of Applied Sciences*. 2009;6(5):824-828.
28. Gupta RD, Arora S, Gupta GD, Sumberia NM. Soil physical variability in relation to soil erodibility under different land uses in foothills of siwaliks in N-W India. *Tropical Ecology*. 2010;51:183-197.
29. Harris RF, Karlen DL, Mulla DJ. A conceptual framework for assessment and management of soil quality and soil health. In: Doran JW, Jones AJ, editors, *Methods for Assessing Soil Quality*, SSSA Special Publication 49, Madison. 1996;61-82.
30. Ahmed M. *Tea history: Science and technology*. 1st ed. 2016;129-130.
31. Ma WCI, Brussaard L, Ridder JA. Long-term effects of nitrogenous fertilizers on grassland earthworms (*Oligochaeta: Lumbricidae*): Their relation to soil acidification. *Agriculture Ecosystem and Environment*. 1990;30(2):71-80.
32. Tabatabai MA, Fu MH, Basta NT. Effect of cropping systems on nitrification in soils. *Comm. Soil Sci. and Plant Analysis*. 1992; 23(13-16):1885-1891.
33. Hossain MF, Islam A, Islam W, Rahaman S. An evaluation of methods for determining available phosphorus in some tea soils of Bangladesh. *Tea Journal of Bangladesh*. 1988;24:8-16.
34. Sanaullah AFM, Ahmed MS. Distribution of active acidity, reserve acidity, clay content and cation exchange capacity of some soils of Sathgaon tea estate, Bangladesh. *J. App. Sci. Res*. 2010;6:1627.
35. Bortoluzzi EC, Tessier D, Rheinheimer DS, Julien JL. The cation exchange capacity of a sandy soil in southern Brazil: An estimation of permanent and pH-dependent charges. *European journal of soil science*. 2006;57(3):356-364.
36. Li ZP, Ding DX. *China Tea*. (in Chinese). 1992;14(6):13-15
37. Campbell CA, Zentner RP, Juma NG. Case studies of soil quality in the Canadian prairies: long-term field experiments. In: Gregorich EG, Carter MR, editors, *Soil Quality for Crop Production and Ecosystem Health*. Elsevier Science B.V., Amsterdam, The Netherlands; 1997;315-392.
38. Haorongbam NS, Rout J, Sethi LN. Assessment of soil fertility status in silcoorie tea estate, assam, north east

- india. International Journal of Current Research. 2014;6(12):10851-10854.
39. Wang XJ, Hu XF, Chen HZ. Some biogeochemical characteristics of tea soils. Pedosphere. 1997;7(3):275-280.
40. Do NQ, Le TK. Tea: Production-processing-marketing. Agricultural Publishing House, Hanoi; (in Vietnamese). 2000;53-129
41. Zhang YL, Luo SH, Zeng YH, Pengo PY. Study nutrient scale of sufficiency or deficiency in tea soils in Hunan province and fertilizing recommendation. J. of Tea Soil. 1997;17(2):161-170.
42. Ranganathan V, Natesan S. Potassium nutrition of tea. In: Munson RD, editor, Potassium in Agriculture. ASA-CSSA-SSSA, USA. 1985;981-1015.
43. Islam MR, Rony M, Afroz H, Islam MS, Hoque MM. Phosphorus fractionation in acid soil of Lakkatura Tea garden, Sylhet. Bangladesh. Journal of Soil and Nature. 2014;7(1):1-6.
44. Ahmed MS, Zamir MR, Sanaullah AFM. Organic carbon, nitrogen and phosphorus contents of some tea soils. Pakistan Journal of Analytical & Environmental Chemistry. 2005;6(1):6.

© 2023 Muzib et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/105494>