



Fertility Status of Soil and Nutrients Content of Spring Groundnut (*Arachis hypogaea* L.) as Influenced by Gypsum, Nitrogen and Phosphorus Application

Akashdeep Singh Brar^{a++} and S. S. Manhas^{a*}

^a Department of Agronomy, Punjab Agricultural University, Ludhiana-141001, India.

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2023/v35i193742

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/106028>

Original Research Article

Received: 01/07/2023
Accepted: 05/09/2023
Published: 09/09/2023

ABSTRACT

The field investigation was carried out for two years at the Students' Research Farm, Department of Agronomy, Punjab Agricultural University, Ludhiana during spring season of 2018 and 2019. The experiment was laid out in a split plot design with four levels of gypsum (0, 125, 175 and 225 kg ha⁻¹) in combination with two gypsum application stages (full at sowing and 50 % at sowing + 50 % at flower initiation stage) in the main plot and three levels of nitrogen and phosphorus (15 kg N ha⁻¹ + 20 kg P₂O₅ ha⁻¹, 25 kg N ha⁻¹ + 30 kg P₂O₅ ha⁻¹ and 35 kg N ha⁻¹ + 40 kg P₂O₅ ha⁻¹) in the sub-plot. Application of 225 kg ha⁻¹ gypsum resulted in maximum kernel N, P, K and S concentration and haulm P and S concentration, which was statistically at par with 175 kg ha⁻¹ and 125 kg ha⁻¹ gypsum, while significantly higher than control. Split application of gypsum recorded

⁺⁺ Mc Student;

^{*}Corresponding author: E-mail: manhas30@gmail.com;

non-significant results on concentration of N, P, K, Ca and S in haulm and kernel during both the year. The concentration of N, P, K, Ca and S in haulm and kernel was maximum with application of 35 kg N ha⁻¹ + 40 kg P₂O₅ ha⁻¹, except for K and Ca in kernel which was significantly higher than lower dose. Soil characteristics such as pH, EC and organic carbon analysed after the harvest of crop were non-significantly affected by the application of different gypsum levels, stage of gypsum application and different levels of nitrogen and phosphorus during both the years. Available phosphorus in soil was highest with 225 kg ha⁻¹ gypsum which was at par with 175 and 125 kg ha⁻¹ gypsum, while significantly higher over control. Available calcium and sulphur of soil were maximum with 225 kg ha⁻¹ gypsum, which was at par with 175 kg ha⁻¹ gypsum while significantly higher than lower levels. Available nitrogen, phosphorus, potassium, calcium and sulphur in soil were non-significantly affected by different gypsum application stages. Available nitrogen and phosphorus in soil was highest with 35 kg N ha⁻¹ + 40 kg P₂O₅ ha⁻¹ and significantly higher over other levels of nitrogen and phosphorus. Available calcium and sulphur in soil were non-significantly affected by different nitrogen and phosphorus levels during both the years. The application of 225 kg ha⁻¹ gypsum, gypsum applied in two splits and application of 25 kg N ha⁻¹ + 30 kg P₂O₅ ha⁻¹ resulted maximum pod yield of spring groundnut.

Keywords: *Groundnut; gypsum; nitrogen; phosphorus; potassium.*

1. INTRODUCTION

Groundnut (*Arachis hypogaea* L.) is the prominent and premier oilseed crop of India and belongs to the family *Leguminosae*. Groundnut covered an area of 4.73 million ha with production of 6.73 million tonnes and productivity of 14.22 quintal ha⁻¹ in India during 2018-19 [1]. In Punjab, groundnut crop was grown over an area of 1.3 thousand ha with an average annual production of 2.6 thousand tonnes and the productivity of 19.8 quintal ha⁻¹ during 2018-19 [2]. Groundnut kernels contain 48-50% edible oil, 25-34% protein, 10-20% carbohydrates and are a rich source of vitamins (E, K and B complex). Since groundnut is a legume-oilseed crop, its requirement of phosphorus, calcium and sulphur is quite high. Therefore, the supply of nutrients in a judicious and balanced manner at appropriate time is very important to realize optimum production of the crop. Gypsum is commonly used as a source of calcium and sulphur for groundnut all over the world. Pod development is enhanced by application of gypsum at flowering stage since it increases the availability of Ca and S in the fruiting zone. Application of gypsum improves soil structure which favours effective pegging in groundnut [3]. Apart from providing Ca and S, gypsum also plays a significant role in the reclamation of alkaline soils. It causes micro-acidification therefore lowering down the soil pH and increasing the nutrient availability in soil [4,5]. Calcium increases the growth and survival of the symbiotic bacteria in groundnut which therefore has a positive influence on biological nitrogen fixation. Zharare et al. [6] conducted a study using hydroponic nutrient solutions

containing various concentrations of Ca and observed that the pod formation would not occur in the solutions without calcium. Deficiency of calcium leads to the production of immature pods, black embryo in seed, weak germination of seeds and increases production potential of aflatoxin and thus, decays peanut pod [7,8]. Sulphur is a component of protein and has an important role to play in oil synthesis. Sulphur finds an important role in the synthesis of sulphur containing amino acids like methionine and cysteine and synthesis of proteins, chlorophyll and oil.

Nitrogen is the main structural component of plant cell. It plays a significant role in plant metabolism and is involved in the synthesis of proteins, amino acids and nucleic acids. Groundnut is a self-fertilizing crop, since its most of the nitrogen requirement is met by the nitrogen-fixing bacteria that are present in the root nodules. Williams [9] suggested that at very high yield levels, the nitrogen requirement of groundnut cannot be met from symbiotic nitrogen fixation alone. To meet the nitrogen requirement during early growth stages, nitrogen could be applied as starter dose [10]. Phosphorus plays an important role in formation of root nodules and fixation of atmospheric nitrogen [11]. Phosphorus is an important structural component of membrane system of the cell, chloroplast and mitochondria. It is an essential constituent of nucleic acid, amino acids, phytin, proteins, nucleoproteins and energy rich phosphate bonds (ADP and ATP). It is involved in the transfer of energy in major metabolic processes like photosynthesis, transformation of sugars and

starch and nutrient movement in plants. Imbalanced and inadequate use of nutrients is the main reason for lower yield of groundnut. However, very less information on the balanced nutrition of spring groundnut is available. Therefore, there is a need to develop a nutrient management strategy to achieve the potential production of spring groundnut. Keeping all these points in view the present study was undertaken to find out the optimum mineral nutrition of spring groundnut.

2. MATERIALS AND METHODS

The field experiments of the present study were conducted at the Students' Research Farm, Department of Agronomy, Punjab Agricultural University, Ludhiana during spring season of 2018 and 2019. Ludhiana is located at an elevation of 247 m above mean sea level and lie at 30°56' latitude and 75°52' longitude, which represents the central agro-climatic zone of Punjab. The average weekly maximum and minimum temperatures during the *spring* season of 2018 were 36.2°C and 22.1°C respectively, whereas during *spring* season of 2019, the corresponding values were 36.1°C and 21.5°C respectively. The weekly mean relative humidity ranged from 21-73% and 29-68% during the cropping seasons of 2018 and 2019, respectively. Total rainfall of 223.6 mm was recorded during the crop season of 2018, whereas the corresponding value for the *spring* season of 2019 was 105.9 mm. The soil of experimental field was loamy sand in texture having pH 8.17, electrical conductivity 0.58 dS m⁻¹ and organic carbon 0.33%. The available nitrogen (N), available phosphorus (P), available potassium (K), available calcium (Ca) and available sulphur (S) of the soil were 156.1 kg ha⁻¹, 16.29 kg ha⁻¹, 323.5 kg ha⁻¹, 114 ppm and 24.8 kg ha⁻¹, respectively. The experiment was laid out in a split plot design replicated three times with four gypsum levels (0, 125, 175 and 225 kg ha⁻¹) in combination with two stages of gypsum application (Full at sowing and 50% at sowing + 50% at flower initiation stage) in the main plot and three levels of nitrogen and phosphorus (15 kg N ha⁻¹ + 20 kg P₂O₅ ha⁻¹, 25 kg N ha⁻¹ + 30 kg P₂O₅ ha⁻¹ and 35 kg N ha⁻¹ + 40 kg P₂O₅ ha⁻¹) in the sub-plot. Second split of gypsum was applied at flower initiation stage *i.e.* at 30 days after sowing while all the doses of nitrogen and phosphorus were applied at the time of sowing. Groundnut variety 'TAG 37A' was sown at the spacing of 30 cm × 15 cm in the second week of March during both the years. All the cultural

practices were followed as per general recommendations. Nitrogen and phosphorus were applied through DAP while the remaining nitrogen requirement was met by using urea.

The samples of groundnut plant (haulm and kernel) were collected at harvest, dried in sun and then oven-dried at 60° C for 24 hours. The samples were grounded and then the samples were analysed for nitrogen, phosphorous, potassium, calcium and sulphur content. The N content in haulm and kernel samples was estimated by using Kjeldahl's distillation method [12]. The P content in haulm and kernel samples was estimated by Vanadomolybdo- phosphoric yellow colour method in HNO₃ as suggested by Jackson [12]. K concentration in plant acid extract was determined by using Flame Photometer after digesting the samples with diacid mixture as suggested by Jackson [12]. Calcium concentration in haulm and kernel samples was estimated by versenate titration with 0.01 N EDTA solution using purpurate indicator after digesting the samples with diacid mixture [13]. Sulphur concentration in haulm and kernel samples was estimated by turbidometric method and the intensity of turbidity was determined using colorimeter 34 at 420 nm, as given by Chesnin and Yien [14]. The soil pH was determined in 1:2 soil-water suspension using an Elico-glass electrode pH meter [12]. Electrical conductivity of soil was determined in 1:2 soil-water suspension equilibrated for 24 hour using a conductivity bridge [12]. Organic carbon was determined by rapid titration method as detailed by Walkley and Black [15]. Available N of soil was estimated by alkaline potassium permanganate method given by Subbiah and Asija [16]. Available P of soil was estimated by 0.5 N NaHCO₃ method suggested by Olsen et al. [17]. Available K of soil was determined by using neutral 1 N ammonium acetate solution as extracting agent as in the method given by Piper [18]. Available Ca in soil was estimated by versenate titration with 0.01 N EDTA solution as given by Cheng and Bray [13]. Available S of soil was determined by turbidometric method given by Chesnin and Yien [14].

3. RESULTS AND DISCUSSION

3.1 Nutrient Concentration in Plant

Application of different gypsum levels exerted a significant influence on the kernel N, P, K, Ca and S concentration as well as haulm P, K, Ca and S concentration of groundnut plants

(Table 1). N content in haulm was influenced non-significantly by different gypsum levels. However, an increasing trend in the N content of haulm was seen with the increasing levels of the applied gypsum. Application of 225 kg ha⁻¹ gypsum resulted in maximum kernel N, P, K and S concentration and haulm P and S concentration, which was statistically at par with 175 kg ha⁻¹ and 125 kg ha⁻¹ gypsum, while significantly higher than control. Application of each of 225 kg ha⁻¹, 175 kg ha⁻¹ and 125 kg ha⁻¹ gypsum gave similar haulm K content, which was significantly higher over control. Both 225 kg ha⁻¹ gypsum and 175 kg ha⁻¹ gypsum, each of them gave highest kernel and haulm Ca content, which was at par with 125 kg ha⁻¹ gypsum, while significantly higher than the control. Rao and Shaktawat [19] supported the findings of the present study and reported that the application of 250 kg ha⁻¹ gypsum significantly increased the N, P, K, Ca and S concentration in kernel and haulm of groundnut over control. Likewise, Pathak et al. [20] reported that seed Ca and S concentration increased with increase in the rates of gypsum application. The increase in nutrient content due to gypsum application might be due to improved nutritional environment in the rhizosphere as well as in the plant system which leads to translocation of N, P and S to reproductive parts which ultimately increased the N, P and S concentration in the kernel of groundnut [4]. The effect of gypsum application stage on the haulm N, P, K, Ca and S concentration and kernel N, P, K and Ca concentration was found to be non-significant (Tables 1 & 2). However, split application of gypsum recorded higher concentration of N, P, K, Ca and S in haulm and kernel as compared to the application of full dose of gypsum at sowing although the difference was non-significant during both the year of study.

Nitrogen and phosphorus levels exerted a significant influence on the haulm N, P, K, Ca and S concentration and kernel N, P and S concentration of groundnut plants (Tables 1 & 2). However, the K and Ca concentration in kernel was non-significantly influenced by nitrogen and phosphorus levels. The maximum N content in kernels was obtained with the application of 35 kg N ha⁻¹ + 40 kg P₂O₅ ha⁻¹, which was significantly higher as compared to the other two doses of nitrogen and phosphorus. The maximum N and P content in haulm was observed with the application of each of 35 kg N ha⁻¹ + 40 kg P₂O₅ ha⁻¹ and 25 kg N ha⁻¹ + 30 kg P₂O₅ ha⁻¹, which was significantly higher than

that of 15 kg N ha⁻¹ + 20 kg P₂O₅ ha⁻¹. Maximum kernel P and S concentration and haulm K, Ca and S concentration was observed with the treatment receiving 35 kg N ha⁻¹ + 40 kg P₂O₅ ha⁻¹, which was at par with 25 kg N ha⁻¹ + 30 kg P₂O₅ ha⁻¹, while significantly higher than that of 15 kg N ha⁻¹ + 20 kg P₂O₅ ha⁻¹. El-Habbasha et al. [21] reported similar observations and reported that the increase in N levels significantly increased N and K content in seeds and haulm of groundnut. Likewise, Gobarah et al. [22] reported that increasing the dose of phosphorus fertilizer significantly increased N and P content in plants of groundnut. These results were supported by Rao and Shaktawat [19] who reported that the application of 60 kg P₂O₅ ha⁻¹ significantly increased the N, P, K, Ca and S concentration in kernel and haulm of groundnut over 20 kg P₂O₅ ha⁻¹. Nitrogen fertilization may increase the cation exchange capacity of plant roots and thus makes them more efficient in absorbing nutrient ions. The application of higher doses of nitrogen was responsible for better root and shoot development and resulted in greater absorption of nutrients in haulm and kernel [23]. Also, the increase in S uptake because of N application could be due to the synergistic effect of N and S in plants. The increased availability of nutrients in root zone coupled with increased metabolic activity at cellular level might have increased nutrient uptake and their accumulation in vegetative plant parts [24]. Application of phosphorus might have led to the development of extensive root system, thus resulting in the improved absorption of nutrients from the soil.

3.2 Effect of Different Treatments on Properties of Soil

3.2.1 Soil pH

Different gypsum levels were observed to have a non-significant effect on the soil pH after crop harvest (Table 3). However, the range of pH observed was 8.22 to 8.29 between various gypsum levels. These results were supported by Chen and Warren [25] who observed that gypsum application improves the pod filling of groundnut without changing the soil pH. Chakrabarti [26] also reported that the soil pH did not show significant variation in gypsum applied soil. The effect of gypsum application stage on pH of soil after crop harvest was non-significant. Nitrogen and phosphorus levels also had a non-significant effect on soil reaction after crop harvest. pH value of 8.27, 8.26 and 8.25 was observed with 15 kg N ha⁻¹ + 20 kg P₂O₅ ha⁻¹, 25

kg N ha⁻¹ + 30 kg P₂O₅ ha⁻¹ and 35 kg N ha⁻¹ + 40 kg P₂O₅ ha⁻¹ respectively.

3.2.2 Electrical conductivity

A non-significant effect of gypsum levels was noticed on the EC of soil after the harvest of crop (Table 3). EC value of soil varied non-significantly between various gypsum levels ranging between 0.62 to 0.63 dS m⁻¹. Similarly, Chakrabarti [26] reported that the electrical conductivity of soil did not show significant variation in gypsum applied soil. EC of the soil after crop harvest was non-significantly influenced by the stage of application of gypsum. Nitrogen and phosphorus levels also had a non-significant effect on the electrical conductivity of soil after crop harvest. EC value of soil varied non-significantly between various nitrogen and phosphorus levels ranging between 0.62 to 0.63 dS m⁻¹.

3.2.3 Organic carbon

Gypsum levels had a non-significant effect on the organic carbon in soil at harvest (Table 3). An increase, though non-significant in OC of soil was observed with increase in the dose of gypsum. The values of organic carbon ranged between 0.34 to 0.35%. Chakrabarti [26] advocated that the soil organic carbon did not show significant variation in gypsum applied soil. Gypsum application stage also had a non-significant effect on the organic carbon of soil at crop harvest. Nitrogen and phosphorus levels exerted a non-significant influence on the organic carbon of soil at harvest. However, a non-significant increase in the organic carbon of soil was observed with increase in the levels of applied nitrogen and phosphorus.

3.2.4 Available nitrogen

The available N in soil after the crop harvest was influenced significantly by different gypsum levels (Table 4). Successive increase in gypsum dose witnessed an increase in the available N of the soil. 225 kg ha⁻¹ dose of gypsum resulted in highest available N (226.1 kg ha⁻¹), which was statistically at par with 175 kg ha⁻¹ gypsum (221.8 kg ha⁻¹) while significantly higher than the other levels of gypsum. These findings are in corroboration with the results of Aulakh et al. [27] who reported that N availability increased with the application of sulphur through gypsum. The increase in available nitrogen may be due to increased supply of sulphur as sulphur may

enhance the nodulation by increasing the supply of sulphur containing proteins, which are essential for multiplication and growth of rhizobia, which fix atmospheric nitrogen. Available N of soil after crop harvest differed non-significantly with different gypsum application stages. However, split application of gypsum resulted in higher available N (219.2 kg ha⁻¹) as compared to application of full dose of gypsum at sowing. Nitrogen and phosphorus levels exerted a significant influence on the available N of soil after crop harvest. Use of 35 kg N ha⁻¹ + 40 kg P₂O₅ ha⁻¹ resulted in highest available N (229 kg ha⁻¹), which was significantly higher as compared to the other two levels of nitrogen and phosphorus. Mohapatra and Dixit [28] reported that the increase in available N in soil could be due to higher amount of N fixation by *Rhizobium* under more favourable conditions of soil and lysis of nodules and secretion of N from these nodules. Also, application of phosphorus may enhance N₂-fixation in groundnut which in turn, improved the N status of the soil [29].

3.2.5 Available phosphorus

Gypsum levels exerted a significant influence on the available P of soil after the harvest of crop (Table 4). Application of 225 kg ha⁻¹ gypsum gave maximum available P (22.84 kg ha⁻¹), which was statistically at par with 175 kg ha⁻¹ (22.39 kg ha⁻¹), while significantly higher than 125 kg ha⁻¹ and control. The effect of gypsum application stage on available P was non-significant. Split application of gypsum resulted in higher available P of soil (21.82 kg ha⁻¹) as compared to basal application of gypsum, although the difference was non-significant. Available P of soil after crop harvest was significantly influenced by the different levels of nitrogen and phosphorus. Use of 35 kg N ha⁻¹ + 40 kg P₂O₅ ha⁻¹ recorded highest available P (23.57 kg ha⁻¹), which was significantly higher over other doses of nitrogen and phosphorus. The increase in phosphorus availability might be due to synergistic effect of nitrogen with phosphorus which increased the availability of P in the soil.

3.2.6 Available potassium

Gypsum levels had non-significant effect on the available K of soil after crop harvest (Table 4). The available K of soil displayed an increase with increased levels of gypsum, although the increase was non-significant. Gypsum application stage also exerted a non-significant influence on the available K of soil. Split

application of gypsum resulted in higher available K of soil (260.7 kg ha^{-1}) as compared to the basal application of gypsum, although the difference was non-significant. Nitrogen and phosphorus levels exercised a significant effect on the available K of soil after the harvest of crop. Maximum available K in soil was recorded with the treatment $35 \text{ kg N ha}^{-1} + 40 \text{ kg P}_2\text{O}_5 \text{ ha}^{-1}$ (267.8 kg ha^{-1}), which was statistically at par with $25 \text{ kg N ha}^{-1} + 30 \text{ kg P}_2\text{O}_5 \text{ ha}^{-1}$ (262.9 kg ha^{-1}), while significantly higher than treatment $15 \text{ kg N ha}^{-1} + 20 \text{ kg P}_2\text{O}_5 \text{ ha}^{-1}$ (248.5 kg ha^{-1}). Parallel findings were observed by Hasan [30] who noticed that the application of $27 \text{ kg N ha}^{-1} + 82 \text{ kg P}_2\text{O}_5 \text{ ha}^{-1}$ resulted in higher potassium content of soil as compared to lower doses of nitrogen and phosphorus. The increase in potassium availability might be due to synergistic effect of nitrogen with potassium which increased the availability of K in the soil [31].

3.2.7 Available calcium

The available Ca of soil after crop harvest was influenced significantly by various gypsum levels (Table 4). Use of 225 kg ha^{-1} gypsum recorded highest available Ca in soil (109.4 ppm), which was statistically at par with that of 175 kg ha^{-1} gypsum (108.1 ppm) while significantly higher than 125 kg ha^{-1} gypsum and control. Similarly, Sharma et al. [32] and Puntamkar et al. [33] observed an increase in the available Ca of soil with application of gypsum. Gypsum application stage had a significant influence on the available Ca of soil after crop harvest. Split application of gypsum recorded significantly higher available Ca of soil (107.2 ppm) in comparison to application of full dose of gypsum at sowing (105.5 ppm). Hallock and Allison [34] also achieved similar results and reported that the application of gypsum at early flowering stage resulted in higher Ca content in soil as compared to control. Available Ca of soil after crop harvest was affected non-significantly by various doses of nitrogen and phosphorus. However, with increase in the levels of nitrogen and phosphorus, an increasing trend of soil available Ca was observed, though the difference was non-significant.

3.2.8 Available sulphur

Application of different gypsum doses had a significant influence on the available S of soil after crop harvest (Table 4). Use of 225 kg ha^{-1}

gypsum recorded highest available S in soil (30.36 kg ha^{-1}), which was statistically at par with that of 175 kg ha^{-1} gypsum (29.42 kg ha^{-1}) while significantly higher than 125 kg ha^{-1} gypsum and control. The results were confirmed by Jat and Ahlawat [35] who reported that with an increase in the dose of gypsum application, the available S content in soil was significantly improved. Gypsum application stage had a non-significant influence on the available S of soil after the harvest of crop. Split application of gypsum resulted in higher available S in soil (28.52 kg ha^{-1}) as compared to application of full dose of gypsum at sowing, though the difference was non-significant. Nitrogen and phosphorus levels exerted a non-significant effect on the available S of soil after crop harvest. However, with increase in the levels of nitrogen and phosphorus, an increasing trend of available S in soil was observed.

3.3 Yield

Application of 225 kg ha^{-1} gypsum resulted in maximum pod yield of 43.04 q ha^{-1} in 2018 and 46.84 q ha^{-1} in 2019 and it was higher as compared to the other doses (Control, 125 and 175 kg ha^{-1}) of gypsum (Fig. 1). Sulphur might have encouraged total biomass production and kernel development in groundnut, which was finally reflected in the improved pod yield. Calcium plays an important role in groundnut pod development and it was necessary for proper filling of pods in adequate quantities in the fruiting zone. Gypsum application stage also significantly influenced the pod yield of spring groundnut during both the years. Highest pod yield of 39.49 q ha^{-1} in 2018 and 42.99 q ha^{-1} in 2019 was observed with the split application of gypsum (50% at sowing + 50% at flower initiation stage) which was higher over application of full dose of gypsum at the time of sowing. Application of $35 \text{ kg N ha}^{-1} + 40 \text{ kg P}_2\text{O}_5 \text{ ha}^{-1}$ gave maximum pod yield of 39.66 q ha^{-1} in 2018 and 43.68 q ha^{-1} in 2019 which was comparable with the pod yield obtained by the application of $25 \text{ kg N ha}^{-1} + 30 \text{ kg P}_2\text{O}_5 \text{ ha}^{-1}$ (38.68 q ha^{-1} in 2018 and 42.36 q ha^{-1} in 2019) while higher than that of $15 \text{ kg N ha}^{-1} + 20 \text{ kg P}_2\text{O}_5 \text{ ha}^{-1}$ (36.60 q ha^{-1} in 2018 and 39.43 q ha^{-1} in 2019) during both the years. The application of increased levels of nitrogen and phosphorus resulted in more nutrient availability to plant and resulted in greater utilization of assimilates into pods and ultimately increased the yield of groundnut.

Table 1. Nitrogen, phosphorus and potassium concentration in haulm and kernel as influenced by gypsum levels, gypsum application stage and nitrogen and phosphorus levels

Treatments	N (%)				P (%)				K (%)			
	Haulm		Kernel		Haulm		Kernel		Haulm		Kernel	
	2018	2019	2018	2019	2018	2019	2018	2019	2018	2019	2018	2019
Gypsum levels												
Control	1.75	1.73	4.16	4.18	0.268	0.268	0.540	0.551	1.53	1.54	0.732	0.727
125 kg ha ⁻¹	1.75	1.75	4.31	4.32	0.278	0.279	0.561	0.566	1.56	1.57	0.761	0.756
175 kg ha ⁻¹	1.76	1.76	4.32	4.33	0.278	0.281	0.562	0.567	1.57	1.57	0.762	0.763
225 kg ha ⁻¹	1.76	1.77	4.32	4.36	0.280	0.281	0.564	0.567	1.57	1.58	0.763	0.768
CD (p=0.05)	NS	NS	0.13	0.13	0.008	0.009	0.019	0.012	0.03	0.03	0.024	0.023
Gypsum application stage												
Full at sowing	1.74	1.75	4.24	4.28	0.276	0.276	0.556	0.560	1.55	1.56	0.753	0.747
50% at sowing + 50% at flower initiation stage	1.76	1.76	4.31	4.31	0.277	0.278	0.558	0.566	1.56	1.57	0.756	0.759
CD (p=0.05)	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Nitrogen and phosphorus levels												
15 kg N ha ⁻¹ + 20 kg P ₂ O ₅ ha ⁻¹	1.72	1.72	4.21	4.17	0.271	0.273	0.545	0.556	1.52	1.55	0.746	0.749
25 kg N ha ⁻¹ + 30 kg P ₂ O ₅ ha ⁻¹	1.77	1.77	4.28	4.29	0.279	0.279	0.562	0.564	1.56	1.58	0.758	0.754
35 kg N ha ⁻¹ + 40 kg P ₂ O ₅ ha ⁻¹	1.77	1.77	4.34	4.43	0.279	0.280	0.564	0.568	1.58	1.58	0.760	0.756
CD (p=0.05)	0.04	0.05	0.05	0.11	0.006	0.005	0.012	0.008	0.03	0.02	NS	NS
Interaction	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS

Table 2. Calcium and sulphur concentration in haulm and kernel as influenced by gypsum levels, gypsum application stage and nitrogen and phosphorus levels


Treatments	Ca (ppm)				S (%)			
	Haulm		Kernel		Haulm		Kernel	
	2018	2019	2018	2019	2018	2019	2018	2019
Gypsum levels								
Control	1.11	1.11	0.360	0.364	0.245	0.248	0.347	0.352
125 kg ha ⁻¹	1.16	1.16	0.376	0.376	0.254	0.255	0.362	0.364
175 kg ha ⁻¹	1.19	1.18	0.376	0.378	0.255	0.257	0.364	0.366
225 kg ha ⁻¹	1.19	1.19	0.377	0.378	0.257	0.258	0.365	0.367
CD (p=0.05)	0.04	0.05	0.010	0.010	0.008	0.007	0.012	0.010
Gypsum application stage								
Full at sowing	1.16	1.16	0.372	0.372	0.252	0.252	0.357	0.360
50% at sowing + 50% at flower initiation stage	1.17	1.17	0.372	0.376	0.253	0.257	0.362	0.365
CD (p=0.05)	NS	NS	NS	NS	NS	NS	NS	NS
Nitrogen and phosphorus levels								
15 kg N ha ⁻¹ + 20 kg P ₂ O ₅ ha ⁻¹	1.14	1.14	0.370	0.371	0.248	0.251	0.352	0.356
25 kg N ha ⁻¹ + 30 kg P ₂ O ₅ ha ⁻¹	1.18	1.16	0.372	0.374	0.254	0.255	0.362	0.365
35 kg N ha ⁻¹ + 40 kg P ₂ O ₅ ha ⁻¹	1.18	1.18	0.375	0.377	0.256	0.258	0.364	0.366
CD (p=0.05)	0.03	0.02	NS	NS	0.005	0.004	0.008	0.006
Interaction	NS	NS	NS	NS	NS	NS	NS	NS

Table 3. pH, EC and OC of soil (after harvest of crop) as influenced by gypsum levels, gypsum application stage and nitrogen and phosphorus levels

Treatments	pH		EC (dS m ⁻¹)		OC (%)	
	2018	2019	2018	2019	2018	2019
Gypsum levels						
Control	8.24	8.19	0.61	0.62	0.33	0.35
125 kg ha ⁻¹	8.25	8.28	0.62	0.63	0.34	0.34
175 kg ha ⁻¹	8.24	8.33	0.64	0.63	0.35	0.34
225 kg ha ⁻¹	8.22	8.31	0.63	0.63	0.35	0.35
CD (p=0.05)	NS	NS	NS	NS	NS	NS
Gypsum application stage						
Full at sowing	8.24	8.27	0.62	0.62	0.34	0.35
50% at sowing + 50% at flower initiation stage	8.24	8.28	0.62	0.63	0.35	0.35
CD (p=0.05)	NS	NS	NS	NS	NS	NS
Nitrogen and phosphorus levels						
15 kg N ha ⁻¹ + 20 kg P ₂ O ₅ ha ⁻¹	8.25	8.28	0.62	0.62	0.34	0.34
25 kg N ha ⁻¹ + 30 kg P ₂ O ₅ ha ⁻¹	8.23	8.27	0.63	0.63	0.34	0.35
35 kg N ha ⁻¹ + 40 kg P ₂ O ₅ ha ⁻¹	8.23	8.28	0.62	0.63	0.35	0.35
CD (p=0.05)	NS	NS	NS	NS	NS	NS
Interaction	NS	NS	NS	NS	NS	NS

Table 4. Available N, P, K, Ca and S of soil (after harvest of crop) as influenced by gypsum levels, gypsum application stage and nitrogen and phosphorus levels

Treatments	Available nutrients									
	N (kg ha ⁻¹)		P (kg ha ⁻¹)		K (kg ha ⁻¹)		Ca (ppm)		S (kg ha ⁻¹)	
	2018	2019	2018	2019	2018	2019	2018	2019	2018	2019
Gypsum levels										
Control	206.0	205.6	18.70	19.82	252.3	253.6	100.3	102.1	23.57	24.42
125 kg ha ⁻¹	211.1	211.6	20.76	22.44	259.9	257.9	105.5	107.9	27.94	28.62
175 kg ha ⁻¹	226.1	217.5	21.92	22.86	264.8	259.5	107.0	109.2	29.03	29.81
225 kg ha ⁻¹	227.7	224.6	22.10	23.56	269.4	260.5	108.1	110.6	30.04	30.68
CD (p=0.05)	NS	NS	1.56	1.27	NS	NS	2.5	2.6	1.93	1.82
Gypsum application stage										
Full at sowing	216.4	210.2	20.52	21.91	261.3	256.2	104.4	106.6	27.06	27.97
50% at sowing + 50% at flower initiation stage	219.1	219.5	21.22	22.43	261.9	259.6	106.1	108.3	28.24	28.80
CD (p=0.05)	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Nitrogen and phosphorus levels										
15 kg N ha ⁻¹ + 20 kg P ₂ O ₅ ha ⁻¹	206.3	200.5	18.95	19.59	249.6	247.4	104.4	107.0	27.00	27.85
25 kg N ha ⁻¹ + 30 kg P ₂ O ₅ ha ⁻¹	217.8	215.1	20.90	22.54	263.6	262.1	105.3	107.4	27.62	28.45
35 kg N ha ⁻¹ + 40 kg P ₂ O ₅ ha ⁻¹	229.1	228.9	22.75	24.37	271.6	264.1	106.0	108.0	28.32	28.85
CD (p=0.05)	10.6	13.2	1.75	1.66	12.7	12.9	NS	NS	NS	NS
Interaction	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS






Fig. 1. Pod yield of groundnut as influenced by (a) gypsum levels, (b) gypsum application stage and (c) nitrogen and phosphorus levels

4. CONCLUSION

The application of gypsum @ 225 kg ha⁻¹ increased the content of major nutrients and improved the soil fertility status in terms of higher availability of major nutrients in the soil under spring groundnut. Application of gypsum to spring groundnut in two split doses *i.e.* half at sowing time and remaining half at flower initiation stage resulted in higher content of major nutrients in crop as well as higher availability of major nutrients in the soil in comparison with the basal application of gypsum. Application of nitrogen and phosphorus @ 35 kg N ha⁻¹ + 40 kg P₂O₅ ha⁻¹ resulted in maximum nutrient content

as well as improvement in the soil fertility status in comparison to the lower doses. Application of gypsum @ 225 kg ha⁻¹ at two split and application of 25 kg N ha⁻¹ + 30 kg P₂O₅ ha⁻¹ also increased the yield of groundnut.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Anonymous. Area, production and productivity. Ministry of Agriculture and Farmers Welfare, Govt. of India; 2019.

- Available:<https://www.indiastat.com/table/agriculture-data/2/groundnut-production/286906/1207870/data.aspx>.
2. Anonymous. Package of Practices for Crops of Punjab, Kharif. Punjab Agricultural University, Ludhiana. 2020; 89-92.
 3. Agasimani CA, Bablad HB, Hosmani MM. Response to time, form and level of calcium application in groundnut (*Arachis hypogaea*). Indian Journal of Agronomy. 1992;37:493-495.
 4. Alcordo IS, Rechcigl, JE. Phosphogypsum in agriculture: A review. In Advances in Agronomy.1993;49:55-118.
 5. Singh AL, Chaudhari VIDYA. Macronutrient requirement of groundnut: Effects on uptake of macronutrients. Indian Journal of Plant Physiology.2007;12:72-77.
 6. Zharare GE, Blamey FPC, Asher CJ. Calcium nutrition of peanut grown in solution culture. II. pod-zone and tissue calcium requirements for fruiting of a virginia and a spanish peanut. Journal of Plant Nutrition.2009;32:1843-1860.
 7. Grichar WJ, Besler BA, Brewer KD. Comparison of agricultural and power plant by-product gypsum for South Texas peanut production. Texas Journal of Agriculture and Natural Resources. 2002;15:44-50.
 8. Murata MR. The impact of soil acidity amelioration on groundnut production on sandy soils of Zimbabwe. *Ph.D. Thesis*, University of Pretoria, Zimbabwe; 2003.
 9. Williams MJ. Growth characteristics of rhizoma peanut and nitrogen fertilized bahiagrass swards. Agronomy Journal. 1994;86:819-823.
 10. Farag IA, Zahran AA. Groundnut (*Arachis hypogaea* L.) growth and yield responses to seed irradiation and mineral fertilization. IOSR Journal of Agriculture and Veterinary Science.2014;7:63-70.
 11. Brady NC, Well RR. The nature and properties of soils. 13 th Edition, Pearson Education Pvt. Ltd., Singapore. 2002;1:05-108.
 12. Jackson ML. Soil Chemical Analysis. Prentice Hall of India (Pvt.) Ltd., New Delhi; 1967.
 13. Cheng KL, Bray RH. Determination of calcium and magnesium in soil and plant material. Soil Science.1951;72:449-458.
 14. Chesnin L, Yien CH. Turbidometric determination of available sulphate. Soil Science Society of America Proceedings. 1951;15:149-151.
 15. Walkley A and Black CA. An examination of the digestion method for the determination of soil organic matter, and a proposed modification of chromic acid titration method. Soil Science. 1934;37: 29-39.
 16. Subbiah BV, Asija GL. A rapid procedure for estimation of available nitrogen in soils. Current Science. 1956;25:259-260.
 17. Olsen SR, Cole CV, Watanable FS, Dean LA. Estimation of available phosphorus by extraction with sodium bicarbonate. United States Department of Agriculture Circular. 1954;9:39.
 18. Piper CS. Soil and Plant Analysis. University of Adelaide, Australia; 1966.
 19. Rao SS, Shaktawat MS. Effect of organic manure, phosphorus and gypsum on nutrient uptake in groundnut. Agropedology. 2005;15:100-106.
 20. Pathak BP, Jain M, Tillman BL, Grusak MA, Gallo M. Effect of gypsum application on mineral composition in peanut pod walls and seeds. Crop Science. 2013;53:1658-1667.
 21. El-Habbasha SF, Taha MH, Jafar NA. Effect of nitrogen fertilizer levels and zinc foliar application on yield, yield attributes and some chemical traits of groundnut. Research Journal of Agriculture and Biological Sciences.2013;9:1-7.
 22. Gobarah ME, Mohamed MH, Tawfik MM. Effect of phosphorus fertilizer and foliar spraying with zinc on growth, yield and quality of groundnut under reclaimed sandy soils. Journal of Applied Sciences Research. 2006;2:491-496.
 23. Patel HK, Patel RM., Desai CK, Patel HB. Response of summer sesamum (*Sesamum indicum* L.) to different spacings and levels of nitrogen under north Gujarat condition. International Journal of Agricultural Sciences.2014;10:336-343.
 24. Sharma YK, Singh H, Mandal N. Effect of nitrogen and copper levels on yield and uptake of nutrients by wheat crop. Annals of Plant Soil Research. 2013;15:43-46.
 25. Chen L, Warren AD. Gypsum as an agricultural amendment. General use guidelines. The Ohio State University Extension; 2011.
 26. Chakrabarti S. Impact of gypsum application on groundnut growing soils of command area. M.Sc. Thesis, Marathwada

- Agricultural University, Parbhani, India; 1990.
27. Aulakh MS, Pasricha NS and Sahota NS. Comparative response of groundnut (*Arachis hypogaea* L.) to three phosphatic fertilizers. *Journal of the Indian Society of Soil Science*. 1980;28:342-346.
 28. Mohapatra AKB and Dixit L. Integrated nutrient management in rainy season groundnut (*Arachis hypogaea*). *Indian Journal of Agronomy*. 2010;55:123-127.
 29. Agboola AA, Fayemi AA. Fixation and excretion of nitrogen by tropical legumes. *Agronomy Journal*. 1972;64:409-412.
 30. Hasan M. Effect of *Rhizobium* inoculation with phosphorus and nitrogen fertilizer on physico-chemical properties of the groundnut soil. *Environment and Ecosystem Science*. 2018;2:04-06.
 31. Bhikane SS. Response of cowpea (*Vigna unguiculata* L.) to application of organic manures with and without inorganic fertilizers. M.Sc. Thesis, Dr.Balasaheb Sawant Konkan Krishi Vidyapeeth. Dapoli, India; 2002.
 32. Sharma ML. Physical and physico-chemical changes in the profile of sodic soil treated with gypsum. *Soil Research*. 1971;9:73-82.
 33. Puntamkar SS, Mehta PC, Seth SP. Effect of gypsum and manure on the growth of wheat irrigated with bicarbonate rich water. *Journal of the Indian Society of Soil Science*. 1972;20:281-285.
 34. Hallock DL, Allison AH. Effect of three Ca sources applied on peanuts II, Soil Ca, K, and Mg levels. *Peanut Science*. 1980;7:26-31.
 35. Jat RA, Ahlawat IPS. Effect of organic manure and sulphur fertilization in pigeonpea (*Cajanus cajan*) + groundnut (*Arachis hypogaea*) intercropping system. *Indian Journal of Agronomy*. 2010;55: 276-281.

© 2023 Brar and Manhas; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/106028>