



# Effect of Seed Priming with Plant Growth Regulators on Growth Parameters, Biochemical Changes of Rice (*Oryza sativa* L.)

Deepak Tripathi <sup>a\*</sup>, Niraj Yadav <sup>a</sup>, Shikha Shahi <sup>b</sup>  
and R. K. Yadav <sup>a</sup>

<sup>a</sup> Department of Crop Physiology, Acharya Narendra Deva University of Agriculture & Technology, Narendra Nagar (Kumarganj), Ayodhya, (U.P.), India.

<sup>b</sup> Department of Crop Physiology, Chandra Shekhar Azad University of Agriculture & Technology, Kanpur, (U.P.), India.

## **Authors' contributions**

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## **Article Information**

DOI: 10.9734/IJPSS/2023/v35i193572

## **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/104645>

**Original Research Article**

**Received: 03/06/2023**

**Accepted: 09/08/2023**

**Published: 22/08/2023**

## **ABSTRACT**

The investigation entitled "Effect of seed priming with plant growth regulators on growth parameters, biochemical changes of rice (*Oryza sativa* L.)" was conducted during *kharif* season, 2018 at Student Instructional Farm (SIF) of Acharya Narendra Deva University of Agriculture & Technology, Kumarganj, Ayodhya-224 229 (U.P.) in randomized block design with eight treatments and three replications on rice variety Sarju-52. The treatments were comprised of priming with plant growth regulators of different concentrations viz salicylic acid (50 ppm, 100 ppm and 150 ppm), KNO<sub>3</sub> (1%, 1.5% and 2%) along with unprimed control and hydro-priming. Seed priming was done for 12 hours in different treatments and nursery was raised. Twenty one days old seedlings were

\*Corresponding author: E-mail: [deepaktripathi25vill@gmail.com](mailto:deepaktripathi25vill@gmail.com);

transplanted. Observations were recorded on various growth parameters at 30, 60, 90 DAT such as leaf area index (LAI), relative water content (RWC), biochemical parameters like chlorophyll content (SPAD Value), catalase activity (EU mg<sup>-1</sup> protein min<sup>-1</sup>), total soluble sugar (mg g<sup>-1</sup> dry weight), at harvest stages. All the plant growth regulators viz salicylic acid (50 ppm, 100 ppm and 150 ppm) and KNO<sub>3</sub> (1%, 1.5% and 2%) influenced positively on the growth characters, chlorophyll content, & yield content in rice crop. Seed priming of KNO<sub>3</sub> @ 2% was found superior followed by salicylic acid 150 ppm over rest of the treatments.

**Keywords:** Rice; seed priming; KNO<sub>3</sub> solution; ppm.

## 1. INTRODUCTION

Rice (*Oryza sativa* L.) is the most important cereal *kharif* season crop as it is a staple food of more than 70 percent of world population. That is why the rice production always holds a key role in the overall food situation of the whole world. Rice is a chief source of carbohydrate. Rice belongs to genus *Oryza* of Gramineae family. The genus *Oryza* includes 24 species, of which 22 wild and two are namely *Oryza sativa* and *Oryza glaberrima* are cultivated. All the varieties found in West Africa belong to species *O. glaberrima*. *Oryza sativa* is a diploid species having 24 chromosomes. The *Oryza* species of the world are commonly grouped into three subspecies namely *indica*, *japonica* and *javanica*. According to advance estimate, rice is cultivated in 43.57 million hectare which is with a production of 104.32 million tonnes and productivity is 2550 kg/ha of our country and state production of rice is 12.95 million tonnes and area is 5.65 million hectare [1] Seed priming was proposed by Heydecker in 1973. Seed priming is an effective technology to enhance rapid and uniformed emergence and to achieve high vigour, leading to better stand establishment and yield [2]. Seed priming by using different chemical solutions (Osmo-priming) has made some considerable effect on plant growth. Seed priming has been successfully demonstrated to improve germination, seed emergence, uniform and vigorous crop stand in seeds of many field crops, such as vegetables and small seeded grasses, wheat, sugar beet, maize, soybean and sunflower [3]. Salicylic acid is a natural compound that plays a central role in certain physiological processes and defense responses in plants [4]. Potassium nitrate has a chemical formula of KNO<sub>3</sub>. It is an ionic salt of potassium ions K<sup>+</sup> and nitrate ions NO<sub>3</sub><sup>-</sup> and is, therefore, an alkali metal nitrate. It is a natural source of nitrate and has been used as a constituent for several different purposes, including food preservatives, fertilizers etc. Priming changes the interior condition of the embryo also. In here it

has reported that KNO<sub>3</sub> priming increased the embryo length in tetraploid watermelon seeds [5].

## 2. MATERIALS AND METHODS

This work was conducted at Student Instructional farm (SIF) of Acharya Narendra Deva University of Agriculture and Technology, Kumarganj, Ayodhya during *kharif* season 2018. The work was laid down in randomized block design with three replications. The treatments were comprised of priming with plant growth regulators of different concentrations viz salicylic acid (50 ppm, 100 ppm and 150 ppm), KNO<sub>3</sub> (1%, 1.5% and 2%) along with unprimed control and hydro-priming. Seed priming was done for 12 hours in different treatments and nursery was raised. Twenty one days old seedlings were transplanted.

### 2.1 Plant Leaf Area

The total number of leaves of two hills was divided into three groups and measured the length and broadest width of each group. The leaf area was calculated by multiplying this area to number of leaves in each group and a factor. This factor is determined on the basis of actual area measured by leaf area meter and recorded by above method. After calculating total leaf area it was divided by ground area in order to get leaf area index. The LAI was recorded three times during the course of investigation of crop at 30, 60 and 90 DAT and then averaged.

### 2.2 Relative Water Content (%)

The relative water content (RWC) was determined by the method described by Turner and beg (1981). Leaf discs were cut from the leaves, weighed and saturated by floating on distilled water in petridish for six hours. The discs were surface dried and weighed. After that discs were kept in oven at 75 ± 5 0 C for 24 hours. After drying, weight of discs was calculated with the help of electronic balance.

### 2.3 Chlorophyll Content (SPAD Value)

Chlorophyll content of leaf was directly measured from intact leaves microprocessor based plant efficiency analyzer model: X55/M-PEA. The values were expressed as SPAD Value. Plant efficiency analyzer can measure three parameters (chlorophyll, nitrogen and leaf temperature) at the same time.

### 2.4 Total Soluble Sugar Content (mg g<sup>-1</sup> Dry Weight)

The total soluble sugar content in shoot was estimated by the method of Yemm and Wills [6].

#### Reagents:

- (1) Ethanol (80%)
- (2) Anthrone (0.2%)
- (3) Glucose

**Anthrone Reagent:** 200 mg Anthrone in 100 ml concentrated H<sub>2</sub>SO<sub>4</sub>

100 mg dried plant sample was homogenized in 10 ml of 80% ethanol centrifuged at 4000 rpm for 20 minutes. Supernatant was collected and residue was re-extracted with 10 ml of 80% ethanol and again centrifuged at 4000 rpm for 20 minutes. Supernatant was collected and both the supernatants were combined and made up to final volume of 20 ml. 0.1-0.2 ml supernatant was dried in a test tube on water bath and cooled to room temperature. One ml of distilled water was added to each test tube and mixed thoroughly. Four ml of anthrone reagent was added along the wall of the test tube and mixed gently and heated on water a bath at 100 0 C for 10 minutes and cooled rapidly under running cold water. Absorbance was measured at 620 nm against the reagent blank. The amount of total carbohydrate present in the extract was calculated using a standard curve prepared from graded concentration of glucose.

### 2.5 Catalase Activity (unit g<sup>-1</sup> Fresh Weight min<sup>-1</sup>)

Catalase activity was assayed calorimetrically according to method given in analytical biochemistry [7]. Catalase facilitates the dismutation of H<sub>2</sub>O<sub>2</sub> to water and O<sub>2</sub> according to the reaction. The enzyme plays an important role in association with SOD as well in photorespiration and glycolate pathway.

#### Reagents:

- I. Phosphate buffer, 0.1 M, pH 7.0
- II. Potassium dichromate acetic acid (5% potassium dichromate glacial acetic acid in 1:3 ratio)
- III. H<sub>2</sub>O<sub>2</sub> 0.2 M

200 mg of fresh leaves material was homogenized with 10 ml of phosphate buffer 0.1 M (pH 7.0) and centrifuged at 10000 rpm for 30 minutes at 4<sup>0</sup>C.

**Table 1. Reaction mixture**

Test	Blank	Reagents
1.25 ml	-	H <sub>2</sub> O <sub>2</sub>
0.50 ml	0.50 ml	Enzyme extract
3.25 ml	4.50 ml	Phosphate buffer
5.00 ml	5.00 ml	

The reaction mixture was taken in Erlenmeyer flask and mixed rapidly at 37 0 C. At 3 minutes interval, 2.0 ml of reaction mixture was withdrawn in test tube and to it, 2.0 ml potassium dichromate acetic acid solution was added and kept on water bath for 10 minutes and colour intensity was measured after cooling at 570 nm on spectronic. Results were expressed as enzyme unit's gram<sup>-1</sup> fresh weight min<sup>-1</sup>

## 3. RESULTS AND DISCUSSION

### 3.1 Leaf Area Index

Data with respect to leaf area index as affected by different treatment of seed priming was recorded at different growth stages have been presented in Table 2 and Fig. 1. The effect of KNO<sub>3</sub> @ 2% and salicylic acid @150ppm was found more pronounced and both the treatments registered significant increase in leaf area index as compared to untreated control at all the stages of observation. The maximum increased in leaf area index was recorded with KNO<sub>3</sub> @ 2% (4.44, 5.28, 5.15 at 30, 60, 90 DAT respectively), followed by salicylic acid @150ppm treatment. However, the minimum leaf area index was (2.23, 2.39, 2.31 at 30, 60, and 90 DAT respectively) was recorded with hydro Priming against control. Sujatha [8] reported that foliar application of salicylic acid (100 ppm) on green gram at 75 DAS increased the plant height (50.4 cm), root length (16.9 cm), number of leaves (18.4) and LAI (1.30). These results are in accordance with the findings of Pour et al. [9], Anisa et al. [10], in rice.

### 3.2 Relative Water Content (RWC) %

The examination of the data presented in Table 3 and Fig. 2 revealed that all the treatments significantly increased in relative water content in leaves at all the stages of the observation with respect to control except hydro-priming. The maximum relative water content (%) was analyzed with  $\text{KNO}_3$  @ 2%, seed priming, (85.48, 82.56, 76.12 at 30, 60, 90, DAT respectively) followed by salicylic acid @150 ppm over rest of the treatments. At 30, 90 DAT salicylic acid @ 150 ppm was found at par with  $\text{KNO}_3$  @ 2% treatment. However, minimum relative water content was recorded with hydro Priming over control. The value of RWC decreasing trends were observed with increasing the crop age. Yildirim et al. [11] reported that ameliorative effect of salicylic acid might be linked to the observable increase in WC, RWC and photosynthetic pigments as well as, leaf area consequently the efficiency of the photosynthetic apparatus was increased due to salicylic acid treatments. Similar finding have been reported by Pour et al. [9] in fenugreek.

### 3.3 Total Chlorophyll Content In Leaves (SPAD Value)

The data presented in Table 4 and Fig. 3 clearly indicate that in general, chlorophyll content increased up to 60 DAT after that decline trend was recorded in primed as well as unprimed (control). All the seed priming treatments improved chlorophyll content in leaf at each stage of observation in comparison to unprimed (control). However, the effect of  $\text{KNO}_3$  @ 2% and salicylic acid @ 150 ppm was found most promising and both the treatments registered significant increase in chlorophyll content in leaf as compared to unprimed control at all the

stages of observation. The higher chlorophyll content was recorded with seed priming of  $\text{KNO}_3$  @ 2% (7.11, 11.38 and 9.78 SPAD value at 30, 60 and 90 DAT, respectively) followed by salicylic acid @ 150 ppm (6.48, 10.34, 8.66 SPAD value) over rest of the treatments. The minimum chlorophyll content was recorded with hydro-Priming (4.54, 8.58, 6.82 at 30, 60, and 90 DAT respectively) against control. These results are in accordance with the findings of El-Tayed [12] in barley, Afzal [13] in tomato.

### 3.4 Total Soluble Sugar ( $\text{mg g}^{-1}$ Dry Weight)

It is evident from the data presented in Table 5 and Fig. 4 that all the seed priming statistically significant improved of total soluble sugar content in dry leaf up to 60 DAT and after that decline trend was recorded. Maximum increase in total soluble sugar content was recorded in case of  $\text{KNO}_3$  @ 2% seed priming at 30, 60, 90 DAT and harvest stages (136.81, 280.31, 250.51, and 239.36  $\text{mg g}^{-1}$  respectively) followed by salicylic acid @ 150ppm over other treatments. At 60 and harvest stages DAT salicylic acid @150ppm was found at par with  $\text{KNO}_3$  @ 2% treatment. While, minimum total soluble sugar was recorded with hydro priming as compare to control. Increase in total soluble sugar by salicylic acid have been reported by Bharti and Malik [14], in mustard, Azooz [15], in barley, Dashtmain et al., [16], in rice.

### 3.5 Catalase Activity ( $\text{EU mg}^{-1}$ Protein $\text{min}^{-1}$ )

The data given in Table 6 and Fig. 5 clearly indicate that in general, leaf catalase content increased up to 90 DAT was recorded in primed as well as control. All the levels and primed seed

**Table 2. Effect of seed priming on leaf area index of rice at different growth duration stages**

Treatments	Leaf area index( LAI)		
	30 DAT	60 DAT	90 DAT
T <sub>1</sub> Control	1.78	2.21	2.17
T <sub>2</sub> Hydro Priming	2.23	2.39	2.31
T <sub>3</sub> SA-50ppm	2.69	2.91	2.85
T <sub>4</sub> SA-100ppm	2.89	3.23	3.11
T <sub>5</sub> SA- 150ppm	4.08	4.78	4.63
T <sub>6</sub> $\text{KNO}_3$ - 1%	3.38	3.48	3.42
T <sub>7</sub> $\text{KNO}_3$ - 1.5%	3.48	3.90	3.85
T <sub>8</sub> $\text{KNO}_3$ -2 %	4.44	5.28	5.15
SEm+ <sub>-</sub>	0.11	0.12	0.12
CD at ( 5%)	0.34	0.35	0.34




Fig. 1. Effect of seed priming on leaf area index of rice at different growth duration stages

Table 3. Effect of seed priming on relative water content of rice at different growth duration stages

Treatments	Relative water content (%)		
	30 DAT	60 DAT	90 DAT
T <sub>1</sub> Control	79.81	75.88	69.90
T <sub>2</sub> Hydro priming	80.12	76.66	70.46
T <sub>3</sub> SA-50ppm	80.32	77.54	71.22
T <sub>4</sub> SA-100ppm	81.83	78.91	72.45
T <sub>5</sub> SA-150ppm	85.02	81.60	75.35
T <sub>6</sub> KNO <sub>3</sub> -1%	82.43	79.31	74.35
T <sub>7</sub> KNO <sub>3</sub> -1.5%	83.45	80.42	73.43
T <sub>8</sub> KNO <sub>3</sub> -2%	85.48	82.56	76.12
SEm+ <sub>-</sub>	0.15	0.29	0.25
CD at (5%)	0.46	0.89	0.77

significantly improved the catalase activity content in leaf at all the stages of observation as compared to control. The maximum increase catalase activity in leaf was registered with @ 2% KNO<sub>3</sub> seed priming at 30, 60, 90 DAT (247.16, 382.58, 413.61 catalase activity,

respectively) followed by salicylic acid @ 150ppm over rest of treatments. However, minimum value was recorded with hydro priming against control. Similar findings have also been reported Chen et al. [17] in rice, Tulio S. Lara et al. [18] in Tomato.

Table 4. Effect of seed priming on chlorophyll content (SPAD Value) of rice at different growth duration stages

Treatments	Chlorophyll content (SPAD Value)		
	30 DAT	60 DAT	90 DAT
T <sub>1</sub> Control	4.39	8.12	6.69
T <sub>2</sub> Hydro Priming	4.54	8.58	6.82
T <sub>3</sub> SA-50ppm	5.18	9.73	7.91
T <sub>4</sub> SA-100ppm	5.37	9.85	8.15
T <sub>5</sub> SA-150ppm	6.48	10.34	8.66
T <sub>6</sub> KNO <sub>3</sub> -1%	5.24	9.96	7.73
T <sub>7</sub> KNO <sub>3</sub> -1.5%	5.97	10.14	8.27
T <sub>8</sub> KNO <sub>3</sub> - 2%	7.11	11.38	9.78
Sem+ <sub>-</sub>	0.21	0.21	0.33
CD at (5%)	0.64	0.63	1.01




Fig. 2. Effect of seed priming on relative water content of rice at different growth duration stages




Fig. 3. Effect of seed priming on chlorophyll content (SPAD Value) of rice at different growth duration stages

Table. 5. Effect of seed priming on total soluble sugar content of rice at different growth duration stages

Treatments	Total soluble sugar content (mg g <sup>-1</sup> dry weight)			
	30 DAT	60 DAT	90 DAT	At harvest
T <sub>1</sub> Control	122.81	265.11	236.32	224.73
T <sub>2</sub> Hydro Priming	123.71	267.81	238.82	225.84
T <sub>3</sub> SA-50ppm	125.60	270.12	241.79	229.15
T <sub>4</sub> SA-100ppm	130.80	274.32	245.13	235.74
T <sub>5</sub> SA-150ppm	136.33	279.80	249.43	238.99
T <sub>6</sub> KNO <sub>3</sub> -1%	133.80	275.15	246.01	236.28
T <sub>7</sub> KNO <sub>3</sub> -1.5%	134.70	277.12	248.37	237.66
T <sub>8</sub> KNO <sub>3</sub> -2%	136.81	280.31	250.51	239.36
SEm+ <sub>1</sub>	0.36	0.33	0.33	0.26
CD at ( 5%)	1.09	1.02	1.00	0.80




Fig. 4. Effect of seed priming on total soluble sugar content of rice at different growth duration stages

Table 6. Effect of seed priming on catalase activity (EU mg<sup>-1</sup> protein min<sup>-1</sup>) of rice at different growth duration stages

Treatments	Catalase activity (EU mg <sup>-1</sup> protein min <sup>-1</sup> )		
	30 DAT	60 DAT	90 DAT
T <sub>1</sub> Control	159.80	298.50	389.67
T <sub>2</sub> Hydro Priming	161.66	311.54	398.50
T <sub>3</sub> SA-50ppm	192.57	321.07	399.64
T <sub>4</sub> SA-100ppm	212.74	332.34	401.29
T <sub>5</sub> SA-150ppm	245.47	381.53	412.78
T <sub>6</sub> KNO <sub>3</sub> -1%	222.29	352.72	404.34
T <sub>7</sub> KNO <sub>3</sub> -1.5%	233.58	361.59	405.69
T <sub>8</sub> KNO <sub>3</sub> -2%	247.16	382.58	413.61
SEm+ <sub>1</sub>	0.76	0.44	0.27
CD at (5%)	2.30	1.35	0.81




Fig. 5. Effect of seed priming on catalase activity (EU mg<sup>-1</sup> protein min<sup>-1</sup>) of rice at different growth duration stages

#### 4. CONCLUSION

The seed priming with KNO<sub>3</sub> and salicylic acid @150ppm was found most promising and both the treatment registered significant increase in chlorophyll content, total soluble sugar content and activity of antioxidant enzyme. The various seed priming sources (KNO<sub>3</sub> and salicylic acid) improved the physiological efficiency of crop which resulted in better growth and yield of rice. Seed priming in salicylic acid @ 150ppm and KNO<sub>3</sub> @ 2% may also be used as a potential tool to enhance 12-16 % yield of rice.

#### CONFERENCE DISCLAIMER

Some part of this manuscript was previously presented in the conference: 6th International Conference on Strategies and Challenges in Agricultural and Life Science for Food Security and Sustainable Environment (SCALFE-2023) on April 28-30, 2023 in Himachal Pradesh University, Summer Hill, Shimla, HP, India. Web Link of the proceeding: <https://www.shobhituniversity.ac.in/pdf/Souvenir-Abstract%20Book-Shimla-HPU-SCALFE-2023.pdf>

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Anonymuos. Agriculture statistics at a glance; 2017-18.
2. Harris D, Rashid A, Hollington A, Jasi L, Riches C. Prospects of improving maize yield with on farm seed priming. In Rajbhandari, N.P. and Ransom J.K. 'Sustainable Maize Production Systems for Nepal'. NARC and CIMMYT, Kathmandu. 2007;180-185.
3. Parera CA, Cantliffe DM. Pre-sowing Brassica seed to improve emergence under different temperatures and soil moisture conditions. Crop Science. 1994;27:1050-1053.
4. Shi Q, Zhu Z. Effects of exogenous salicylic acid on manganese toxicity, element contents and antioxidant system in cucumber. Environ. and Experimental Bot. 2008;63:317-326.
5. Nerson H, Govers A. Salt priming of muskmelon seeds for low temperature germination. Scientia Horticulture. 1985; 28:85-91.
6. Yemma E, Willis A. The estimation of carbohydrates in plants extract by anthrone, Biochem. J. 1954;57:508-514.
7. Sinha AK. Colorimetric assay of catalase. Analytical Biochemistry. 1972;47:38-42.
8. Sujatha KB. Effect of foliar spray of chemicals and bio-regulators on growth and yield of green gram (*Vigna radiate* L.). M.sc. thesis, Tamil Nadu Agric. Univ., Coimbatore; 2001.
9. Pour AP, Farahbakhsh H, Saffari M, Keramat B. Effects of seed priming on germination and seedling growth under salinity stress in fenugreek. IJACS. 2012;4-12:779-786.
10. Ruttanruangboworn A, Chanprasert W, Tobunluepop P, Onwimol D. Effect of seed priming with different concentrations of potassium nitrate on the pattern of seed imbibition and germination of rice (*Oryza sativa* L.). J.I. A. 2017;16(3):605-613.
11. Yildirim B, Yaser F, Ozpay T, Ozypay DT, Turkozu D, Terziodlu O, Tamkoo A. Variations in response to salt stress among field pea genotypes (*Pisum sativum* sp. arvense L.). J. Anim. Veter. Adv. 2008;7:907-910.
12. El-Tayeb MA. Response of Barley grains to the interactive effect of salinity and salicylic acid. Plant Growth Regulation J. 2005;45:215-224.
13. Afzal I, Hussain B, Basra, SMA, Habibullah, S. Halo-priming triggers higher germination potential and early seedling growth of tomato. J. of Agri. & Social Sci. 2011;7(3):105-108.
14. Bharti J, Malik CP. Evaluation of biochemical parameters in nitrate-hardened seeds of Indian mustard subjected to salt stress. J. of oil. Brassica. 2013;4(1):11-18.
15. Ajouri AA, Haben BM. Seed priming enhances germination and seedling growth of barley under conditions of P and Zn deficiency. J. of Plant Nutrition and Soil Sci. 2004;167:630-636.
16. Dashtmain FG, Hosseini MK, Esfahani M. Improving rice seedling physiological and biochemical processes under low temperature by seed priming with salicylic acid. I.J.P.A.E.S. 2014;565.
17. Chen Z, Silva H, Klessig DF. Active oxygen species in the induction of plant



- systemic acquired resistance by salicylic acid. *Science*. 1993;262:1883-1886.
18. Lara TS, Lira JMS, Rodrigues AC, Rakocevic M, Alvarenga AA. Potassium Nitrate Priming Affects the Activity of Nitrate Reductase and Antioxidant Enzymes in Tomato Germination. *J. A. S.* 2014;6(2).

© 2023 Tripathi et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*  
*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/104645>