



Response of Irrigation and Mulching on Yield, Water Use Efficiency, Nutrient Use Efficiency and Economy of Mango Cv. Banganpalli in (Semiarid Region) Southern Telangana Region

Guvvali Thirupathaiah^{1*}, A. Bhagwan², A. Kiran Kumar³, K. Avil Kumar⁴ and D. Vijaya⁵

¹Dept. of Fruit Science, COH, Rajendranagar, SKLTSHU, Hyderabad -30, India.

²Fruit Research Station, Sangareddy, SKLTSHU, Telangana, India.

³College of Horticulture, Rajendranagar, SKLTSHU, Hyderabad -30, India.

⁴Dept. of Agro, PJTSAU, Hyderabad -30, India.

⁵Grape Research Station, Hyderabad, SKLTSHU, Hyderabad -30, India.

Authors' contributions

This work was carried out in collaboration among all authors. Authors GT and AB designed the study, performed the statistical analysis, wrote the protocol and wrote the first draft of the manuscript. Authors AKK and KAK managed the analyses of the study. Author DV managed the literature searches. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IRJPAC/2020/v21i2330320

Editor(s):

(1) Dr. Richard Sawadogo, Research Institute for Health Sciences, Burkina Faso.

Reviewers:

(1) Bala Gambo Jahun, Abubakar Tafawa Balewa University, Nigeria.

(2) Diler Olivera Viciado, São Paulo State University (UNESP), Brazil.

(3) Silvana Fátim Didoné Farmácia, Universidade de Passo Fundo (UPF), Brazil.

Complete Peer review History: <http://www.sdiarticle4.com/review-history/63259>

Original Research Article

Received 02 October 2020

Accepted 07 December 2020

Published 30 December 2020

ABSTRACT

An investigation was carried out on sandy loam soils of semi arid regions of southern Telangana to analyze the potential of drip irrigation along with mulch on mango fruit yield, water and nutrient use efficiency. The treatments of present study were comprised of two levels mulching (M₀-without mulch and M₁-with mulching), with silver Polyethylene of 100 micron thickness were used and two levels of irrigation (I₁ -75% and I₂ 100% ETc through drip), here irrigation levels and mulching together constituting four treatment combinations with five replications under 2x2 factorial randomized block design and the treatment combinations are: I₁M₀-75% ETc + No mulching, I₁M₁-

*Corresponding author: E-mail: thiruicar@gmail.com;

75% ETc + with mulch, I₂M₀- 100% ETc + No mulching and I₂M₁- 100 % ETc + with mulch. The study revealed that drip irrigation 100% ETc along with silver polythene mulch showed better performance in terms of yield, water use efficiency, nutrient use efficiency as well as economics. Maximum yield of 89.11 kg tree⁻¹ combination of 75 % ETc + with mulching has proven the maximum water use efficiency (5.54 g liter⁻¹ water consumption) and fertilizers use efficiency (0.89 q kg⁻¹ fertilizer application) however along with maximum net return of 400973.90 rupees ha⁻¹, net return of 346873.90 ₹ ha⁻¹ per hectare and B: C ratio of 7.41 was recorded in I₂M₁ (100 % ETc + with mulch). So the experiment suggests that drip irrigation along with mulch has the potential to provide greater benefit by optimizing the use of water resources. But the interaction of 75 % ETc + with mulching has proven the maximum water use efficiency (6.28 g liter⁻¹ water consumption).

Keywords: Yield; water and nutrient use efficiency and economy.

1. INTRODUCTION

Mango (*Mangifera indica* L.) is one the most luscious fruit since time immemorial in the tropical and subtropical region of the world and is native of South East Asia (Indo-Burma region). It is designated as the 'King of Fruits' [1] because of its excellent flavour, attractive fragrance, beautiful shades of color and delicious taste with high nutritive value. Presently it is grown in India, China, Thailand, Mexico, Indonesia, Pakistan, Brazil, Egypt, Bangladesh and Nigeria [2]. India is the leader, sharing 41% of the world's mango production [3] with an area of 2.26 million ha with the production of 21.82 m Mt. and average productivity is 9.7 t ha⁻¹ [4].

In Telangana mango occupies a prime position in cultivation among the fruit crops, i.e. 70 per cent of total fruit cultivated area and 56% of total production devoted to mango, but the average productivity of India and Telangana (9.7 t ha⁻¹ and 9.3 t ha⁻¹, respectively) is very low compared to Uttar Pradesh (India) and Bangladesh (17.0 t ha⁻¹ to 17.63 t ha⁻¹) and it indicating that, there is huge potential to increasing the productivity of mango [4]. However, the mango productivity has declined due to several biotic and abiotic factors. Out of several biotic and a-biotic factors responsible, optimum water management is one of the most important factors that significantly influence productivity and quality, even though mango trees are tolerant to drought and occasional flooding. Water stress during the critical stages of fruit growth and development is the main reason for low productivity [5]. In such situation, water management, especially during development stage plays an important role in improving yield and quality. However, the amount and quality of available irrigation water of the arid and semi- arid regions are being main limiting factors for productivity of mango [5].

As the water and land are the important, indispensable resources for agricultural development and economic up-gradation of any country cannot be regarded as available in abundance and free forever. The demand for these two resources will continue to grow due to ever enhancing population. Increasing the water shortage has caused us to investigate the sustainable use of irrigation water. For this, specific effective water-saving irrigation techniques without detrimentally affecting crop production need to be developed urgently. In this context a shift in focus is indispensable for the development of additional water resources along with the efficient use of already developed water resources [6]. As the mango is irrigated by basin and drip irrigation systems without any concern on crop water requirement and schedule; that has leads to inefficient utilization of available water and it must be converted from the basin irrigation system to drip irrigation with a proper scheduled manner, their by the water conserved and it would helps in increase the acreage with available water [7].

In addition to above the mulching is another important conservative cultural practice. As many studies have shown that mulches can improve soil qualities, weed control [8] and also helps in efficient water management under such situation, the drip irrigation along with mulching is the best practice which saves 25-30% irrigation water [9]. Sing et al. [10] suggested that, the irrigation requirement met through drip irrigation along with polyethylene mulch gave the 164 per cent greater yield compared to ring basin irrigation in guava. With an above brief background, an experiment was carried to find out the effect of irrigation scheduling and mulching on yield, water use efficiency, nutrient use efficiency and economy of mango.

2. MATERIALS AND METHODS

2.1 The Study Area

The field experiment was conducted during June-2017 to May-2018 and June-2018 to May-2019 was executed in the private mango orchard at Sathapur (Kollapur area) and the experimental farm is located at 16° 30' N latitude and 78° 19' E longitude at an elevation of 550 m above MSL, representing Southern Telangana zone, is mainly covered by red sandy soil, is locally known as 'Chelka soils' and the soil properties are mentioned in Table 1. Experiment comprised of 40 number of uniform size healthy trees of mango Cv. Banganpalli, 12 years old plants were selected and the plants were planted with spacing of 10x10 m. Two plants were selected for each treatment in each replication.

All plants were given similar cultural operations except irrigation and mulching. The application of recommended dose of fertilizers (RDF *i.e.*, 1000:1000:1000 g of NPK and 50 kg of farm yard manure (FYM) plant⁻¹ year⁻¹ for more than ten years old plants) are applied during third week of June *i.e.* half of doses of nitrogen, potassium, total FYM and phosphorus were applied. Another half dose of nitrogen and potassium was applied through soil, along with the micronutrients [ZnSO₄ (50 g) + FeSO₄ (25 g) + Borax (20g) + MgSO₄ (30 g) tree⁻¹] are applied through foliar spray fruit at marble. Here, two levels M₀ and M₁ (*i.e.* without mulch and with mulching, respectively), mulch with silver Poly Ethylene of 100-micron thickness were used and two levels of irrigation (I₁ -75% and I₂ 100% ETc through drip). Here is a bifactorial 2x2 design were used and There were two levels of irrigation as main

plot and two levels of mulching in the sub plot together constituting four treatment combinations with five replications under factorial randomized block design (OPSTAT programme by COBS&H CCS HAU, Hisar; [11]). Taking into account the scheduling of irrigation at different levels of ETc and mulching were used in the treatment combination: I₁M₀-75% ETc + No mulching, I₁M₁-75% ETc + with mulch, I₂M₀- 100% ETc + No mulching and I₂M₁- 100 % ETc + with mulch.

Irrigation was scheduled by calculating ET using pan evaporation measurements adjusted by crop coefficient. The experimental plots were irrigated by using drip irrigation system and the crop water requirement was calculated daily with the help of meteorological data recorded by meteorological observatory of KVK, Palem, Nagarkurnool, and PJTSAU-Telangana. The irrigation water requirement was estimated by Pan "A" evaporation method, where the daily water requirement was calculated using the formula [12]:

$$V = Kp \times Kc \times Sp \times Sr \times Wp$$

Where,

V = Volume of water required (liter day⁻¹ plant⁻¹),
 Ep = Pan evaporation as measured by Class-A pan evaporimeter mm day⁻¹.
 Kc = Crop co-efficient (co-efficient depends but for fully grown plants is 0.85)
 Kp = Pan co-efficient (0.7)
 Sp = Plant to plant spacing (m) Sr = Row to row spacing (m)
 Wp = Fractional wetted area, which varies with different growth stage (0.3 for wider spaced crops).

Table 1. Physical and chemical properties of experimental soil

Particulars	Value for 0-60 cm depth
A. Physical properties	
Sand (%)	70
Silt (%)	19
Clay (%)	11
Textural class	Red sandy lome
B. Chemical properties	
pH	7.5
Electrical conductivity (dS m ⁻¹ at 25°C)	0.33
Organic carbon (%)	0.56
Available N (kg ha ⁻¹)	185
Available P ₂ O ₅ (kg ha ⁻¹)	23
Available K ₂ O (kg ha ⁻¹)	240

2.2 Water Use Efficiency

Water use efficiency (WUE) was computed from the following relationship and has been expressed as g liter⁻¹ water.

$$\text{Water use efficiency (WUE)} = \frac{Y}{\text{LWA}}$$

Where;

Y- Yield (g L⁻¹)

LWA- liter of water applied

2.3 Nutrient Use Efficiency (NEU- q kg⁻¹ Fertilizer)

The word “efficiency” generally indicates “the level of output per unit of input.” Referred to the plant system, “efficiency” defines the “growth, physiological activity, yield or harvested yield (output) per unit of land, water, nutrient, or energy (input).” While focusing the attention on the nutrients, the term NUE is defined as “the plant growth, physiological activity, yield or harvested yield per unit of nutrient.” Even within this simple ratio, NUE has been defined in 18 different ways in diverse contexts which are primarily grouped in “Agronomic” (uptake efficiency) and “Physiological” (utilization efficiency) terms.

$$\text{NEU (Q kg}^{-1}\text{ fertilizer)} = \frac{\text{Fruit yield in treated plant}}{\text{Nutrient applied for treated plant}}$$

2.4 Economics

The cost of cultivation of mango was worked out for one hectare area and the details of economics are presented in Table 4.

3. RESULTS AND DISCUSSION

3.1 Fruit Yield (kg tree⁻¹)

Significant differences have been recorded with respect to fruit yield (kg tree⁻¹) of mango, among the different treatments (Table 2) and the maximum fruit yield tree⁻¹ (78.83 kg) was recorded in (100 % ETc). However, minimum fruit yield tree⁻¹ (66.48 kg) was recorded in I₁ (75 % ETc). The average increase in the yield per tree is about 15.61 per cent I₂ over I₁. The results are in conformity with the findings of Srinivas [13]; Dixit et al. [14] and Dinesh et al. [15] in mango. Similarly, Shirgure et al. ([16], [17]) noted the maximum fruit yield with irrigation equivalent to 0.8 of open pan evaporation in citrus under Nagpur conditions. Drip irrigation provides a consistent moisture regime in the soil due to

which root remains active throughout the season, resulting in optimum availability of water helps proper translocation of food materials which accelerates the fruit growth and development. Pavel and Villers [18] observed increased fruit yield per tree in mango due to drip irrigation and Coelho and Borges [19] also emphasized the importance of drip irrigation in mango for better yield.

The average data on the effect of mulching on fruit yield tree⁻¹ of mango Cv. Banganpalli had shown the significant difference and the maximum fruit yield tree⁻¹ (82.45 kg) was recorded in M₁ (with mulching), the minimum of fruit yield tree⁻¹ (62.87 kg) was recorded in M₀ (without mulching) and the average increase in the fruit yield per tree is 23.75 per cent. Increase yield was directly related to the reduced weed density, weed control efficiency under mulched condition that resulted in increased availability of soil-water and nutrients to the plants that, subsequently enhanced fruit weight and fruit yield. The present observations were in line with the findings of Kumar [20].

The maximum fruit yield tree⁻¹ (89.11 kg) was recorded in I₂M₁ (100% ETc + with mulching) followed by I₁M₁ (75% ETc + with mulching) with 75.78 kg fruits tree⁻¹. However the minimum fruit yield tree⁻¹ (57.18 kg) was recorded in I₁M₀ (75 % ETc + without mulching). The average fruit yield per tree increased by 35.83 percent over 75 % ETc+ without mulch to at 100 % ETc +with mulching. Similar kind of findings were reported by Kumar et al. [20] under drip irrigated mango Cv. Arka Anmol has obtained the maximum yield of 65.72 kg tree⁻¹ at 75 per cent PER. Similarly, Sharma et al. [21] while working with drip irrigation in guava revealed that 164 per cent greater yield was found in case of drip irrigated guava as compared to ring basin irrigation. Pradhan et al. [22] in mango got a maximum yield under 80 per cent irrigation through drip and plastic mulch. Joshi et al. [23] also reported maximum yield under high density litchi at 100 per cent estimated irrigation water requirement along with black plastic mulch. Several other scientists working with drip irrigation and mulch on various fruit crops also stressed on the significant improvement in yield parameters viz., Dixit et al. [14], Srinivas [13] and Coelho and Borges [19] in mango; Shirgure et al. [24] in Nagpur mandarin, Sulochanamma et al. [25] in pomegranate and Ramniwas et al. [31] in guava etc. Here, the possible reason for highest yield under 100% ETc and mulching, maintained

optimum soil moisture throughout fruit growing period and mulching also has helped in soil moisture and nutrient conservation and favorable hygrothermal balance, which ultimately helps in better yield.

3.2 Water Use Efficiency (g liter⁻¹ water)

Current findings evident that, the water use efficiency (WUE) was found to be influenced variably by different irrigation and mulching treatments (Table 3). The data on main effect of irrigation indicate the WUE to the tune of 5.51 g liter⁻¹ water under I₁ (75 % ETc) and minimum WUE (4.90 g liter⁻¹ water) was recorded in I₂ (100 % ETc). Similarly, maximum WUE (5.91 g liter⁻¹ water) registered with M₁ and minimum (4.50 g liter⁻¹ water) was noted in M₀ (without mulching). Similarly, the maximum WUE (6.28 g liter⁻¹ water) was noted with conjugation of mulching and 75% ETc (I₁M₁). However, minimum WUE (4.26 g liter⁻¹ water) was noted in without mulching and 100 % ETc (I₂M₀). The results suggested that, 75 % ETc was registered highest yield per unit water consumption. From present investigation the highest WUE might be due to the effect of improved soil, microclimate, weed free environment, low evaporation and higher moisture availability in the root zone that helped

in better nutrient uptake by plant resulting in earlier and better vegetative growth, which then enhanced the photosynthesis rate and translocation of synthesized food from leaves to fruits, resulted in early harvesting and increased number of fruits per plant under black plastic mulch. Among the different treatments the conjugation of irrigation and mulching has proven higher WUE i.e. 25 percent water saving over 100 % ETc + un-mulching condition, without adverse effect on yield and similar finding are suggested by Panigrahi *et al.* [26] in mango with 60% water through drip with black polythene mulch and Srivastava *et al.* [27] in banana. Here, the lower WUE in 100 % ETc without mulching due to the water lost through evaporation which lead to least WUE that of 75 % ETc with mulching. Similarly, Tiwari *et al.* [28] also noted higher WUE with 80% of irrigation met through drip irrigation along with plastic mulching in sapota. From the present study it was indicated that I₁ has recorded 12.45 percent more water use efficiency compared to I₂. The mulching has also registered 23.86 % higher WUE compared to unmulched condition. Similarly the interaction of irrigation and mulching the treatment I₁M₁ has noted 32.17 % more WUE compared to I₂M₀ and 14.44 % better WUE with I₁M₁ compared to I₂M₁.

Table 2. Effect of irrigation and mulching on fruit yield (kg tree⁻¹) of mango Cv. Banganpalli*

Particulars	Fruit yield (kg tree ⁻¹)		
	M ₀	M ₁	Mean
I ₁ (75% ETc)	57.18 ^d	75.78 ^b	66.48 ^b
I ₂ (100% ETc)	68.56 ^c	89.11 ^a	78.83 ^a
Mean	62.87 ^B	82.45 ^A	
Factors	F- Test	SEm _±	CD at 5%
Irrigation (I)	*	2.346	7.308
Mulching (M)	*	2.346	7.308
Interaction (IXM)	*	3.317	10.61

(Average values of two years data)

Note: M₀- Without mulch; M₁-With mulch; F-Test (*)-significant; CD at 5 per cent level of significance; NS-Non-significant

Table 3. Water use efficiency of mango (g liter⁻¹ water)*

Particulars	WUE (g liter ⁻¹ water)		
	M ₀	M ₁	Mean
I ₁ (75% ETc)	4.74	6.28	5.51 ^a
I ₂ (100% ETc)	4.26	5.54	4.90 ^b
Mean	4.50 ^b	5.91 ^a	
Factors	F- value	SEm _±	CD at 5%
Irrigation (I)	*	0.17	0.53
Mulching (M)	*	0.17	0.53
Interaction (IXM)	NS	0.24	-

(*Average values of two years data)

F-value (*)-significant; CD at 5 per cent level of significance; NS-Non-significant; M₀- without mulch; M₁-With mulch

Table 4. Fertilizer use efficiency (q kg⁻¹ fertilizer)

Particulars	FUE (q kg ⁻¹ fertilizer)		
	M ₀	M ₁	Mean
I ₁ (75% ETc)	0.57	0.76	0.66 ^b
I ₂ (100% ETc)	0.69	0.89	0.79 ^a
Mean	0.63 ^b	0.82 ^a	
Factors	F- value	SEm _±	CD at 5%
Irrigation (I)	*	0.023	0.073
Mulching (M)	*	0.023	0.073
Interaction (IXM)	NS	0.033	-

(*Average values of two years data)

F-value (*)-significant; CD at 5 per cent level of significance; NS-Non-significant; M₀- without mulch; M₁-With mulch

3.3 Fertilizer Use Efficiency (q kg⁻¹ Fertilizer)

The present study resulted that, the fertilizer use efficiency (FUE) has significantly influenced by different irrigation and mulching treatments (Table 4). The data of main effect of irrigation was indicated the FUE to the tune of 0.79 q kg⁻¹ fertilizer under I₂ (100 % ETc) and minimum FUE (0.66 q kg⁻¹ fertilizer) was recorded in I₁ (75 % ETc). Similarly, maximum FUE (0.82 q kg⁻¹ fertilizer) registered with M₁ and minimum (0.69 q kg⁻¹ fertilizer) was noted in M₀ (without mulching). Similarly, the maximum FUE (0.89 q kg⁻¹ fertilizer) was noted with conjugation of 100 % ETc and mulching (I₂M₁). However, minimum FUE (0.57 q kg⁻¹ fertilizer) was noted in without mulching and 75% ETc (I₁M₀). Present study suggests that, application 100 % ETc along with mulching proven to higher FUE compared to other treatments it may be due to the plastic mulch increase soil moisture by reducing loss of water through evaporation, increase nutrient use efficiency by reducing loss of nutrients through leaching, surface run off or volatilization, eliminates weed growth at the vicinity of tree, thereby providing congenial environment for enhanced tree growth, fruit yield and quality, besides ameliorating leaf nutrient contents and water use efficiency Neilsen et al. [29] in apple and Barman et al. [30] in guava Cv. Lalit. From our study I₂ (100% ETc) has registered 16.46% better FUE as compared to I₁ (75 % ETc). Regarding to mulching, the mulched plants was found 23.17% efficient in fertilizer usage. Similarly the interaction of mulching along with 100% ETc has gave 35.96 % higher FUE over 75 % ETc + without mulching.

3.4 Economics

The economics of mango influenced by different levels of irrigation and fertilization had

shown significant among the treatments and the data is presented in Table 4 and the detail of cost of cultivation is given in Table 5 and 6.

The data on cost of cultivation per hectare revealed that, the irrigation treatment I₂ has recorded maximum gross returns and net returns (3.55 and 3.06 lakh rupees ha⁻¹, respectively) and least values were registered with I₁ (3.00 and 2.50 lakh rupees ha⁻¹), similarly maximum gross returns and net returns (3.71 and 3.16 lakh rupees ha⁻¹, respectively) was registered with M₁ (with mulching) and minimum was noted with M₀ (2.82 and 2.38 lakh rupees ha⁻¹, respectively). The maximum benefit cost ratio (BCR) was registered with I₂ (7.20) and minimum BCR was noted with I₁ (6.07). Here, 100 % ETc gave 15.67 % higher BCR compared to 75 % ETc. Similarly, maximum benefit cost ratio (BCR-6.86) was registered with M₁ (with mulching) and minimum BCR (6.41) was noted with M₀ (without mulching) and the mulched plants has registered 6.56 % more BCR compared to unmulched plants. Similarly, maximum gross returns and net returns (4.00 and 3.46 lakh rupees ha⁻¹, respectively) was registered with I₂M₁ (with mulching) and minimum was noted with I₁M₀ (2.57 and 2.13 lakh ha⁻¹, respectively) and the maximum BCR (7.41) was noted with conjugation of 100% ETc and mulching (I₂M₁). However, minimum BCR (5.83) was noted in without mulching and 75% ETc (I₁M₀). Present study suggests that, application 100 % ETc along with mulching proven to higher BCR compared to other treatments it may be due to the plastic mulch increase soil moisture by reducing loss of water through evaporation, increase nutrient use efficiency by reducing loss of nutrients through leaching, surface run off or volatilization, eliminates weed growth at the vicinity of tree, thereby providing congenial environment for enhanced tree growth, fruit yield resulted higher

Table 5. Cost of cultivation for mango Cv. Banganpalli with different levels of irrigation, fertigation and mulching (₹ =Rupees)

No.	Materials/works	I ₁ M ₀	I ₁ M ₁	I ₂ M ₀	I ₂ M ₁
I.	Inputs	30,941.88	40,941.88	30,941.88	40,941.88
1.	Recommended dose (RDF) is 1000:1000:1000 g of NPK and 50kg FYM plant ⁻¹ year ⁻¹ for >10 years old plants (1.33 kg of Urea; 2.2 kg of DAP and 2.0 kg of SOP per plant)	16480.88	16480.88	16480.88	16480.88
	i. Urea ₹ 5.52 kg ⁻¹	732.00	732.00	732.00	732.00
	ii. DAP ₹ 10.4 kg ⁻¹	2261.00	2261.00	2261.00	2261.00
	iii. SOP ₹ 17.44 kg ⁻¹	348.00	348.00	348.00	348.00
	iv. FYM ₹ 2.00kg ⁻¹	10,000.00	10,000.00	10,000.00	10,000.00
	v. Zinc ₹ 100 kg ⁻¹	500.00	500.00	500.00	500.00
	vi. Iron ₹ 80 kg ⁻¹	200.00	200.00	200.00	200.00
	vii. Boron ₹ 90 kg ⁻¹	180.00	180.00	180.00	180.00
	viii. Magnesium ₹ 80 kg ⁻¹	240.00	240.00	240.00	240.00
	ix. Mulch material 2000 ₹ Roll ⁻¹ (400 x 0.9 m) to cover 20 m canopy area plant ⁻¹ ha ⁻¹	--	10000.00	--	10000.00
II.	Cultural operations, plant protection and labour charges	48,800.0	48,800.0	48,800.0	48,800.0
2.	Spraying of micronutrients 4 labour (300 ₹ . per day)	1200.00	1200.00	1200.00	1200.00
3.	Bunds formation, Weeding, irrigation and fertilizer application 75 labours	21,000.00	21,000.00	21,000.00	21,000.00
4.	Plant protection chemicals and measures (700 litte water ⁻¹ ha ⁻¹)	8,800.00	8,800.00	8,800.00	8,800.00
	i. 16 labours for 4 sprays (300 ₹ . per day)	4,800.00	4,800.00	4,800.00	4,800.00
	ii. Neem oil @ 2.5 ml l ⁻¹ (1.75 l Neem oil and 400 ₹ l ⁻¹)	700.00	700.00	700.00	700.00
	iii. Thiomethoxam @ 0.5 ml l ⁻¹ (0.35 l ha ⁻¹ and 1,600 ₹ l ⁻¹)	560.00	560.00	560.00	560.00
	iv. Imidachloprid @ 0.5 ml l ⁻¹ (0.35 l ha ⁻¹ and 2,000 ₹ l ⁻¹)	700.00	700.00	700.00	700.00
	v. Profenophos +cypermetrin 0.5 ml l ⁻¹ (0.35 l ha ⁻¹ and 1,200 ₹ l ⁻¹)	420.00	420.00	420.00	420.00
	vi. SAAF (Carbendazim 12 % + Mancozeb 63 % wp) @ 2.0 g l ⁻¹ (2.00 kg l ha ⁻¹ (600 ₹ kg ⁻¹)	1200.00	1200.00	1200.00	1200.00
	a. Hexaconazole @ 0.5 ml l ⁻¹ (0.35 l ha ⁻¹ and 1,200 ₹ l ⁻¹)	420.00	420.00	420.00	420.00
5.	15- Labours for each harvesting in control (4,500 .00 ₹ harvest ⁻¹)	9,000.00	9,000.00	9,000.00	9,000.00
III.	Total cost of cultivation	79, 7741.88	89, 7741.88	79, 7741.88	89, 7741.88

(Note: Marketable price of mango Cv. Banganpalli 45 ₹ kg-1)

Table 6. Economy of mango influenced by irrigation and mulching treatments*(□=Rupees)

Particulars	Gross income (₹ ha ⁻¹)			Net income (₹ ha ⁻¹)			BCR		
	M ₀	M ₁	Mean	M ₀	M ₁	Mean	M ₀	M ₁	Mean
I ₁ (75% ETc)	257,298.40	341,028.90	299,163.70	213,198.40	286,928.90	250,063.70	5.83	6.30	6.07 ^B
I ₂ (100% ETc)	308,499.20	400,973.90	354,736.50	264,399.20	346,873.90	305,636.50	7.00	7.41	7.20 ^A
Mean	282,898.80	371,001.40		238,798.80	316,901.40		6.41	6.86	
Factors	F- value	SEm±	CD at 5%	F- value	SEm±	CD at 5%	F- value	SEm±	CD at 5%
Irrigation (I)	*	10,554.73	32,882.56	*	10,554.74	32,882.60	*	0.21	0.64
Mulching (M)	*	10,554.73	32,882.56	*	10,554.74	32,882.60	NS	0.21	-
Interaction (IXM)	NS	14,926.64	-	NS	14,926.66	-	NS	0.29	-

(*Average values of two years data)

F-value (*)-significant; CD: at 5 per cent level of significance; NS-Non-significant; M0- without mulch; M1-With mulch

BCR in mango [22]. From this the interaction of 100 % ETc along with mulching has gave 21.32% higher BCR compared to 75% ETc + unmulched condition, 14.98% more compared to 75% ETc + mulched plants and 5.3% higher BCR compared to 100% ETc + unmulched condition.

4. CONCLUSION

From the investigations on the effect of irrigation and mulching suggested that, the maximum fruit yield, water and nutrient use efficiency, and the higher BCR were also obtained with the application of 100 % ETc along with mulching. But the WUE was highest with 75 % ETc + with mulching, and it is also suggested the during water shortage condition the combination of 75 % ETc + with mulching is an effective way to manage irrigation without any adverse on fruit yield, quality and economy of mango Cv. Banganpalli. In this instance efficient water use is possible with crop improvements by enabling crops to grow successfully under drought environments and use of advanced water-saving irrigation techniques and measures can significantly reduce water consumption, improve agronomic water use efficiency, save labor and costs, reduce the adverse effects of agricultural water resource availability from climate change and relieve the crisis of water resource and so on.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Purselove JW. Mangoes west of India. Acta. Hort. 1972;24:170-174.
2. Pooja Devi, Singh CP, Mansi Darmwa. Drip fertigation and it's implication on vegetative growth and leaf nutrient content of mango cv. Pant Sinduri. International Journal of Chemical Studies 2019;7(4):1836-1839.
3. Ganeshamurthy AN*, Rupa TR, Shivananda TN. Enhancing Mango Productivity through Sustainable Resource Management. Journal of horticultural science. 2018;13(1):1-31.
4. Anonymous. Fruits. Indian Horticulture Database, National Horticulture Board, Ministry of Agriculture, Govt. of India; 2018.
5. Available: <http://nhb.gov.in>
5. Adak T, Kumar K, Singh VK, Singh Vinod Kumar, Singh AK. Effect of different soil moisture regimes on soil moisture rd dynamics, yield and quality in mango. In: Proceedings of 3 International Agronomy Congress on "Agriculture Diversification, Climate Change Management and Livelihoods", held during November 26-30th, 2012 at IARI, New Delhi, India. 2012;1416-1418.
6. Daniel PL, Eelco VB. Water Resources Planning and Management: An Overview. Water Resource Systems Planning and Management. 2017;1-49.
7. Panigrahi P, Srivastava AK. Deficit irrigation (DI) scheduling for matured Nagpur mandarin (*Citrus reticulata* Blanco) trees of central India. Indian Journal of Soil Conservation. 2011;39(2):149-154.
8. Ross OC. Reflective mulch effects on the grapevine environment, Pinot noir vine performance, and juice and wine characteristics. Dissertation, Lincoln University; 2010.
9. Santosh K. Influence of Drip Irrigation and Mulch on Leaf Area Maximization, Water Use Efficiency and Yield of Potato (*Solanum tuberosum* L.). urnal of Agricultural Science. 2012;4(1):71-80.
10. Singh K. Fall in Water Table in Central Punjab – How Serious? The PunjabState Farmers Commission, Chandigarh. 2006;13.
11. Sheoran OP, Tonk DS, Kaushik LS, Hasija RC, Pannu RS. Statistical Software Package for Agricultural Research Workers. Recent Advances in information theory, Statistics & Computer Applications by D.S. Hooda & R.C. Hasija Department of Mathematics Statistics, CCS HAU, Hisar. 1998;139-143.
12. Shukla KN, Singh PK, Singh KK. Crop water requirement under drip irrigation. Plasticulture Development Centre, GBPUAT, Pantnagar. Journal of Geoscience and Environment Protection. 2001;5(7):22-30.
13. Srinivas K. Drip irrigation studies in mango for growth and yield. Ann. Rep., I.I.H.R., Bangalore. 2005;20-21.
14. Dixit A, Sharma D, Agrawal N, Dubey P. Effect of drip irrigation and mulch on mango yield and its fruits quality characters. Plant Archives. 2003;3(2):199-202.

15. Dinesh K, Pandey V, Vishal N. 2008. Effect of drip irrigation regimes on growth, yield and quality of mango hybrid Arka Anmol. Indian Journal Horticulture. 2008;65(4):409-412.
16. Shirgure PS, Srivastava AK, Singh Shyam, Pimple AR. Drip irrigation scheduling, growth, yield and quality of acid lime (*Citrus aurantifolia* Swingle). Indian Journal of agricultural Sciences. 2003;73:209-11.
17. Shirgure PS, Srivastava AK, Singh Shyam. Growth, yield and quality of acid lime under pan evaporation based drip irrigation scheduling. Indian Journal of Soil Conservation. 2004;32:32-35.
18. Pavel EW, de Villiers AJ. Response of mango trees to reduced irrigation regimes. International symposium on irrigation and water relation in grape vine and fruit trees. Acta Horticulture. 2004;648.
19. Coelho EF, Borges AL. Irrigation and fertigation in mango. International Symposium Brazil. Acta Horticulture. 2004;646:121-123.
20. Kumar D, Pandey V, Nath V. Effect of organic mulches and irrigation scheduling through drip on growth and yield of 'Lal Sundari' mango (*Mangifera indica*) in Eastern region of India. Indian Journal of Agricultural Sciences. 2008;78(5):385-388.
21. Sharma S, Patra SK, Ray R. Effect of drip fertigation on growth and yield of guava. Environment and Ecology. 2011;29(1-2):34-38.
22. Pradhan PC, Mishra JN, Panda SC, Bahera B. Technological feasibility and economical viability of drip irrigation and polyethylene mulch in mango (*Mangifera indica*). Indian Journal of Dry Land Agriculture Research and Development. 2010;25(2):91-94.
23. Joshi G, Singh PK, Singh SK, Srivastava PC. Effect of drip fertigation and mulching on water requirement, yield and economics of high density litchi. Progressive Horticulture. 2011;43(2):237-242.
24. Shirgure PS, Srivastava AK, Singh S. Growth, yield and quality of Nagpur mandarin (*Citrus reticulata* Blanco) in relation to irrigation and fertigation. Indian Journal of Agricultural Sciences. 2000;71(8):547-550.
25. Sulochanamma BN, Reddy TY, Reddy GS. Effect of basin and drip irrigation on growth, yield and water use efficiency in pomegranate Cv. Ganesh. Acta Horticulture. 2005;696:277-79.
26. Panigrahi HK, Narendra Agrawal R, Agrawal, Saket Dubey, Tiwari SP. Effect of drip irrigation and polythene mulch on the fruit yield and quality parameters of mango (*Mangifera indica* L.). Journal of Horticultural Sciences. 2010;5(2):140-143.
27. Srivastava PK, Parikh MM, Sawani NG, Raman S. Response of banana to drip irrigation, mulches and irrigation scheduling in South Gujarat. Agricultural Engineering Today. 1999;23:29-38.
28. Tiwari KN, Kumar M, Santosh DT, Singh VK, Maji MK, Karan AK. Influence of Drip Irrigation and Plastic Mulch on Yield of Sapota (*Achras zapota*) and Soil Nutrients. Irrigation and Drainage Engineering. 2014;3:116-117.
29. Neilsen GH, Hogue EJ, Forge T, Neilsen D. Mulches and biosolids affect vigour, yield and leaf nutrition of fertigated high density apple. Hort Science. 2003;38(1):41-45.
30. Barman P, Kanchan K. Srivastava, Tarun A. response of guava cv. Lalit to plastic mulch, fertigation and micro-nutrient spray under raised bed. Indian Journal of Agriculture and Allied Sciences. 2017;3(4):1-8.
31. Ramniwas, Kaushik RA, Sarolia DK, Pareek S, Singh V. Effect of irrigation and fertigation scheduling on growth and yield of guava (*Psidium guajava* L.) under meadow orcharding. African Journal of Agricultural Research. 2012;7(47):6350-6356.

© 2020 Thirupathaiah et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<http://www.sdiarticle4.com/review-history/63259>