



Appraising the Biochemical Responses in Wheat (*Triticum aestivum* L.) Seedlings under Various Seed Treatments

Sultan Singh ^{a*}, Virender Singh Mor ^a, Axay Bhuker ^a,
Digamber ^a, Rahul Kumar ^a and Gagandeep Singh ^a

^a Department of Seed Science and Technology, Chaudhary Charan Singh Haryana Agricultural University, Hisar-125004, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JEAI/2023/v45i112243

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/109238>

Original Research Article

Received: 03/09/2023

Accepted: 10/11/2023

Published: 16/11/2023

ABSTRACT

A study was conducted to analyse the effect of pesticides and biofertilizers on enzymatic activity of dehydrogenase, catalase, peroxidase and superoxide dismutase (SOD) in seedlings of wheat (*Triticum aestivum* L.) varieties (WH1105 and WH1124) and their old and fresh seed lots. This experiment was conducted in the laboratories at the Department of Seed Science and Technology, CCS Haryana Agricultural University in Hisar during 2020. A total of 16 treatment combinations were applied to the seeds in the experiment, including control, which were subsequently used for germination. The seedlings of the wheat cultivars were germinated in between paper at controlled temperature 20°C. 4 days old germinated seedlings were used for the analyses of the biochemical parameters. This finding revealed that the seedlings germinated from seeds treated with T₅-

*Corresponding author: E-mail: sultan.hau@gmail.com;

Azotobacter+PSB exhibited highest biochemical activity, followed by T₄-Phosphate Solubilizing Bacteria. The application of biofertilizers enhanced the dehydrogenase, catalase, peroxidase and superoxide dismutase enzyme activities, whereas the T₁-Chlorpyrifos reduced the biochemical activities of the wheat seedlings. The results demonstrated that Chlorpyrifos had an adverse effect on the biochemical activities and inhibiting seedlings optimal growth and inducing stress-related challenges for wheat seedlings.

Keywords: Biochemical; catalase; dehydrogenase; enzyme activity; peroxidase; superoxide dismutase; wheat.

1. INTRODUCTION

Plants are equipped with a robust antioxidant defense system, consisting of both enzymatic and non-enzymatic elements. Among the non-enzymatic components are carotenoids, ascorbate, glutathione, and tocopherols (Vitamin E). During various stress conditions, it has been observed that reactive oxygen species, including hydrogen peroxide (H₂O₂), superoxide (O²⁻), and hydroxyl radicals (OH), are produced. In concurrence with these, there are antioxidants enzymes like superoxide dismutase (SOD), catalase, and those involved in the ascorbate-glutathione cycle. SOD plays a fundamental role in this defense system by transforming two molecules of superoxide into a lesser amount of harmful substances, namely hydrogen peroxide (H₂O₂) and molecular oxygen (O₂) [1]. In plants, three distinct types of superoxide dismutase (SOD) enzymes (EC 1.15.1.1) have been identified, each characterized by its specific metal cofactor. These are Manganese SOD (MnSOD), which is located in the mitochondrial matrix. Copper/Zinc SOD (Cu/ZnSOD) primarily localized in the cytosolic. Iron SOD (FeSOD), which is localized in the chloroplastic stroma [2]. Additionally, some plants may also have Cu/Zn SOD in their chloroplasts. It's worth observing that Cu/ZnSOD is sensitive to both hydrogen peroxide (H₂O₂) and potassium cyanide (KCN), while FeSOD is responsive to H₂O₂ but not KCN. On the other hand, MnSOD shows insensitivity to both H₂O₂ and KCN, making it distinct from the other two SOD types [3]. Plant peroxidases have served as valuable biochemical indicators for different forms of biotic and abiotic stresses. This is attributed to their essential involvement in critical physiological processes, such as regulating growth through lignification, forming cross-links in pectins and structural proteins within cell walls, and participating in the breakdown of auxins [4]. Catalases and superoxide dismutases stand out as the most

effective antioxidant enzymes [5]. These enzymes include superoxide dismutase (SOD, EC 1.15.1.1) which reacts with superoxide radicals and converts them to O₂ and H₂O₂. H₂O₂ is then detoxified by catalase (CAT, EC 1.11.1.6). The endosperm of mature seeds contains over 250 proteins that play a role in 13 different biochemical processes, including reactions related to ATP inter-conversion, during germination [6]. As respiration becomes more intense during germination, there is a concurrent increase in the release of inorganic phosphorus and cellular energy. This heightened respiration plays a crucial role in generating the energy needed for the normal progression of metabolic processes. Following this phase, the synthesis of new proteins, nucleic acids, and lipids occurs, while the stored reserves serve as nutrients for the growing seedling [7]. This study is focused on assessing the impact of various environmental pollutants resulting from the extensive use of substances like insecticides and fungicides, and how the role played by certain biofertilizers in the plant development process. Therefore, the current research aims to investigate how Chlorpyrifos, Vitavax, *Azotobacter* and PSB, individually and in various combinations, influence enzymatic activity of dehydrogenase, catalase, peroxidase and superoxide dismutase of the wheat seedlings.

2. MATERIALS AND METHODS

This study was carried out during the year 2020 at the laboratories of the Department of Seed Science and Technology and the seed material were obtained from breeder seed store, Department of Seed Science and Technology, CCS Haryana Agricultural University, Hisar, Haryana, situated in the semi-tropical region in the north-western zone of India. For this study, wheat varieties WH1105 and WH1124 were taken, along with one-year-old harvested seeds and freshly harvested seed lots for each variety.

2.1 Seed Treatment

The seeds were primed with different treatments, including Chlorpyrifos 20EC (1.5 ml/kg of seeds), Vitavax (2g/kg of seeds), *Azotobacter* (5ml/kg of seeds), Phosphate solubilizing bacteria (PSB) (5ml/kg of seeds) and their combinations of total 16 treatments including control as shown in Table 1. To enhance the adhesion of biofertilizers to the seed surface, a 10 percent jaggery solution was used, followed by shade exposure for some time to the treated seeds. Afterward, the treated seeds were packaged in malleable zippered bags and used for sowing purpose.

Seedlings were grown from seeds in the between paper at controlled temperature at 20°C. In this study, enzymatic activity of dehydrogenase (DHA), catalase, peroxidase (POX) and superoxide dismutase (SOD) were studied on the 4th day seedlings during germination after the various treatments *i.e.*, Chlorpyrifos, vitavax, *Azotobacter* and PSB with their combinations. The germinated seedlings were used for enzyme extraction.

2.2 Tissue Extraction

The steps, which involve the extraction process, were carried out at a temperature range of 0-4°C. The 1 g of plant tissue from 4th day germinated seedlings were finely ground in a chilled mortar and pestle while being mixed with 5 ml of 0.1 M phosphate buffer with a pH of 7.5. Three sets of biological samples (leaf tissue) were collected to create enzyme extracts. This buffer solution also contained 5% (w/v) polyvinylpyrrolidone (PVP), 10mM β -mercaptoethanol, and 1mM EDTA. The resulting homogenate was subjected to centrifugation using a centrifuge operating at 10,000 rpm for 20 minutes at 4°C. Following the centrifugation

process, the liquid portion, known as the supernatant, was carefully separated and utilized as a crude enzyme preparation, following the method outlined by Sinha [8]. These extracts were used for all the enzyme assays except dehydrogenase (DHA).

2.3 Determination of the Dehydrogenase Activity (O D g⁻¹ ml⁻¹)

The Dehydrogenase activity (DHA) test was conducted following the method outlined by Kittock and Law [9]. Each seedling lot, represented by one gram of seedlings and replicated thrice, was ground properly. The resulting 200 mg of extract was soaked in 5 ml of a 0.5% tetrazolium solution at 38°C for 3 hours. Afterward, it was centrifuged at 10,000 rpm for 3 minutes, and the supernatant was carefully decanted. The formazan, a red-colored compound, was extracted from the mixture using 10 ml of acetone over a 16-hour period. Following this, another round of centrifugation was performed, and the absorbance of the resulting solution was measured at 480 nm using an Eppendorf Biospectrometer. These observations were expressed as changes in optical density per gram per milliliter (OD g⁻¹ ml⁻¹).

2.4 Determination of the Catalase activity (CAT)

In this process of Catalase activity (CAT) (EC 1.11.1.6), a mixture was created by combining 0.55 ml of 0.1 M potassium phosphate buffer at pH 7.0, 0.4 ml of 0.2 M H₂O₂, and 50 μ l of enzyme extract. This mixture was thoroughly mixed and incubated for one minute. Subsequently, 3.0 ml of a 5% potassium dichromate: acetic acid (1:3) solution was added. A control sample was also prepared, which included 0.6 ml of assay buffer and

Table 1. Various treatments and their combinations

S. No.	Treatments	S. No.	Treatments
T ₀	Control	T ₈	Vitavax+ <i>Azotobacter</i>
T ₁	Chlorpyrifos 20EC	T ₉	Vitavax+ <i>Azotobacter</i> +PSB
T ₂	Vitavax	T ₁₀	Chlorpyrifos+ <i>Azotobacter</i>
T ₃	<i>Azotobacter</i>	T ₁₁	Chloropyrifos+PSB
T ₄	Phosphate Solubilizing Bacteria (PSB)	T ₁₂	Chlorpyrifos+ <i>Azotobacter</i> +PSB
T ₅	<i>Azotobacter</i> +PSB	T ₁₃	Chlorpyrifos+Vitavax+ <i>Azotobacter</i>
T ₆	Vitavax+Chlorpyrifos	T ₁₄	Chlorpyrifos+Vitavax+PSB
T ₇	Vitavax+PSB	T ₁₅	Chloropyrifos+Vitavax+ <i>Azotobacter</i> +PSB

0.4 ml of 0.2 M H₂O₂, but without the enzyme extract, alongside the test samples. The tubes containing these mixtures were then placed in a boiling water bath for 10 minutes, allowed to cool, and their absorbance was measured at 570 nm using the dichromate: acetate solution as a reference. Calculations were carried out by subtracting the absorbance of the samples from that of the control, and the quantity of H₂O₂ was determined using a standard curve. One catalase (CAT) unit is defined as the amount of enzyme required to decompose one μ mol of H₂O₂ per minute or per milligram of protein.

2.5 Determination of the Peroxidase activity (POX)

In Peroxidase activity (POX) (EC 1.11.1.7) a glass cuvette was filled with 2.15 ml of 0.1 M Tris-HCl buffer at pH 7.0 was carefully measured. To this, 0.6 ml of a 1% guaiacol solution and 0.1 ml of the enzyme extract were added. The solution was thoroughly mixed and adjusted to achieve 100% light transmission at 470 nm. Then, 0.15 ml of 100 mM H₂O₂ was introduced and mixed, and then the initial reading was taken at 470 nm. Subsequently, the increase in absorbance was recorded at 15-second intervals for duration of 3 minutes. The enzyme activity was calculated using the linear segment of the optical density (OD) change, utilizing a molar extinction coefficient of 26.6 mM⁻¹ cm⁻¹ for guaiacol oxidation. One unit of peroxidase (POX) activity is defined as the amount of enzyme needed to oxidize one nmol of guaiacol per minute, per milliliter, or per milligram of protein.

2.6 Determination of the SOD Activity

To measure superoxide dismutase (SOD, EC 1.15.1.1) (SOD) activity, the method employed involved evaluating its capacity to impede the photochemical reactions involving nitro blue tetrazolium (NBT), following the procedure outlined by Beauchamp and Fridovich [10]. In this procedure, a reaction mixture of 3.0 ml was prepared, consisting of 2.5 ml of 60 mM Tris-HCl at pH 7.8, along with 0.1 ml each of 420 mM L-methionine, 1.80 mM NBT, 90 μ M riboflavin, 3.0 mM EDTA, and 0.1 ml of the enzyme extract. Riboflavin was introduced at the end of the preparation. The tubes were thoroughly shaken and positioned 30 cm below a light source comprising three 20 W fluorescent lamps. The reaction was initiated by switching on the light and halted after a 40-minute incubation period by turning off the light source. To shield the tubes

from further light exposure, they were covered with a black cloth. A control was maintained, consisting of a non-irradiated reaction mixture that did not exhibit any color change. The maximum color development was observed in the reaction mixture without the enzyme extract. The absorbance was measured at 560 nm, and one enzyme unit was defined as the amount of enzyme required to inhibit the photo-reduction of one μ mol of NBT. To express the enzyme activity in terms of units per gram of fresh weight, it was calculated using the standard formula by Giannopolitis and Ries [11], which is commonly used for assessing kinetic and regulatory properties.

$$\text{SOD units} = R - r \text{ divided by } r$$

Where,

R = Rate of assay reaction in absence of SOD.

r = Rate of assay reaction in presence of SOD.

However, calculate the percent inhibition by the following formula of Asada et al. [12]:

$$\text{Per cent inhibition} = R - r \text{ divided by } R \times 100$$

2.7 Statistical Analysis

The final data attained was subjected to the analysis using online statistical analysis tool OPSTAT by Sheoran [13].

3. RESULTS AND DISCUSSION

This study focuses on evaluating the effects of different seed treatments, both when applied individually and in various combinations like *Azotobacter*, PSB, Chlorpyrifos, and Vitavax, on the biochemical activities of wheat seedlings. The dehydrogenase activity of seedlings, germinated from treated seeds is presented in Table 2 of wheat in both the lots old and fresh of WH1105 and WH1124. Dehydrogenase (DHA) activity was enhanced with treatment of biofertilizers, the activity indicated as (optical density) OD value (O.D. g⁻¹ ml⁻¹). The overall highest DHA activity was recorded in treatment T₅-*Azotobacter*+PSB (0.250), followed by T₄-PSB (0.238) and lowest was observed in T₁-Chlorpyrifos (0.183) as compared to T₀-control (0.200). The mean performance includes the both lots of the variety WH1105, the maximum enhancement of DHA activity was observed in treatment T₅-*Azotobacter*+PSB (0.248), followed

by T₄-PSB (0.237) and lowest was observed in T₁-Chlorpyrifos (0.181) as compared to T₀-control (0.199). A same performance was recorded in both the lots of the variety WH1124, the maximum DHA activity was observed in treatment T₅-*Azotobacter*+PSB (0.251), followed by T₄-PSB (0.239), while the lowest observed T₁-Chlorpyrifos (0.185) in comparison of T₀-control (0.202).

Table 3 displays the catalase activities of seedlings that have germinated from treated seeds reported in $\mu\text{moles g}^{-1}$ fresh weight of wheat in old and fresh lots of WH1105 and WH1124. The overall maximum catalase activity was observed in treatment T₅-*Azotobacter*+PSB (5637), followed by T₄-PSB (5400) and minimum was recorded in T₁-Chlorpyrifos (4305) as compared to T₀-control (4430). The mean observations involves the both the lots of WH1105, the maximum catalase activity in treatment T₅-*Azotobacter*+PSB (5524), followed by T₄-PSB (5300) and lowest was observed in T₁-Chlorpyrifos (4257) as compared to T₀-control (4420). The same findings were observed in both the lots of the variety WH1124, the maximum catalase activity was recorded in treatment T₅-*Azotobacter*+PSB (5750), followed by T₄-PSB

(5501), while the minimum observed T₁-Chlorpyrifos (4353) in comparison of T₀-control (4439). These findings confirms by Singh et al. [14] Catalase (CAT) activity showed an increase in maize leaves when they were inoculated with *Azotobacter chroococcum* and the combination of *Azotobacter chroococcum* with *Bacillus polymyxa*. Seyed Sharifi et al. [15] revealed that the application of biofertilizers and nano-oxides enhanced the proline content, soluble sugars, catalase, peroxidase, and polyphenol oxidase enzyme activities of wheat.

Table 4 comprises the peroxidase activities (POX) of seedlings that have germinated from the various treatment combinations, presented as $\mu\text{moles per gram of fresh weight}$, in both old and fresh seed lots of WH1105 and WH1124 wheat varieties. The overall maximum peroxidase activity was observed in T₅-*Azotobacter*+PSB (31.71) followed by T₄-PSB (29.21) whereas the minimum peroxidase activity was observed in T₁-Chlorpyrifos (11.26) as compared to T₀-control (13.53). The mean results includes the lots of WH1105, the maximum peroxidase activity in treatment T₅-*Azotobacter*+PSB (29.84), followed by T₄-PSB (27.51) and lowest was observed in

Table 2. Effect of various treatments on dehydrogenase activity (DHA) (O.D. $\text{g}^{-1} \text{ml}^{-1}$) in wheat seedlings

S. No.	Variety(V)						Overall Mean
	WH1105			WH1124			
	Old	Fresh	Mean	Old	Fresh	Mean	
T ₀	0.195	0.202	0.199	0.198	0.206	0.202	0.200
T ₁	0.177	0.185	0.181	0.181	0.188	0.185	0.183
T ₂	0.203	0.210	0.207	0.207	0.214	0.211	0.209
T ₃	0.231	0.236	0.234	0.233	0.237	0.235	0.234
T ₄	0.234	0.239	0.237	0.236	0.241	0.239	0.238
T ₅	0.243	0.253	0.248	0.246	0.256	0.251	0.250
T ₆	0.205	0.213	0.209	0.209	0.218	0.214	0.211
T ₇	0.211	0.217	0.214	0.214	0.220	0.217	0.216
T ₈	0.208	0.214	0.211	0.209	0.218	0.214	0.212
T ₉	0.224	0.228	0.226	0.227	0.231	0.229	0.228
T ₁₀	0.197	0.204	0.201	0.203	0.206	0.205	0.203
T ₁₁	0.209	0.215	0.212	0.212	0.219	0.216	0.214
T ₁₂	0.214	0.221	0.218	0.218	0.226	0.222	0.220
T ₁₃	0.181	0.187	0.184	0.183	0.193	0.188	0.186
T ₁₄	0.201	0.208	0.205	0.205	0.211	0.208	0.206
T ₁₅	0.184	0.190	0.187	0.186	0.194	0.190	0.189
Mean	0.207	0.214		0.210	0.217		
	V	L	T	VxL	VxT	LxT	VxLxT
C.D. (P=0.05)	0	0	0.001	NS	NS	0.001	NS
SE.m (\pm)	0	0	0	0	0.001	0.001	0.001

Table 3. Effect of various treatments on catalase activity ($\mu\text{moles g}^{-1}$ fresh weight) in wheat seedlings

S. No.	Variety(V)						Overall Mean
	WH1105			WH1124			
	Old	Fresh	Mean	Old	Fresh	Mean	
T ₀	4,392	4,448	4,420	4,405	4,473	4,439	4,430
T ₁	4,194	4,320	4,257	4,277	4,428	4,353	4,305
T ₂	4,442	4,630	4,536	4,474	4,726	4,600	4,568
T ₃	4,927	5,320	5,123	5,060	5,491	5,276	5,200
T ₄	5,039	5,562	5,300	5,226	5,775	5,501	5,400
T ₅	5,282	5,767	5,524	5,553	5,947	5,750	5,637
T ₆	4,467	4,706	4,586	4,505	4,822	4,664	4,625
T ₇	4,642	4,985	4,814	4,840	5,043	4,942	4,878
T ₈	4,495	4,808	4,652	4,677	4,935	4,806	4,729
T ₉	4,838	5,174	5,006	4,951	5,329	5,140	5,073
T ₁₀	4,401	4,460	4,430	4,424	4,491	4,458	4,444
T ₁₁	4,532	4,911	4,722	4,715	4,989	4,852	4,787
T ₁₂	4,723	5,036	4,880	4,893	5,135	5,014	4,947
T ₁₃	4,245	4,357	4,301	4,311	4,432	4,372	4,336
T ₁₄	4,403	4,518	4,461	4,438	4,591	4,515	4,488
T ₁₅	4,311	4,406	4,359	4,380	4,437	4,409	4,384
Mean	4,583	4,838		4,696	4,940		
	V	L	T	VxL	VxT	LxT	VxLxT
C.D. (P=0.05)	3.53	3.53	9.98	4.99	14.11	14.11	19.95
SE.m (\pm)	1.26	1.26	3.57	1.78	5.04	5.04	7.13

Table 4. Effect of various treatments on peroxidase activity ($\mu\text{moles g}^{-1}$ fresh weight) in wheat seedlings

S. No.	Variety(V)						Overall Mean
	WH1105			WH1124			
	Old	Fresh	Mean	Old	Fresh	Mean	
T ₀	9.66	14.04	11.85	12.14	18.27	15.21	13.53
T ₁	7.84	11.32	9.58	10.50	15.37	12.94	11.26
T ₂	12.14	19.19	15.67	15.24	22.14	18.69	17.18
T ₃	20.20	30.47	25.33	24.27	33.36	28.82	27.07
T ₄	22.46	32.55	27.51	26.34	35.49	30.92	29.21
T ₅	24.47	35.22	29.84	28.77	38.37	33.57	31.71
T ₆	12.95	21.23	17.09	16.84	23.56	20.20	18.65
T ₇	15.41	25.31	20.36	19.32	27.87	23.60	21.98
T ₈	13.43	22.76	18.10	18.22	25.20	21.71	19.90
T ₉	17.42	28.56	22.99	22.25	31.47	26.86	24.92
T ₁₀	9.69	15.42	12.55	13.18	19.35	16.26	14.41
T ₁₁	14.18	23.88	19.03	18.87	26.26	22.57	20.80
T ₁₂	16.18	27.31	21.75	20.23	29.15	24.69	23.22
T ₁₃	8.37	12.31	10.34	11.09	16.70	13.90	12.12
T ₁₄	10.17	18.13	14.15	13.44	21.23	17.34	15.74
T ₁₅	9.06	13.07	11.07	11.93	17.20	14.57	12.82
Mean	13.98	21.92		17.66	25.06		
	V	L	T	VxL	VxT	LxT	VxLxT
C.D. (P=0.05)	0.041	0.041	0.117	0.058	0.165	0.165	0.233
SE.m (\pm)	0.015	0.015	0.042	0.021	0.059	0.059	0.083

T₁-Chlorpyrifos (9.58) as compared to T₀-control (11.85). The similar result was observed in both the lots of WH1124, the highest peroxidase

activity in treatment T₅-Azotobacter+PSB (33.57), followed by T₄-PSB (30.92) and lowest was observed in T₁-Chlorpyrifos (12.94) as compared

Table 5. Effect of various treatments on superoxide dismutase activity ($\mu\text{moles g}^{-1}$ fresh weight) in wheat seedlings

S. No.	Variety(V)						Overall Mean
	WH1105			WH1124			
	Old	Fresh	Mean	Old	Fresh	Mean	
T ₀	90.01	93.95	91.98	91.24	98.19	94.71	93.35
T ₁	87.73	90.06	88.90	88.09	91.16	89.63	89.26
T ₂	91.70	97.15	94.42	94.84	99.87	97.35	95.89
T ₃	102.33	108.38	105.36	107.48	112.55	110.02	107.69
T ₄	103.55	110.83	107.19	108.52	115.46	111.99	109.59
T ₅	106.81	113.61	110.21	110.95	117.55	114.25	112.23
T ₆	93.63	99.15	96.39	95.22	101.16	98.19	97.29
T ₇	97.07	103.46	100.27	101.22	105.22	103.22	101.75
T ₈	95.07	101.17	98.12	97.25	101.91	99.58	98.85
T ₉	99.77	105.52	102.65	105.23	109.54	107.39	105.02
T ₁₀	90.92	94.45	92.69	92.85	95.20	94.03	93.36
T ₁₁	95.83	102.25	99.04	99.23	102.48	100.86	99.95
T ₁₂	98.17	104.34	101.25	103.38	107.31	105.34	103.30
T ₁₃	88.67	91.35	90.01	89.27	92.10	90.69	90.35
T ₁₄	91.35	95.82	93.59	93.37	98.19	95.78	94.68
T ₁₅	89.23	92.17	90.70	90.09	93.07	91.58	91.14
Mean	95.12	100.23		98.01	102.56		
	V	L	T	VxL	VxT	LxT	VxLxT
C.D. (P=0.05)	0.054	0.054	0.152	0.076	0.215	0.215	0.304
SE.m (\pm)	0.019	0.019	0.054	0.027	0.077	0.077	0.109

to T₀-control (15.21). Several reports confirms the results in terms of biochemical enzyme activities, the use of a double full dose of organic manure, in combination with the *Azotobacter* and *Pseudomonas* inoculum, resulted in an increase in both peroxidase and catalase activities. These findings in sorghum supported by Amal et al. [16]. These results also confirmed in wheat by Babaei et al. [17].

The data shown in Table 5 presents the superoxide dismutase (SOD) activities in germinated seedlings using various treatments, measured in $\mu\text{moles per gram}$ of fresh weight. It specifically focuses on superoxide dismutase activities in both old and fresh seed lots of WH1105 and WH1124 wheat varieties. The overall mean performance superoxide dismutase activities (SOD) was recorded highest in T₅-*Azotobacter*+PSB (112.23) treatment, followed by T₄-Phosphate Solubilizing Bacteria (109.59) and minimum in T₁-Chlorpyrifos (89.26) in comparison of T₀-control (93.35). The mean performance of superoxide dismutase activities of both the lots of WH1105, the maximum SOD activity was observed in treatment T₅-*Azotobacter*+PSB (110.21) which was at par with T₄-PSB (107.19) and minimum in T₁-Chlorpyrifos (88.90) as compared to T₀-control (91.98). A same performance was recorded in both the lots

of the WH1124, the maximum superoxide dismutase activity was observed in treatment T₅-*Azotobacter*+PSB (114.25), followed by T₄-PSB (111.99), while the lowest observed T₁-Chlorpyrifos (89.63) as compared to T₀-control (94.71). The fresh seed lot of WH1124 recorded the highest enzymatic activities among other lots. In maize, the presence of microbes such as, *Serendipita indica*, *Rhizophagus intraradices*, and *Azotobacter chroococcum* led to increase in the activities of antioxidant enzymes, including peroxidase (POX), catalase (CAT), polyphenol oxidase (PPO), and superoxide dismutase (SOD), when compared to control plants, these results were supported by Tyagi et al. [18]. *Paenibacillus* sp.+*Bacillus subtilis* (PGPR) (biofertilizers) showed a maximum increase in superoxidase dismutase (SOD), peroxidase dismutase (POD), catalase (CAT) in wheat. These results confirmed by Iqbal et al. [19]. It has been reported by Srivalli [20] that the activity of superoxide dismutase and peroxidase was recorded higher as compared to the control depending on plant species, tissue and stage of development. The present study revealed that the combined application of biofertilizers enhanced the biochemical activities of wheat seedlings. The maximum enzymatic activity of dehydrogenase, catalase, peroxidase and superoxide dismutase was observed with the

treatment T₅-Azotobacter+PSB. The increasing amount of chemical pesticides and fertilizers has had a negative impact on the ecosystem. To minimize this, we should be more and more aware of the need to reduce their dependence on chemical fertilizers. Instead of chemicals the efforts should be focused on utilizing the biofertilizers and increase biological activity of microorganisms in the rhizosphere to increase soil fertility and promote crop growth.

4. CONCLUSION

The combined application of the biofertilizers recommended for high quality and profitable wheat production. Seedlings germinated from seeds treated with T₅-Azotobacter+PSB exhibited highest biochemical activity. The application of biofertilizers enhanced the dehydrogenase, catalase, peroxidase and superoxide dismutase enzyme activities, whereas the T₁-Chlorpyrifos reduces the biochemical activities of the wheat seedling. The rise in enzyme activity leads to enhanced stress tolerance in wheat seedlings. This coordinated defense mechanism assists the plants in sustaining their growth and recuperating from stress related challenges.

ACKNOWLEDGEMENTS

The authors would like to thank CCS Haryana Agricultural University, Hisar for providing necessary facilities to conduct this study.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Ceron-Garcia A, Vargas-Arispuro I, Aispuro-Hernandez E, Martinez-Tellez MA. Oligoglucan elicitor effects during plant oxidative stress. Cell Metabolism. In Cell Homeostasis and Stress Response; 2012.
2. Bowler C, Montagu MV, Inze D. Superoxide dismutase and stress tolerance. Annual Review of Plant Biology. 1992;1:83-116.
3. Beauchamp C, Fridovich I. Isozymes of superoxide dismutase from wheat germ. Biochimica et Biophysica Acta (BBA)-Protein Structure. 1973;317(1):50-64.
4. Gaspar T, Penel C, Thorpe T, Greppin H. Peroxidases 1970-1980. A survey of their biochemical and physiological roles in higher plants. Peroxidases 1970-1980. A survey of their biochemical and physiological roles in higher plants; 1982.
5. Scandalios JG, Tsaftaris AS, Chandlee JM, Skadsen RW. Expression of the developmentally regulated catalase (Cat) genes in maize. Developmental Genetics. 1983;4(4):281-93.
6. Weitbrecht K, Müller K, Leubner-Metzger G. First off the mark: Early seed germination. Journal of Experimental Botany. 2011;62(10):3289-309.
7. Burzo I, Toma S, Craciun C, Voican V, Dobrescu A, Delian E. Fiziologia plantelor de cultura, Volum I - Procesele fiziologice din plantele de cultura, Chisinau: Întrep. Editorial - Poligrafica Stiinta; 1999.
8. Sinha AK. Colorimetric assay of catalase. Analytical Biochemistry. 1972;47(2):389-94.
9. Kittock DL, Law AG. Relationship of seedling vigor to respiration and tetrazolium chloride reduction by germinating wheat seeds 1. Agronomy Journal. 1968;60(3):286-8.
10. Beauchamp C, Fridovich I. Superoxide dismutase: improved assays and an assay applicable to acrylamide gels. Analytical Biochemistry. 1971;44(1):276-87.
11. Giannopolitis CN, Ries SK. Superoxide dismutases: I. Occurrence in higher plants. Plant Physiology. 1977;59(2):309-14.
12. Asada K, Takahashi MA, Nagate M. Assay and inhibitors of spinach superoxide dismutase. Agricultural and Biological Chemistry. 1974;38(2):471-3.
13. Sheoran OP. Online statistical analysis (OPSTAT) software developed by Chaudhary Charan Singh Haryana Agricultural University, Hisar, India. 2010. Available:<http://14.139.232.166/opstat/>
14. Singh NB, Singh D, Singh A. Biological seed priming mitigates the effects of water stress in sunflower seedlings. Physiology and Molecular Biology of Plants. 2015; 21:207-14.
15. Seyed Sharifi R, Khalilzadeh R, Pirzad A, Anwar S. Effects of biofertilizers and nano zinc-iron oxide on yield and physicochemical properties of wheat under water deficit conditions. Communications in Soil Science and Plant Analysis. 2020; 51(19):2511-24.
16. Amal GA, Orabi S, Gomaa AM. Bio-organic farming of grain sorghum and its effect on growth, physiological and yield

- parameters and antioxidant enzymes activity. Research Journal of Agriculture and Biological Sciences. 2010;6(3): 270-9.
17. Babaei K, Seyed Sharifi R, Pirzad A, Khalilzadeh R. Effects of bio fertilizer and nano Zn-Fe oxide on physiological traits, antioxidant enzymes activity and yield of wheat (*Triticum aestivum* L.) under salinity stress. Journal of Plant Interactions. 2017; 12(1):381-9.
 18. Tyagi J, Mishra A, Kumari S, Singh S, Agarwal H, Pudake RN, et al. Deploying a microbial consortium of *Serendipita indica*, *Rhizophagus intraradices*, and *Azotobacter chroococcum* to boost drought tolerance in maize. Environmental and Experimental Botany. 2023;206:105142.
 19. Iqbal Z, Bushra, Hussain A, Dar A, Ahmad M, Wang X, et al. Combined use of novel endophytic and rhizobacterial strains upregulates antioxidant enzyme systems and mineral accumulation in wheat. Agronomy. 2022;12(3):551.
 20. Srivalli B, Sharma G, Khanna-Chopra R. Antioxidative defense system in an upland rice cultivar subjected to increasing intensity of water stress followed by recovery. Physiologia Plantarum. 2003; 119(4):503-12.

© 2023 Singh et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/109238>