



Cardiac Tamponade: An Unusual First Case Presentation of Systemic Lupus Erythematosus

**Dghoughi Sarah ^{a*}, Jama Dounia ^a, Charif Hanae ^a,
Mounaouir Karim ^a, Bouziane Maha ^b, Haboub Meryem ^b,
Bennouna El Ghali ^b, Drighil Abdenasser ^b
and Habbal Rachida ^b**

^a Department of cardiology, Ibn Rochd University Hospital, Casablanca, Morocco.

^b Department of Cardiology, Faculty of Medicine and Pharmacy, Ibn Rochd University Hospital,
Casablanca, Hassan II University of Casablanca, Casablanca, Morocco.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJMPCR/2024/v17i2374

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/115531>

Case Report

Received: 08/02/2024

Accepted: 15/04/2024

Published: 19/04/2024

ABSTRACT

This case is reported for the rarity of cardiac tamponade as a first manifestation of systemic lupus erythematosus (SLE). Systemic lupus erythematosus (SLE) is a multisystem autoimmune disease with worldwide prevalence, affecting almost every organ in the body. A 27-year-old woman presented to our hospital after two months of progressive shortness of breath, generalized edematous syndrome and diffuse, non-migratory arthralgias; the patient's shortness of breath was exacerbated on exertion and associated with a vague sensation of chest tightness. Transthoracic echocardiography revealed a large circumferential pericardial effusion with an echo-free space of

*Corresponding author: E-mail: sarah.dghoughi2@gmail.com;

34 mm in apical view and 36 mm in subcostal view with an oscillating heart and diastolic collapse of the right ventricle and atrium. This case is reported for the rarity of cardiac tamponade as a first manifestation of SLE. Because pericarditis is more common in SLE patients, the low frequency of tamponade may be partially explained by the extensive use of non-steroidal anti-inflammatory medications (NSAIDs) and steroids, which effectively reduce pericardial inflammation. We emphasize the importance of including SLE in the differential diagnosis of patients presenting with cardiovascular symptoms, as cardiac tamponade and pleural effusion are uncommon early presentations of SLE. Early detection and diagnosis are essential for prompt intervention and better patient outcomes.

Keywords: Systemic lupus erythematosus; cardiac tamponade; echocardiography; pericardial effusion.

1. INTRODUCTION

“Systemic lupus erythematosus (SLE) is a multisystem autoimmune disease with worldwide prevalence that affects almost all organs of the body” [1]. “The disease is more prevalent among younger women, with a female-to-male prevalence ratio of approximately 10:1” [2]. “Cardiac involvement is common and can affect the pericardium, myocardium, valvular structures, and the conduction system, pericardial involvement is the most common cardiac manifestation of SLE” [3,4]. The progression to developing cardiac tamponade is very rare, with an incidence between 1% and 3% [5], and as the first presentation, it is rarer [6], it is a potentially fatal complication if not treated promptly. In this case report, we describe a 27-year-old Moroccan woman whose initial manifestation of SLE was cardiac tamponade. This case is reported for the rarity of cardiac tamponade as the first manifestation of SLE.

2. CASE PRESENTATION

A 27-year-old female presented to our hospital with a two-month history of progressive breathlessness, generalized edematous

syndrome and diffuse, non-migratory arthralgias, the patient's shortness of breath was exacerbated with exertion and associated with vague chest tightness. She had been on symptomatic treatment for psoriasis for 8 years and had been taking no medication for a year before the presentation. “She had no history suggestive of heart failure, renal failure, or liver disease. She had no hair loss, oral ulcers, or rashes. She was not a known patient with diabetes mellitus and/or hypertension. She doesn't drink alcohol or smoke cigarettes” [7].

On examination, the patient's vital signs showed a blood pressure of 90/70 mmHg, a heart rate of 135 beats per minute, a temperature of 36,6°, and oxygen saturation of 98% on room air. The patient was ill-appearing and was in moderate distress. She had facial puffiness, generalized edema. On cardiovascular examination, the heart sounds were muffled, and the jugular veins were distended. A respiratory examination revealed decreased breath sounds at the right lung base.

The electrocardiogram showed sinus tachycardia, low voltages and electrical alternans (Fig. 1). The chest radiograph showed an increased cardiac silhouette (Fig. 2).




Fig. 1. EKG demonstrating sinus tachycardia with low voltage QRS and electrical alternans




Fig. 2. Chest X-ray shows an enlarged cardiac silhouette and small right pleural effusion



Fig. 3. n (A) Transthoracic 2-dimensional echocardiogram (apical view) showing a large pericardial effusion (PE) and collapse of the right atrium (orange arrow). (B) Transthoracic 2-dimensional echocardiogram (subcostal view) showing large pericardial effusion opposite the right atrium (green arrow) and posterior wall of the left ventricle (blue arrow)




Fig. 4. Significant inspiratory decrease in the mitral and tricuspid valves E wave velocity on pulse wave Doppler tracing




Fig. 5. CT scan shows pericardial effusion and small right pleural effusion



Fig. 6. Favorable clinical response with a significant reduction in systemic congestion after initiating the treatment

The transthoracic echocardiography revealed a large circumferential pericardial effusion (PE) with an echo-free space of 34 mm in the apical view and 36 mm in the subcoastal view measured at end-diastole with swinging heart and diastolic collapse of the right atrium (Fig. 3) and the right ventricle. There was also a significant inspiratory decrease in the mitral and the tricuspid valves E wave velocity on pulse wave Doppler tracing, 44% and 61% respectively

(Fig. 4) and an IVC dilated to 26 mm non-complying, which confirmed a tamponade.

The patient was initially admitted to our intensive care unit in a half-seated position, given a saline infusion, and had 1700 ml of yellow-citrine fluid removed during pericardiocentesis. For the pericardial effusion, we have begun symptomatic treatment with colchicine 0.5 mg daily and aspirin 3g daily along with a proton pump inhibitor. The

evolution was marked in our patient by the recurrence of a large posterior effusion with signs of tamponade. Given the rapid re-accumulation of pericardial fluid, recurrent episodes of tamponade, and the location of the effusion, predominantly opposite the posterior wall of the LV, a surgical drainage with a pleuro-pericardial window was indicated. The pericardial biopsy showed inflammatory fibrous pericarditis and no malignancy. A biochemical, cytological and bacteriological study was carried out on the punctured fluid, finding an exudative fluid with lymphocytic predominance (90%) and no germs, the culture was sterile and tuberculosis PCR was negative. Complete blood count showed a normochromic normocytic anemia (hemoglobin 10.1 gr/dl), thrombocytopenia at 120000/mm³ and leukopenia at 1230/mm³ with an inflammatory syndrome with a CRP of 209 mg/l the erythrocyte sedimentation rate was 55 mm and hyperfibrinogenemia at 5.95 g/l. Additional laboratory testing was done to further determine the cause of the pericardial effusion. Anti-double stranded DNA (anti-dsDNA) was positive, as were antinuclear antibodies (ANA) at a titer 1280. Complement C3 of 0,45 g/L, and complement C4 <0,06g/l. Positive proteinuria at 0.53 g/24h. Direct Coombs testing was positive. TSH us and blood urea nitrogen (BUN) was within the normal range. Additional tests revealed an elevation in serum complement levels and cancer antigen 125 (CA-125), but other malignant markers are negative. Serologic tests for hepatitis C virus (HCV), hepatitis B surface antigen (HBsAg), and human immunodeficiency virus (HIV). A CT scan revealed a large pericardial effusion associated with a small right pleural effusion (Fig. 5).

The patient had a score of 39 according to the 2019 European Alliance of Associations for Rheumatology/American College of Rheumatology (EULAR/ACR) classification criteria for SLE, confirming the diagnosis of SLE. These included low levels of C4 and C3, affirmative anti-dsDNA antibodies, proteinuria > 0.5g/24h; tenderness in at least two joints; pericardial effusion and acute pericarditis, thrombocytopenia and autoimmune hemolysis. The patient was started on methylprednisolone at a dose of 1 mg/kg/day. She was also administered hydroxychloroquine orally 200mgx2/day. She had a favorable clinical response with a significant reduction in systemic congestion (Fig. 6) as well as biological with a CRP at 10 mg/l vs 164 and a 24h proteinuria at 0.16 vs 0.53 g/24h and also regression of the pericardial effusion. She is currently undergoing

outpatient follow-up in the rheumatology and cardiology clinics.

3. DISCUSSION

“Our patient had severe serositis as an initial presentation of SLE that presented as cardiac tamponade and small pleural effusion. Cardiac tamponade is a pericardial syndrome known as medical emergency characterised by impairment of the diastolic filling of the ventricles causing reduction of cardiac output, and if not treated promptly, can lead to cardiac arrest” [4,8]. “All the causes of pericardial effusion are also possible causes of cardiac tamponade, but the main causes of cardiac tamponade are: cardiac interventions, cancer, infections/inflammatory causes, mechanical complications of myocardial infarction, aortic dissection” [9]. “The rate and acuity at which these fluids accumulate within the pericardial space are the primary drivers for the development of tamponade physiology” [4]. “The heart is frequently involved in SLE. Approximately half of SLE patients will experience cardiac complications” [3]. “Cardiac tamponade as the first sign of SLE is very rare and only limited to case reports and series” [5]. “In 1949, Curtis and Horne described the first confirmed cases of lupus erythematosus with pericardial effusion as the main manifestation” [10]. The diagnosis of cardiac tamponade is a clinical one based on the combination of a suggestive history and symptoms (typically dyspnea, orthopnea, chest pain), associated with pulsus paradoxus, jugular venous distension, hypotension and decreased heart sounds, confirmed by echocardiography [5-9], the core echocardiographic findings of pericardial tamponade are pericardial effusion, signs of right atrial (sensitive) and/or right ventricular compression (specific) [11], abnormal respiratory variation in right and left ventricular dimensions and in mitral and tricuspid valve flow velocities, it is significant when respiratory changes of the mitral E velocity are above 25%, and tricuspid E velocity are above 40%, usually associated with inferior vena cava plethora [5-9]. “The low incidence of tamponade despite the high prevalence of pericarditis in SLE may be attributable in part to the widespread use of steroids and nonsteroidal anti-inflammatory drugs (NSAIDs), which effectively reduce pericardial inflammation” [5]. “Several studies have been carried out in an attempt to determine which patients are likely to develop cardiac tamponade. One retrospective study showed that patients with pericardial effusions who developed

tamponade had statistically lower C4 complement levels than those who did not develop physiological tamponade” [12]. “Moreover, A retrospective study showed that the presence of pleuritis and anti-nucleosome antibody positivity are significant predictors of progression to cardiac tamponade in SLE patients” [13]. “In treating known cardiac tamponade, efforts to withdraw the pericardial fluid by pericardiocentesis are the treatment of choice” [14]. The diagnosis of SLE, in our patient, was established based on positive clinical and immunological findings. According to the 2019 classification “criteria of the European League Against Rheumatism (EULAR)”. “Generally, the prognosis of lupus serositis is good; it responds well to corticosteroids and immunosuppressants. However, relapse and progression to fibrosis are uncommon” [15]. We highlight the importance of including SLE in the differential diagnosis of patients who present with cardiovascular symptoms because cardiac tamponade and pleural effusion are uncommon early presentations of SLE. For prompt intervention and improved patient outcomes, early detection and diagnosis are essential.

4. CONCLUSION

SLE affects any organ or system, and its onset ranges from non-specific symptoms to uncommon conditions such as cardiac tamponade, which is a potentially fatal condition and can lead to admission to the emergency department. Clinicians should consider the possibility of SLE when evaluating patients with pericardial effusion. Accurate and rapid diagnosis of this rare form of SLE could save lives. The aim of this case report is to raise awareness of these unusual appearances, to determine the best management approaches and to better understand the mechanisms underlying the different manifestations, hence the need for further research.

CONSENT

As per international standards or university standards, patient(s) written consent has been collected and preserved by the author(s).

ETHICAL APPROVAL

As per international standards or university standards written ethical approval has been collected and preserved by the author(s).

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Goswami RP, Sircar G, Ghosh A, Ghosh P. Cardiac tamponade in systemic lupus erythematosus. *QJM : An International Journal of Medicine*. 2017;111(2):83–87.
2. Alghareeb R, Hussain A, Maheshwari M v, Khalid N, Patel PD: Cardiovascular complications in systemic lupus erythematosus. *Cureus*. 2022;14:26671.
3. Emorinken A, Dic-Ijiewere MO, Izirein HO. Cardiac tamponade, an unusual first presentation of systemic lupus erythematosus: A case report in a rural tertiary hospital. *Cureus*; 2022.
4. Gomez Casanovas J, Bartl M, Rincon-Rueda L, Loftis CE, Dulgheru E. At the heart of the diagnosis : A case of systemic lupus erythematosus presenting as cardiac tamponade; 2023.
5. Amro AM, Deeb S, Rije R, Deeb N, Qunaibi YY, Amro B, Irzeqat K, Alhadad B, Emar A. Systemic lupus erythematosus presenting as cardiac tamponade and pleural effusion: A Case Report. *Cureus*; 2024.
6. Almousa S, Wannous H, Khedr K, Qasem H. Unusual case presentation of systemic lupus erythematosus in a young woman. *Rheumato*. 2022;2(4):93–97.
7. Emorinken A, Dic-Ijiewere MO, Izirein HO. Cardiac tamponade, an unusual first presentation of systemic lupus erythematosus: a case report in a rural tertiary hospital. *Cureus*. 2022 Aug 14;14(8).
8. Maharaj SS, Chang SM: Cardiac tamponade as the initial presentation of systemic lupus erythematosus: A case report and review of the literature. *Pediatr Rheumatol Online J*. 2015; 13:9.
9. Imazio M, De Ferrari GM. Cardiac tamponade: An educational review. *European Heart Journal: Acute Cardiovascular Care*. 2020; 204887262093934.
10. Curtis AC, Horne SF. Disseminated lupus erythematosus with pericardial effusion. *Ann Intern Med*. 1949 Jan;30(1):209-17.
11. Alerhand S, Adrian RJ, Long B, Avila J. Pericardial tamponade: A comprehensive emergency medicine and

- echocardiography review. The American Journal of Emergency Medicine; 2022.
12. Rosenbaum E, Krebs E, Cohen M, Tiliakos A, Derk CT: The spectrum of clinical manifestations, outcome and treatment of pericardial tamponade in patients with systemic lupus erythematosus: A retrospective study and literature review. *Lupus*. 2009;18:608-12.
 13. Cauduro SA, Moder KG, Tsang TS, Seward JB: Clinical and echocardiographic characteristics of hemodynamically significant pericardial effusions in patients with systemic lupus erythematosus. *Am J Cardiol*. 2003;92:1370-2.
 14. Carrion DM, Carrion AF: Cardiac tamponade as an initial manifestation of systemic lupus erythematosus. *BMJ Case Rep*. 2012;2012:bcr0320126126.
 15. Almousa S, Wannous H, Khedr K, Qasem H. Unusual case presentation of systemic lupus erythematosus in a young woman. *Rheumato*. 2022;2(4):93–97.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/115531>