



# Above Ground Biomass and Carbon Sequestration of Urban Green Spaces in Danang City, Vietnam

Ngoc-Son Tran <sup>a\*</sup> and Chi Cuong Doan <sup>a</sup>

<sup>a</sup> The University of Danang—University of Science and Education, 459 Ton Duc Thang St, Danang City, 550000, Vietnam.

## Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

## Article Information

DOI: 10.9734/AJOB/2024/v20i6409

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/115743>

Short Research Article

Received: 25/02/2024

Accepted: 29/04/2024

Published: 04/05/2024

## ABSTRACT

Green space in cities is an essential component of the urban environment. Different forms of urban green spaces, with varying populations, plant communities, and roles, offer varying urban ecological advantages. Using the i-Tree Eco model, this study examined the structural characteristics of urban green trees and calculated the biomass storage capacity and carbon sequestration in urban green spaces in Da Nang City. The results revealed a considerable diversity of species composition among 7,513 trees belonging to 42 species of 20 families studied in four types of green spaces, with *Terminalia catappa*, *Cyrtostachys renda*, and *Roystonea regia* dominating. The average Shannon biodiversity index was 2.9, with the institution having the highest at 2.6 and the park having the lowest at 2.2. The total leaf biomass per hectare was 588.0 kg/ha, and the carbon sequestration rate was 2,845.30 kg/ha. *Ficus subpisocarpa* was the species with the highest value for these two advantages, and the park was the green space with the highest benefit in terms of biomass and carbon sequestration.

\*Corresponding author: E-mail: [tinson@ued.udn.vn](mailto:tinson@ued.udn.vn), [tranngocsondhsp@gmail.com](mailto:tranngocsondhsp@gmail.com);

**Keywords:** *I-Tree Eco; green spaces; biomass; carbon sequestration; dominant species; Danang; Vietnam.*

## 1. INTRODUCTION

Climate change is defined as a shift in the climate through time caused by natural circumstances and human activity [1,2]. Global warming, rising sea levels, and increasing severe hydrometeorological occurrences are all symptoms of current climate change [3]. Climate change is regarded as the greatest threat to human health [4] and a major challenge confronting humanity, with human activity being the primary cause, owing to the production of one of the main greenhouse gases causing climate change, carbon dioxide, by the combustion of fossil fuels such as coal, oil, and natural gas. As a result, limiting carbon emissions is the most crucial answer to climate change.

Planting trees is one of the top priorities for addressing the challenge of reducing carbon emissions. By sequestering carbon and lowering vehicular pollution, urban trees play a vital ecological role in cities [5]. By fixing carbon during photosynthesis and storing carbon in the form of biomass, urban trees operate as a carbon dioxide reservoir [6]. Planting trees, preserving and growing urban green spaces provides chances for enhanced carbon removal from the atmosphere [7,8], as well as a variety of additional co-benefits for city inhabitants, such as improved health, reduced pollution, and increased infrastructure life [9]. Furthermore, trees in urban green spaces provide economic and social benefits such as energy savings, community inclusion, and outdoor leisure [10].

Numerous studies in Vietnam have been conducted to calculate biomass reserves and carbon buildup for trees. However, the majority of the work has been still done using traditional methods. The i-Tree Eco model was used in this study to evaluate the structure and quantify the value of biomass and carbon accumulation of trees [10] in four green spaces in Da Nang City, Vietnam, with the goal of assessing the current status of tree structure and species composition, as well as determining the biomass and carbon stocks of tree species in green spaces. The findings of the study are being utilized to track the city's carbon footprint.

## 2. MATERIALS AND METHODS

The research was conducted in four green spaces totaling 68.93 acres in Da Nang City, Vietnam. For the field data collection, we used the standard plot size for an Eco analysis is a 0.1 acre circular plot with radius of 11.4 m (USDA Forest Service, 2021). A total of 134 circular (Table 1) were created with public parks (36 plots), residential green spaces (28 plots), educational Institution green spaces (50 plots), and industrial green spaces (20 plots). Each plot has an area of 408.07 m<sup>2</sup> as recommended by Nowak et al [6].

During biomass estimate, a non-destructive sampling approach that does not affect plants was utilized, which requires the application of a generic or specialized correlation formula for a given type of vegetation [11]. Calculations for biomass and carbon sequestration based on i-Tree Eco model.

**Table 1. Characteristic sample plots of the four types of urban green spaces**

Four kind of urban greenspaces	Area of greenspaces(ha)	Number of sample plots	Locations
Public Park	9.13	36	29/3 Park
Residential spaces	7.71	28	Nguyen_Duc_Trung Residence, Phu_Loc Residence, Hoa_Minh Residence.
Educational institutions	20.83	50	Le_Van_Tam primary school, University of Sport Danang, The University of Danang (Danang University of Science and Education, Danang University of Science and Technology
Industrial zone	31.26	20	Hoa_Khanh industrial zone
<b>Total</b>	<b>68.93</b>	<b>134</b>	

The i-Tree Eco program (formerly known as UFORE, Urban Forest Effect Modeling) is intended to employ standardized field data from sample plots or complete inventory, as well as climatic data. The biomass and carbon sequestration of trees at four green spaces were calculated using local hourly visualization and air pollution for thorough characterization of urban green spaces structure and multi-ecosystem quantification (www.itreetools.org).

### 3. RESULTS AND DISCUSSION

#### 3.1 Plant Community Composition

The study surveyed 7,513 trees from 42 different families, with 24.4% coverage in four types of green spaces: parks, institution, industrial, and residential areas, with three species predominating: *Terminalia catappa*, *Cyrtostachys renda*, and *Roystonea regia*. Furthermore, several plant species, however few in number, are predominantly found in four green spaces, including *Samanea saman*, *Bauhinia variegata*,

*Plumeria rubra*, *Lagerstroemia speciosa*, *Alstonia scholaris*, and *Peltophorum pterocarpum* (Table 2).

The difference in the area of four types of green spaces has resulted in an uneven distribution of the plant system in terms of quantity and types, specifically, the industrial zone with an area of 31.26 hectares has 17 species with 2,399 trees (accounting for 31.93%), followed by the institution with an area of 20.83 ha has 22 species with 2,109 trees (28.07%), the park with an area of 9.13 ha has 31 species with 2,331 trees (31.03%), and the residential zone with an area of 7.71 ha has 23 species with 674 trees (accounting for 8.97%). When the Shannon diversity index is compared among the green spaces, it is found that the total score of four green spaces was at an average of 2.9, with the greatest being 2.6 for the institution and the lowest being 2.2 for the park. Industrial and residential zones have the same diversity index of 2.3 (Table 3).

**Table 2. Plant species composition at 4 green spaces**

Species	Families	Park	Residential area	Industry	Institution
<i>Acacia auriculiformis</i>	Fabaceae	✓	✓	✓	
<i>Alstonia scholaris</i>	Apocynaceae	✓	✓	✓	✓
<i>Areca catechu</i>	Arecaceae	✓			
<i>Artocarpus altilis</i>	Moraceae		✓		
<i>Artocarpus heterophyllus</i>	Moraceae		✓	✓	
<i>Barringtonia acutangula</i>	Lecythidaceae		✓		✓
<i>Bauhinia variegata</i>	Fabaceae	✓	✓	✓	✓
<i>Casuarina equisetifolia</i>	Casuarinaceae			✓	
<i>Cassia fistula</i>	Fabaceae				✓
<i>Ceiba pentandra</i>	Malvaceae		✓		
<i>Cinnamomum camphora</i>	Lauraceae	✓			
<i>Cocos nucifera</i>	Arecaceae	✓			✓
<i>Coccoloba uvifera</i>	Polygonaceae	✓			
<i>Cyrtostachys renda</i>	Arecaceae	✓	✓	✓	✓
<i>Chukrasia tabularis</i>	Meliaceae		✓		
<i>Dalbergia tonkinensis</i>	Fabaceae	✓	✓		
<i>Delonix regia</i>	Fabaceae	✓		✓	✓
<i>Dimocarpus longan</i>	Sapindaceae	✓	✓		
<i>Dracontomelon duperreanum</i>	Anacardiaceae	✓			
<i>Erythrina fusca</i>	Fabaceae		✓		✓
<i>Ficus benjamina</i>	Moraceae	✓			
<i>Ficus elastica</i>	Moraceae	✓			✓

Species	Families	Park	Residential area	Industry	Institution
<i>Ficus racemosa</i>	Moraceae	✓	✓		✓
<i>Ficus subpisocarpa</i>	<i>Ficus subpisocarpa</i>	✓			
<i>Ficus religiosa</i>	Moraceae		✓		
<i>Hopea odorata</i>	Dipterocarpaceae	✓	✓		✓
<i>Khaya senegalensis</i>	Meliaceae	✓			✓
<i>Lagerstroemia speciosa</i>	Lythraceae	✓	✓	✓	✓
<i>Mangifera indica</i>	Anacardiaceae		✓		✓
<i>Manilkara zapota</i>	Sapotaceae			✓	
<i>Mimusops elengi</i>	Sapotaceae			✓	
<i>Peltophorum pterocarpum</i>	Fabaceae	✓	✓	✓	✓
<i>Plumeria rubra</i>	Apocynaceae	✓	✓	✓	✓
<i>Pinus kesiya</i>	Pinaceae	✓			
<i>Roystonea regia</i>	Arecaceae	✓			✓
<i>Samanea saman</i>	Fabaceae	✓	✓	✓	✓
<i>Salix babylonica</i>	Salicaceae	✓			
<i>Senna siamea</i>	Fabaceae	✓			✓
<i>Syzygium samarangense</i>	Myrtaceae	✓	✓		
<i>Tamarindus indica</i>	Fabaceae	✓	✓		
<i>Terminalia catappa</i>	Combretaceae	✓	✓	✓	✓
<i>Terminalia mantaly</i>	Combretaceae	✓			



(*Terminalia catappa*)



(*Cyrtostachys renda*)



(*Roystonea regia*)

Fig. 1. Three common species at 4 green spaces

Table 3. Distribution of trees in four green spaces

Green spaces	Area (ha)	Number of plant species	Number of trees	Quantity ratio (%)	Shannon index
Industrial zone	31.26	17	2,399	31.93	2.3
Institution	20.83	22	2,109	28.07	2.6
Park	9.13	31	2,331	31.03	2.2
Residential zone	7.71	23	674	8.97	2.3
<b>Total</b>	<b>68.93</b>		<b>7,513</b>		<b>2.9</b>




Fig. 2. The DBH distribution of trees in four green spaces




Fig. 3. DBH distribution of trees in each green space

### 3.2 Tree Trunk Diameter Distribution

The distribution of trunk diameter (DBH) in the four green spaces was uneven, with DBH ranged from 7.6 to 15.2 cm accounting for 39.70%, followed by DBH from 0 - 7.6 cm (38.94%), DBH from 15.2 - 30.5 cm (20.73%), DBH from 30.5 - 45.7 cm (0.38%), and DBH greater than 45.5 cm (0.25%) (Fig 2).

When the distribution of DBH among trees in the four green spaces was compared, it was discovered that, with the exception of the park, the trees in the other green spaces were generally in the young and growing phases. Trees in the park have the most complete trunk diameter distribution structure, though there was still a difference in the distribution ratio between

DBHs, specifically: DBH from 0 - 7.6 cm accounted for 30.65%, 7.6 - 15.2 cm was 34.95%, 15.2 - 30.5 cm was 33.6%, 30.5 - 45.7 cm was 0.27%, and greater than 45.7 cm was 0.54%; followed by trees in the institute with DBH range from 0 - 7.6 cm accounting for 40.63%. Trees in residential and industrial areas exhibited a substantial difference in trunk diameter distribution between trees with BDH ranging from 0 to 7.2 cm and trees with DBH ranging from 15.2 to 30.5 cm (as shown in Fig 3).

*Samanea saman* has the most constant trunk diameter distribution among the 10 prominent species in the study region, whereas the majority of the other dominant species had a DBH ranged from 0 to 30.5 cm (Fig 4).




Fig. 4. Distribution of DBH of 10 dominant species in four green spaces

Table 4. Total biomass and carbon sequestration of trees at four green spaces

Type of urban green space	Tree density (trees/ha)	Leaf biomass density (kg/ha)	Carbon storage (kg/ha)	Net carbon sequestration density (kg/yr/ha)	Gross carbon sequestration density (kg/yr/ha)	CO <sub>2</sub> equivalent (kg/yr/ha)
Park	255.30	1,643.70	13,421.40	2,587.90	2,682.91	9,838.23
Residential	87.40	627.80	639.60	221.11	225.45	826.73
Industrial	76.70	432.30	744.30	233.83	240.95	883.57
Institution	101.30	344.10	2,179.00	637.10	657.03	2,409.32
Average	109.00	588.00	2,845.30	666.08	688.40	2,524.35

**Table 5. The top ten species have the highest levels of carbon storage and sequestration**

No.	Leaf biomass			Carbon storage				
	Species	Number of trees	Biomass (kg)	Average mass per tree (kg)	Species	Number of trees	Biomass (kg)	Average mass per tree (kg)
1	<i>Terminalia catappa</i>	1,333	11,458	8.60	<i>Roystonea regia</i>	1,316	82,320	62.55
2	<i>Roystonea regia</i>	1,316	4,505	3.42	<i>Samanea saman</i>	430	24,620	57.26
3	<i>Samanea saman</i>	430	3,952	9.19	<i>Ficus subpisocarpa</i>	13	16,490	1,268.46
4	<i>Dalbergia tonkinensis</i>	123	2,729	22.19	<i>Terminalia catappa</i>	1,333	11,560	8.67
5	<i>Lagerstroemia speciosa</i>	373	1,899	5.09	<i>Peltophorum pterocarpum</i>	406	11,070	27.27
6	<i>Delonix regia</i>	323	1,836	5.68	<i>Khaya senegalensis</i>	107	9,650	90.19
7	<i>Ficus subpisocarpa</i>	13	1,570	120.77	<i>Delonix regia</i>	323	6,710	20.77
8	<i>Peltophorum pterocarpum</i>	406	1,513	3.73	<i>Cocos nucifera</i>	141	3,860	27.38
9	<i>Hopea odorata</i>	431	1,440	3.34	<i>Ficus elastica</i>	16	3,620	226.25
10	<i>Dracontomelon duperreanum</i>	163	914	5.61	<i>Hopea odorata</i>	431	3,500	8.12

### 3.3 Carbon Storage and Sequestration

The overall tree density in the four green spaces was around 109.00 trees/ha, with the density in the park being the greatest at 255.30 trees/ha and the industrial zone being the lowest at 76.70 trees/ha. The four green spaces obtained a total biomass density of 588.00 kg/ha, with trees in the park reaching 1,643.70 kg/ha, followed by residential areas at 627.80 kg/ha, industrial zones at 432.30 kg/ha, and institutions at 344.10 kg/ha (Table 4).

The average carbon storage at four green spaces was 2,845.30 kg/ha, with the park having the most at 13,421.40 kg/ha and the residential having the lowest at 639.60 kg/ha. The densities of total net sequestration and carbon sequestration were 666.08 kg/year/ha and 688.40 kg/year/ha, respectively. The park had the highest net sequestration and carbon sequestration, at 2,587.90 kg/year/ha and 2,682.91 kg/year/ha, respectively; the institution had 637.10 kg/year/ha and 657.03 kg/year/ha, respectively; the industrial zone had 233.83 kg/ha/year and 240.95 kg/year/ha, respectively; and the residential area had 221.11 kg/year/ha and 225.45 kg/year/ha. The overall quantity of carbon dioxide equivalent was 2,524.35

kg/year/ha, with the park being the greenest space at 9,838.23 kg/year/ha and the residential area being the least at 826.73 kg/year/ha (Table 4).

The species with the greatest biomass value (11,458 kg) was *Terminalia catappa*, followed by *Roystonea regia* (4,505 kg) and *Samanea saman* (3,952 kg). However, when evaluating the average weight of a tree, *Ficus subpisocarpa* has the greatest biomass value (120.77 kg/tree). *Roystonea regia* was the species with the highest carbon storage value (82,320 kg), followed by *Samanea saman* (24,620 kg), *Ficus subpisocarpa* (16,490 kg). The highest average mass per tree belonged to *Ficus subpisocarpa* (1,268.46 kg/tree) (as shown in Table 5).

### 4. CONCLUSION

The study found the rich diversity in species composition among 7,513 trees belonging to 42 species from 20 families surveyed in four green spaces, with the major species being *Terminalia catappa*, *Cyrtostachys renda*, and *Roystonea regia*. The total Shannon biodiversity index was 2.9, with the institution area having the greatest at 2.6 and the park having the lowest at 2.2. The total biomass per hectare was 588.00 kg/ha, and

the total carbon sequestration was 2,845.30 kg/ha. *Ficus subpisocarpa* was the species with the highest value for biomass and carbon sequestration, and the park was the green space with the highest value for these two advantages.

## ACKNOWLEDGEMENTS

We are grateful to the U.S. Forest Service, GreenViet Biodiversity Conservation Centre and Frankfurt Zoological Society for providing us with technical training and funding.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Walther G, Post E, Convey P, Menzel A, Parmesan C, Beebee TJC., Fromentin J, IOH, Bairlein F. Ecological response to recent climate change. *Nature*. 2002;416:389–395.
2. Constance M. Climate change and forests of the future: managing in the face of uncertainty. *Ecological Applications*. 2007;17(8):2145–2151.
3. Thang NV. Climate change scenario. *Jurnal Penelitian Pendidikan Guru Sekolah Dasar*. 2016;128.
4. Patz JA, Campbell-Lendrum D, Holloway T, Foley JA. Impact of regional climate change on human health. *Nature*. 2005;438(7066):310–317. Available: <https://doi.org/10.1038/nature04188>
5. Kiran GS, Kinnary S. Carbon Sequestration By Urban Trees on Roadsides of. *International Journal of Engineering Science and Technology (IJEST)*. 2015;3(4):3066–3070.
6. Nowak DJ, Greenfield EJ, Hoehn RE, Lapoint E. Carbon storage and sequestration by trees in urban and community areas of the United States. *Environmental Pollution*. 2013;178:229–236. Available: <https://doi.org/10.1016/j.envpol.2013.03.019>
7. Niemelä J, Saarela SR, Söderman T, Kopperoinen L, Yli-Pelkonen V, Väre S, Kotze DJ. (). Using the ecosystem services approach for better planning and conservation of urban green spaces: A Finland case study. *Biodiversity and Conservation*. 2010;19(11):3225–3243. Available: <https://doi.org/10.1007/s10531-010-9888-8>
8. Livesley SJ, McPherson EG, Calfapietra C. The Urban Forest and Ecosystem Services: Impacts on Urban Water, Heat, and Pollution Cycles at the Tree, Street, and City Scale. *Journal of Environmental Quality*. 2016;45(1):119–124. Available: <https://doi.org/10.2134/jeq2015.11.0567>
9. Steenberg JWN, Ristow M, Duinker PN, Lapointe-Elmrabti L, MacDonald JD, Nowak DJ, Pasher J, Flemming C, Samson C. (). A national assessment of urban forest carbon storage and sequestration in Canada. *Carbon Balance and Management*. 2023;18(1):1–13. Available: <https://doi.org/10.1186/s13021-023-00230-4>
10. Song P, Kim G, Mayer A, He R, Tian G. Assessing the ecosystem services of various types of urban green spaces based on i-Tree Eco. *Sustainability (Switzerland)*. 2020;12(4). Available: <https://doi.org/10.3390/su12041630>
11. Rahmawaty, Sitorus NA, Rauf A. Distribution, above-ground biomass and carbon stock of the vegetation in Taman Beringin Urban Forest, Medan City, North Sumatra, Indonesia. *Malaysian Forester*. 2017;80(1), 73–84.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

### Peer-review history:

The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/115743>