



Effect of Foliar Application of Nano Urea, Boron and Zinc Sulphate on Growth, Yield and Quality of Guava (*Psidium guajava* L.) cv. Allahabad Surkha

Amit Kumar ^{a++*}, Annjoe V. Joseph ^{a#} and Vijay Bahadur ^{at}

^a Department of Horticulture, Naini Agricultural institute, SHUATS, Prayagraj, Uttar Pradesh, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/jabb/2024/v27i6887>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/116807>

Original Research Article

Received: 07/03/2024

Accepted: 10/05/2024

Published: 13/05/2024

ABSTRACT

The present experiment entitled "Effect of Foliar Application of Nano Urea, Boron and Zinc Sulphate on Growth, Yield and Quality of Guava (*Psidium guajava* L.) cv. Allahabad Surkha" was conducted at Department of Horticulture, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj during the session 2022 - 2024. The experiment was laid out in randomized block design with three replications, and the study consists of ten treatment combinations including control. The best treatment was T9 (Nano urea 2.0% + zinc sulphate 0.6% + boron 0.6%) & T8 (Nano urea 2.0% + zinc sulphate 0.4% + boron 0.4%) which

⁺⁺ M.Sc. Scholar;

[#] Assistant Professor;

[†] Head & Associate Professor;

^{*}Corresponding author: E-mail: amitjakhar183@gmail.com;

Cite as: Kumar, A., Joseph, A. V., & Bahadur, V. (2024). Effect of Foliar Application of Nano Urea, Boron and Zinc Sulphate on Growth, Yield and Quality of Guava (*Psidium guajava* L.) cv. Allahabad Surkha. *Journal of Advances in Biology & Biotechnology*, 27(6), 285–292. <https://doi.org/10.9734/jabb/2024/v27i6887>

shows the highest values in all the parameters viz., number of flower/tree (86.47), Days taken to fruit harvesting (49.19), fruit set (74.00%), Number fruit/tree (63.25), Fruit length (7.42 cm), fruit weight (178.73g), fruit diameter (7.12 cm), fruit yield/tree (11.29 kg). Increased flowering, fruit yield and quality might be due to the increased absorption of nutrients when given as foliar application.

Keywords: Guava; zinc sulphate; psidium; guajava; boron.

1. INTRODUCTION

Guava (*Psidium guajava* L.) known as the “Apple of the tropics” belongs to the family Myrtaceae and is a highly remunerative crop. It is an important fruit of tropical and subtropical areas of the world and was introduced in India in the 17th century by the Portuguese. The genus *Psidium* belongs to a myrtle family and emerged from Central America and Mexico. The genus *guajava* consists of different types like - yellow fruited cherry guava, strawberry and red apple guava. The commonly consumed guava is with yellow skin and either white or pink flesh. It is a fruit with high production in tropical and subtropical countries. The principal producers of guava in the world are India, Pakistan and Brazil.

Guava is mainly grown in the states of Uttar Pradesh, Madhya Pradesh, Maharashtra and Bihar. The excellent quality guava fruit in the world is produced in Allahabad district of Uttar Pradesh. Uttar Pradesh occupied first rank in the production of guava in India with production of 4,86,700 Metric tons. Approximately 30 – 50 percent of losses are seen in post-harvest handling due to the lack of marketing and storage facilities [1]. It contains remarkable mineral levels that include calcium, phosphorus, iron and vitamins such as niacin, pantothenic acid, thiamin, riboflavin and vitamin C. It has large amounts of antioxidant properties due to the existence of polyphenolic compounds and carotenoids in it.

Micronutrients like Zinc and Boron perform a specific role in the growth and development of plants, quality produce and uptake of major nutrients. Boron is much required for cell division and development in the growth regions of the plant near the tips of shoots and roots [2,3]. A shortage of Boron also causes cracking and distorted growth in fruit with the appearance of red spots on the newly emerged leaves. It can be managed by foliar application of Borax. Because boron is essential for the translocation of carbohydrates, auxin production, enhanced pollen viability, and fertilisation, borax produced a more beneficial reaction. Fruit set, pollen tube

expansion, and ovule development all depend on boron. Cell membranes contain boron, which is necessary for cell division. It aids in the plant's uptake of nitrogen and the translocation of sugar, and it regulates the ratio of potassium to calcium in the plant. In plants, boron improves the availability of nitrogen. It is thought that boron causes T9 growth hormone to become inactive by forming a complicated molecule.

Zinc is necessary for the production of tryptophan, which is a precursor to auxin and helps prevent fruit drop [4]. Applying zinc at a greater concentration raises the zinc content in the leaves, which in turn promotes the body's natural synthesis of auxin, which decreases fruit drop. It is crucial for the metabolism of starch and serves as a cofactor for several enzymes that impact protein formation, nucleic acid metabolism, and photosynthetic reactions. Low levels of auxin in tissue cause the severe stunting of leaves and shoots that characterises crops lacking in zinc. Young leaves are often the most damaged; they are typically tiny, thin, chlorotic, and frequently rosetted as a result of the shoot's inability to lengthen. Zinc is essential for growth and development, as well as for the growth, quality, and blooming of fruits. It also raises the chlorophyll content of leaves and plays a significant part in enzymatic activities.

Nitrogen is essential for oxidation-reduction processes and plays a role in the development of porphyrin ring structure, which forms chlorophyll molecules. An increase in the nitrogen content of leaves is linked to the development of green colour in those leaves. In stress situations as well as plant development, production, and quality, potassium is essential for a number of physiological and biochemical processes [5]. It is important for metabolic activities such as the synthesis of carbohydrates and the growth of meristematic tissue. In addition, potassium controls the transpiration process and water absorption while promoting the lignification of the cell walls. In a similar vein, zinc plays a crucial part in the activation of enzymes like catalase, peroxidase, and cytochrome oxidase as well as raising the chlorophyll content of leaves.

Applications of zinc, potassium, and urea had positive effects on fruit crop output and vegetative development. Therefore, for the guava crop to thrive and produce more effectively, the prudent administration of both macro and micronutrients is necessary.

2. MATERIALS AND METHODS

The area of Prayagraj district comes under subtropical belt in the South east of Uttar Pradesh, which experience extremely hot summer and fairly cold winter. The maximum temperature of the location reaches up to 46°C – 48°C and seldom falls as low as 4°C – 5°C. The relative humidity ranged between 20 to 94 percent. The average rainfall in this area is around 1100 mm annually. Prayagraj has a subtropical and semi- arid climate with rain mostly during July- September.

The experiment was laid out in Randomized Block Design Which consisting of ten treatments with T₀ – Control (N:P:K 0.8 % : 0.2% : 0.2%), T₁ – Nano urea 1.0% + zinc sulphate 0.2% + boron 0.2%, T₂ – Nano urea 1.0% + zinc sulphate 0.4% + boron 0.4%, T₃ - Nano urea 1.0% + zinc sulphate 0.6% + boron 0.6%, T₄ - Nano urea 1.5% + zinc sulphate 0.2% + boron 0.2%, T₅ - Nano urea 1.5% + zinc sulphate 0.4% + boron 0.4%, T₆ - Nano urea 1.5% + zinc sulphate 0.6% + boron 0.6%, T₇ - Nano urea 2.0% + zinc sulphate 0.2% + boron 0.2%, T₈ Nano urea 2.0% + zinc sulphate 0.4% + boron 0.4%, T₉ - Nano urea 2.0% + zinc sulphate 0.6% + boron 0.6%.The biometrical observation were recorded on five randomly selected plants of each replication to assess the morphological characters, *i.e.* Flowering Parameters, Physical Characteristics of fruit, Chemical Characteristics of fruit and Economics.

3. RESULTS AND DISCUSSION

3.1 Flowering Parameters

3.1.1 Number of days to anthesis

The data on number of days to anthesis of guava as influenced by Foliar Application of Nano-urea Boron and Zinc Sulphate are summarized in Table 1.

The data reveals that the number of days to anthesis of guava decreased significantly by the application of foliar application of nano urea, boron, zinc and sulphate under experimentation

over the control. The minimum number of days to anthesis (12) was recorded with treatments 9 (nano urea 2.0% + zinc sulphate 0.6% + boron 0.6%). While the maximum number of days to anthesis (27) was recorded under control. A decrease in number of days to anthesis might be due to the increased absorption of nutrients.

The minimum in number of days to anthesis of guava due to treatment might be due to fact that nano-urea regulate the growth by causing cell division and cell elongation in plant system. These results are conformed with Nikbakht et al. [6]. reported decreases in number of days to anthesis of guava with the use of nano urea Abobatta and Ahmed [7] reported increase in growth parameters with nano urea in guava. While the maximum value in treatment T₁ (control) may owes to its inhibitory effect because this treatment occupies only the recommended dose of fertilizers in guava, Increase in growth parameters (number of days to anthesis) use of nano urea may be due to its effect in cell division and cell enlargement Bhatti et al. [8].

3.1.2 Number of flowers/trees

The data on number of flowers/trees guava as influenced by Foliar Application of Nano Urea, Boron and Zinc Sulphate are summarized in Table 2.

The data reveals that the number of flowers/trees of guava increased significantly by the application of foliar application of nano-urea, boron, zinc and sulphate under experimentation over the control. The maximum number of flower/tree (86.47) was recorded with treatment 9 (nano urea 2.0% + zinc sulphate 0.6% + boron 0.6%). While the minimum Number of flower/tree (62.53) was recorded under control. The interaction effect of nano urea, boron, zinc sulphate significantly influenced the number of flowers/trees in guava.

The maximum increase in number of flowers per tree of guava due to treatment might be due to fact that zinc sulphate and boron regulate the growth by causing cell division and cell elongation in plant system. The beneficial effect of zinc sulphate and boric acid on fruit set and reducing fruit drop might be due to the higher availability of photosynthates. These chemicals are also associated with hormone metabolism which promotes synthesis of auxin, These results are in conformity with Zagade et al. [9] reported

Table 1. Effect of Foliar Application of Nano Urea, Boron and Zinc Sulphate on flowering parameters and fruit parameters of Guava (*Psidium guajava* L.) cv. Allahabad Surkha

Treatment Symbol	Treatment combinations	No. of days to anthesis	No. of flowers/tree	Number of Days taken to fruit harvest	Fruit set (%)	Number of fruit/tree
T ₀	Control (N:P:K 0.8% : 0.2% : 0.2%)	27	62.53	59.28	54.78	35.50
T ₁	Nano urea 1.0% + zinc sulphate 0.2% + boron 0.2%	16	69.88	60.20	58.58	39.18
T ₂	Nano urea 1.0% + zinc sulphate 0.4% + boron 0.4%	17	74.98	59.35	58.16	44.53
T ₃	Nano urea 1.0% + zinc sulphate 0.6% + boron 0.6%	19	76.74	58.17	58.56	66.40
T ₄	Nano urea 1.5% + zinc sulphate 0.2% + boron 0.2%	21	78.14	57.26	62.99	46.99
T ₅	Nano urea 1.5% + zinc sulphate 0.4% + boron 0.4%	24	81.73	56.11	67.29	51.99
T ₆	Nano urea 1.5% + zinc sulphate 0.6% + boron 0.6%	19	63.87	54.37	67.56	56.05
T ₇	Nano urea 2.0% + zinc sulphate 0.2% + boron 0.2%	14	82.87	54.24	68.01	60.48
T ₈	Nano urea 2.0% + zinc sulphate 0.4% + boron 0.4%	13	85.31	51.53	70.82	60.84
T ₉	Nano urea 2.0% + zinc sulphate 0.6% + boron 0.6%	12	86.47	49.19	74.00	63.25
	F-test	S	S	S	S	S
	SEm(±)	1.55	2.19	1.64	2.23	1.55
	CD (p=0.05)	0.58	6.52	4.89	6.63	4.60

Table 2. Effect of Foliar Application of Nano Urea, Boron and Zinc Sulphate on physical characteristics of fruit of guava (*Psidium guajava* L.) cv. Allahabad Surkha

Treatment Symbol	Treatment combinations	Fruit length (cm)	Diameter of fruit (cm)	Fruit weight (g)	Fruit yield /tree(kg)
T ₀	Control (N:P:K 0.8% : 0.2% : 0.2%)	5.59	5.99	142.36	5.03
T ₁	Nano urea 1.0% + zinc sulphate 0.2% + boron 0.2%	5.83	5.99	147.23	5.77
T ₂	Nano urea 1.0% + zinc sulphate 0.4% + boron 0.4%	6.98	6.19	153.70	6.84
T ₃	Nano urea 1.0% + zinc sulphate 0.6% + boron 0.6%	5.92	6.38	158.60	7.45
T ₄	Nano urea 1.5% + zinc sulphate 0.2% + boron 0.2%	6.55	6.47	163.80	8.51
T ₅	Nano urea 1.5% + zinc sulphate 0.4% + boron 0.4%	6.78	6.56	169.40	9.47
T ₆	Nano urea 1.5% + zinc sulphate 0.6% + boron 0.6%	6.98	6.73	173.56	10.49
T ₇	Nano urea 2.0% + zinc sulphate 0.2% + boron 0.2%	7.28	6.82	175.80	10.69
T ₈	Nano urea 2.0% + zinc sulphate 0.4% + boron 0.4%	7.36	6.94	175.93	11.11
T ₉	Nano urea 2.0% + zinc sulphate 0.6% + boron 0.6%	7.42	7.12	178.73	11.91
	F-test	S	S	S	S
	SEm(±)	0.20	0.19	4.81	0.30
	CD (p=0.05)	0.58	0.57	14.28	0.88

increase in number of flowers per tree of guava with the use of zinc Sulphate and boric acid. Abobatta and Ahmed [7] reported increase in growth parameters with nano urea in guava. While the minimum value in treatment T1 (control) may owe to its inhibitory effect because this treatment occupies only recommended dose of fertilizers in guava. Increase in growth parameters (number of flowers per trees) use of nano urea may be due to its effect in cell division and cell enlargement Sharif et al. [10].

3.1.3 Days number of taken to fruit harvesting

The data reveals that the days taken to fruit harvesting of guava increased significantly by the application of foliar application of nano urea, boron, zinc and sulphate under experiment over the control. The minimum number days taken to fruit harvesting (49.19) was recorded with treatments 9 (Nano urea 2.0% + zinc sulphate 0.6% + boron 0.6%) while the maximum number days taken to fruit harvesting (60.20) was recorded under control. Further, the interaction effect of nano urea, boron, zinc sulphate significantly influenced the number of days taken to fruit harvesting.

The minimum number days taken to fruit harvesting of guava due to treatment might be due to fact that zinc sulphate and boron regulate the growth by causing cell division and cell elongation in plant system. The beneficial effect of zinc sulphate and boric acid on fruit set and reducing fruit drop might be due to the higher availability of photosynthates. These chemicals are also associated with hormone metabolism which promotes synthesis of auxin, These results are in conformity with Raj Kumar et al. [11] reported decreases in number of days taken to fruit harvesting of guava with the use of zinc Sulphate and boric acid. Singh et al. [12] reported increase in growth parameters with nano urea in guava.

While the maximum value in treatment T1 (control) may owes to its inhibitory effect because this treatment occupy only recommended dose of fertilizers in guava. Increase in growth parameters (number of days taken to fruit harvesting) use of Nano urea may be due to its effect in cell division and cell enlargement Yadav et al. [13].

3.2 Fruit Set Percentage

The data reveals that the Fruit set percent of guava increased significantly by the application

of foliar application of nano urea, boron, zinc and sulphate under experimentation over the control. The maximum fruit set (74%) was recorded in treatments 9 (Nano urea 2.0% + zinc sulphate 0.6% + boron 0.6%). while the minimum fruit set (54.78%) was recorded under control. The interaction effect of nano urea, boron, zinc sulphate significantly influenced the fruit set per cent in guava.

Increase in fruit set percent with the use of zinc might be due to fact that benzyle adenine causes the production of large number of fruits with rapid elongation of peduncle, leading to full development of flower buds having all reproductive parts functional which increases the fruit set and number of berries per plant. It could also be due to the fact that zinc application accelerates the development of differentiated inflorescence.

Singh and Tripathi [14] observed increase in yield parameters following use of treatment Nano urea and zinc in guava. Similar results have been reported by Saima et al. [15].

3.2.1 Number of fruit/tree

The data reveals that the Number of Fruit/tree of guava increased significantly by the application of foliar application of nano urea, boron, zinc and sulphate under experimentation over the control. The maximum Number of fruits/tree (66.40) was recorded with treatments 9 (nano urea 2.0% + zinc sulphate 0.6% + boron 0.6%) while the minimum Number of fruits/tree (35.50) was recorded under control. The interaction effect of nano urea, boron, zinc sulphate significantly influenced the fruit set per cent in guava. Increase number of fruit plant was might be due to the increased duration of flowering. This might be due to the fact their more food reserves were available for less number of flower buds.

Increase in number of fruits per tree with the use of zinc might be due to fact that benzyl adenine causes the production of large number of fruits with rapid elongation of peduncle, leading to full development of flower buds having all reproductive parts functional which increases the fruit set and number of berries per plant. It could also be due to the fact that zinc application accelerates the development of differentiated inflorescence.

Bhatti et al. [8] observed increase in yield parameters following use of treatment Nano urea

and zinc in guava. Similar results have been reported by Saima et al. [15].

3.3 Physical Characteristics of Fruit

3.3.1 Fruit length (cm)

The data reveals that the fruit Length of guava increased significantly by the application of foliar application of nano urea, boron, zinc and sulphate under experimentation over the control. The maximum fruit length (7.42 cm) was recorded with treatments 9 (nano urea 2.0%+ zinc sulphate 0.6% + boron 0.6%) while the minimum fruit Length (5.59 cm) was recorded under control. The interaction effect of nano urea, boron, zinc sulphate significantly influenced the fruit length in guava.

The maximum Increase in fruit length (mm) following use of zinc might be due to its effect in cell division and cell elongation. Zinc is also reported to promote growth by increasing plasticity of cell wall followed by hydrolysis of starch into sugar which reduces cell wall potential, resulting in the entry of water into the cell and causing its elongation (Richard, 2006).

Singh et al. [12] reported maximum length and minimum days to first harvesting in guava and 'Brighton' with zinc 2%. These results are in close similarity with the findings of Lal and Das [16]; Yadav et al. [13].

3.3.2 Diameter of fruit

The data reveals that the diameter of fruit of guava increased significantly by the application of foliar application of nano urea, boron, zinc and sulphate under experimentation over the control. The maximum diameter of fruit (7.12 cm) was recorded with treatments 9 (Nano urea 2.0% + zinc sulphate 0.6% + boron 0.6%) while the minimum diameter of fruit (5.99 cm) was recorded under control. The interaction effect of Nano Urea, boron, zinc sulphate significantly influenced the fruit set per cent in guava.

The maximum increase in fruit diameter following use of zinc sulphate and boric acid might be due to its effect in cell division and cell elongation. The beneficial effect of zinc sulphate and boric acid on fruit set and reducing fruit drop might be due to the higher availability of photosynthates. These chemicals are also associated with hormone metabolism which promotes synthesis of auxin, These results are in conformity with

Rajkumar et al. [11] reported increase in fruit diameter of guava with the use of zinc Sulphate and boric acid. Abobatta and Ahmed [7] reported increase in growth parameters with Nano urea in guava.

While the minimum value in treatment T1 (control) may owes to its inhibitory effect because this treatment occupy only recommended dose of fertilizers in guava. Increase in growth parameters (fruit diameter) use of nano urea may be due to its effect in cell division and cell enlargement Sharif et al. [10].

3.3.3 Fruit weight (g)

The data reveals that the fruit weight of guava increased significantly by the application of foliar application of nano urea, boron, zinc and sulphate under experimentation over the control. The maximum fruit weight (178.73g) was recorded with treatment 9 (Nano urea 2.0% + zinc sulphate 0.6% + boron 0.6%) while the minimum fruit weight (142.37g) was recorded under control. The interaction effect of nano urea, boron, zinc sulphate significantly influenced the fruit weight in guava.

The maximum increase in fruit weight following the use of zinc might be due to its effect in cell division and cell elongation. Zinc is also reported to promote growth by increasing plasticity of cell wall followed by hydrolysis of starch into sugar which reduces cell wall potential, resulting in the entry of water into the cell and causing its elongation Shreekant *et al.* (2017).

Zagade et al. [9] reported maximum fruit weight guava with nano urea and zinc 2%. These results are in close similarity with the findings of Singh et al. (2013).

3.3.4 Fruit yield/tree

The data reveals that the fruit yield/tree of guava increased significantly by the application of foliar application of nano urea, boron, zinc and sulphate under experimentation over the control. The maximum fruit yield/tree (11.29kg) was recorded with treatments 9 (Nano urea 2.0% + zinc sulphate 0.6% + boron 0.6%) while the minimum fruit yield/tree (5.03kg) was recorded under control. The interaction effect of nano urea, boron, zinc sulphate significantly influenced the fruit yield/tree in guava.

The maximum increase in fruit yield/tree following use of zinc Sulphate and boric acid

might be due to its effect in cell division and cell elongation. The beneficial effect of zinc sulphate and boric acid on fruit set and reducing fruit drop might be due to the higher availability of photosynthates. These chemicals are also associated with hormone metabolism which promotes synthesis of auxin, These results are in conformity with Chander et al. [17] reported increase in fruit yield per tree of guava with the use of zinc Sulphate and boric acid. Abobatta and Ahmed [7].

While the minimum value in treatment T1 (control) may owe to its inhibitory effect because this treatment occupies only the recommended dose of fertilizers in guava. Increase in growth parameters (fruit yield per tree) use of zinc Sulphate and boric acid may be due to its effect in on cell division and cell enlargement Sharif et al. [10].

4. CONCLUSION

Based on the above findings it can be concluded that guava with the application of nano urea along with the application of micronutrient ZnSO₄ and boron recorded the highest number of days to anthesis, number of flower tree⁻¹, days taken to fruit harvesting, fruit set percent, number of fruits tree⁻¹, fruit length, diameter of fruit, fruit weight, fruit yield tree⁻¹.

ACKNOWLEDGEMENT

The authors are thankful to Department of Horticulture, Naini Agricultural Institute, Prayagraj, Sam Higginbottom University of Agriculture Technology And sciences, (U.P) India for providing necessary facilities to undertaken the studies.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Pooja J, Ghanshyam DS, Vijay K. Preparation and evaluation of Guava jelly (*Psidium guajava*). Journal of Pharmacognosy and Phytochemistry. 2020;9(6):2061-2063.
2. Kumar J, Kumar R, Rai R, Mishra DS. Response of pant prabhat guava trees to foliar sprays of zinc, boron, calcium and potassium at different plant growth stages.

- International Quarterly Journal of Life Science. 2015;10(2):495-498.
3. Tripathi P, Dubey NK, Shukla AK, Use of some essential oils as post-harvest botanical fungicides in the management of grey mould of grapes caused by *Botrytis cinerea*. World Journal of Microbiology and Biotechnology. 2008;24:39-46.
4. Hassan HSA, Sarrwy SMA, Mostafa EAM. Effect of foliar spraying with liquid organic fertilizer, some micronutrients and gibberellins on leaf minerals content, fruit set, yield, and fruit quality of "Hollywood" plum trees. Agriculture and Biology Journal of North America. 2010;1:638-643.
5. Marschner H. Mineral nutrition in higher plants. Academic Press, New York. 1995;299- 312.
6. Nikbakht M, Solouki M, Aran M. Effects of foliar application of nanonitrogen and urea fertilizers on quantity and quality properties of bitter apple (*Citrullus colocynthis* L.). Journal of Crops Improvement. 2021;23(1):155-168.
7. Abobatta WF, Ahmed FK. Effect of urea and nano-nitrogen spray treatments on some citrus rootstock seedlings. Horticulture Research Journal. 2023;1(1):68-84.
8. Bhatti D, Varu DK, Dudhat M. Effect of different doses of urea and nano-urea on yield and quality of guava (*Psidium guajava* L.) cv. Lucknow-49. The Pharma Innovation Journal. 2023;12(6):4151-4156.
9. Zagade PM, Munde GR, Shirsath AH. Effect of foliar application of micronutrients on yield and quality of Guava (*Psidium guajava* L.) Cv. Sardar. Journal of Pharmacy and Biological Sciences. 2017;12(5):56-58.
10. Sharif BS, Neasham J, Hinton OR, Adams AE. A computationally efficient doppler compensation system for underwater acoustic communications. IEEE Journal of Oceanic Engineering. 2000;25(1):52-61.
11. Rajkumar JP, Tiwari Shant Lal, Mohit Kumar, Anshuman Singh, Ashwani Kumar. Effect of boron and zinc application on nutrient uptake in Guava (*Psidium guajava* L.) cv. Pant Prabhat Leaves Int. J. Curr. Microbiol. App. Sci. 2014;6(6):1991-2002.
12. Singh YK, Singh SS, Singh R, Prasad VM, Yadav A. Assess the effect of different levels of micronutrient on quality attributes of Guava (*Psidium guajava* L.)" Cv. Allahabad Safeda. Journal of

- Pharmacognosy and Phytochemistry. 2017;6(6):1340-1345.
13. Yadav A, Verma RS, Ram RB, Kumar V, Yadav R. Effect of foliar application of micronutrients on physical parameters of winter season Guava (*Psidium guajava* L.) cv. Lalit. Plant Archives. 2017;17(2):1457-1459.
 14. Saxena A, Tripathi RM, Singh RP. Biological synthesis of silver nanoparticles by using onion (*Allium cepa*) extract and their antibacterial activity. Dig J Nanomater Bios. 2010;5(2):427-432.
 15. Saima Z, Sharma A, Umar I, Wali VK, Effect of plant bio-regulators on vegetative growth, yield and quality of strawberry cv. Chandler. African journal of agricultural research. 2014;9(22):1694-1699.
 16. Lal N, Das RP. Effect of plant growth regulators on yield and quality of guava (*Psidium guajava* L.) cv. Allahabad Safeda. Int.J.Curr. Microbiol. App. Sci. 2017;6(5):857- 863.
 17. Chander S, Jain MC, Pareek PK, Bola PK, Meena RR, Sharma YK, Renuka. Effect of foliar feeding of borax, zinc sulphate and urea on fruiting and yield of guava (*Psidium guajava* L.) cvs. Lalit and Shweta under high density planting system. Chem Sci Rev Lett. 2017;6(22):874-883.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/116807>