



# Response of Starter Nitrogen and Phosphorus Fertilizer Rates to Nodulation, Nitrogen Fixation, and Net Nitrogen Balance on Groundnut (*Arachis hypogaea* L.) in the Semi-Arid Tigray Region of Ethiopia

Kinfe Tekulu <sup>a\*</sup>

<sup>a</sup> Department of Soil Fertility Research, Tigray Agricultural Research Institute, Shire Soil Research Centre, P.O. Box 40, Shire, Ethiopia.

## Author's contribution

Author KT initiated the research, wrote the thesis research proposal, conducted the research, did data entry and analysis, and wrote the manuscript.

## Article Information

DOI: <https://doi.org/10.9734/asrj/2024/v8i4158>

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/120966>

Original Research Article

Received: 13/06/2024  
Accepted: 17/08/2024  
Published: 23/08/2024

## ABSTRACT

For small-scale farmers in developing countries, biological nitrogen fixation through legumes is an affordable method of reducing reliance on nitrogen fertilizer. The study aimed to look at how groundnut nodulation, biological nitrogen fixation, atmospheric nitrogen, and the net nitrogen balance were affected by starter nitrogen and phosphorus fertilizer dosages. The three times

\*Corresponding author: E-mail: [tekulukinfe@gmail.com](mailto:tekulukinfe@gmail.com);

**Cite as:** Tekulu, Kinfe. 2024. "Response of Starter Nitrogen and Phosphorus Fertilizer Rates to Nodulation, Nitrogen Fixation, and Net Nitrogen Balance on Groundnut (*Arachis Hypogaea* L.) in the Semi-Arid Tigray Region of Ethiopia". *Asian Soil Research Journal* 8 (4):8-20. <https://doi.org/10.9734/asrj/2024/v8i4158>.

replicated treatments of three nitrogen levels (0, 15 and 30 kg N ha<sup>-1</sup>) and four phosphorus levels (0, 23, 46 and 69 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup>) were arranged in a factorial combination using a randomized complete block design (RCBD). The results of the analysis demonstrated that, in comparison to plots without nitrogen fertilizer, the number of nodules per plant was significantly increased by the single application of nitrogen alone at a dose of 15 kg ha<sup>-1</sup>. Compared to 69 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup>, biological nitrogen fixation rose by 88.41% at 46 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup>; nonetheless, it was significantly similar to zero and 23 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup>. The nitrogen derived from the atmosphere (50.9%) was significantly higher in the control treatments. The interactive application of 15 kg of nitrogen ha<sup>-1</sup> and zero phosphorus resulted in a significantly greater net nitrogen balance (-4 kg ha<sup>-1</sup>), whereas the co-application of 30 kg ha<sup>-1</sup> of nitrogen and zero phosphorus resulted in the lowest measurement of net nitrogen balance. Therefore, large nitrogen fertilizer application rates are not necessary to promote BNF in groundnuts.

*Keywords: Biological nitrogen fixation; groundnut; net nitrogen balance; nitrogen; phosphorus fertilizers; nodulation.*

## 1. INTRODUCTION

Unbroken cereal cultivation has led to the loss of soil fertility and, thus, low yields [1,2]. Crop rotation is more productive than continuous monoculture [3,4]. In crop rotation, legumes play an important role in cereals due to their contribution of residual nitrogen (N) from biological nitrogen fixation (BNF). The biosphere's primary mechanism for sustaining life on earth is the BNF, ranked second in significance only to photosynthesis [5]. This is due to the role played by legumes in fixing nitrogen, restoring soil fertility through rhizobia-legume symbiosis, and breaking the cycles of pests and diseases that attack crops [6]. Nitrogen derived from the atmosphere can reduce the use of synthetic N fertilizer in cereal crops in rotation after legumes [5,6]. Legumes are therefore becoming a viable alternative to N fertilizer for resource-poor farmers [7] by constantly adding N to the soil to maintain existing N reserves in the soils [8].

Groundnut is a legume crop that produces its N via BNF and transfers large amounts of N to grains to produce protein and also provides the soil with residual N, which is essential for rotational cereal crops [9]. When residues are present and incorporated into the soil, a higher content of mineralized N is found in the soil, which subsequently ensures a promising yield for the successor cereal crops [10]. Biological nitrogen fixation is a renewable N cycle and a source of N for agriculture, and it allows host plants to grow in environments with low soil N, reducing N losses through denitrification, volatilization and leaching, thereby promoting sustainable agriculture [11]. The discovery by Bado et al. [12], confirmed that groundnut can fix

N in an amount equivalent to 35 kg ha<sup>-1</sup> and that subsequent cereal crops achieve higher yields when the stover residues of groundnut are returned to the soil [13]. Further findings from Yakubu et al. [14] reported that groundnut can produce 27.19 kg N ha<sup>-1</sup>. Bado et al. [15] observed that groundnut sequester has an average of 8 to 23 kg N ha<sup>-1</sup> and that the percentage of N derived from the atmosphere ranges from 27 to 34%. Although legumes produce N, they can suffer from N deficiency under field conditions, especially when soil N levels are very low, until nodules become functional [16,17]. In such situations, it may be necessary to add a starter N fertilizer at a rate of 20 kg N ha<sup>-1</sup> [18]. Many researchers have also reported that P fertilizer is essential for healthy root growth and profuse nodulation, which in turn may influence the BNF potential of legumes [19,20]. Nodulation of legumes requires energy, so P nutrition is desirable to provide energy for the BNF process and plant development, including energy transfer from legumes during BNF [21], hence the application of 40 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> in groundnut plots increased nodule number by 160%, N content by 147% and the amount of fixed N by 169% over the unfertilized plots. An increase in soil available N was observed in P treated groundnut plots, suggesting that P increased the BNF of groundnut and consequently led to the formation of residual N in the soil [14].

Farmers in the research area typically grow groundnut in rotation with finger millet, sorghum, and *Eragrostis tef* because legumes increase the grain yields of subsequent crops [3]. In contrast to sorghum and wheat monoculture, Bado et al. [12] and Angus et al. [22] observed increased yields of both sorghum and wheat following

legume rotation. It is estimated that American farmers might save \$200 million to \$300 million by employing alfalfa legumes in their crop production [23]. Legumes can generally be used to lessen impoverished farmers' reliance on artificial fertilizers [24]. The objectives of this study were to ensure proper management, offer some additional insights, and fully realize the advantages of legumes as a source of nitrogen nutrients for agricultural systems. Due to a lack of research on these subjects, this study set out to determine the impacts of starter N and P fertilizer treatments on BNF, net nitrogen balance, and groundnut nodulation.

## 2. MATERIALS AND METHODS

### 2.1 Site Description

The experiment was conducted during the 2017 growing season in Medhin village, Mereb Leke district, Tigray region of Ethiopia. Geographically, the study location was between 14° 18' 48" and 14° 25' 26" N and 38° 42' 15" and 38° 48' 30" and with an altitude of 1390 m above sea level, and it is located 42 km from the historical administrative zone of the city of Aksum. Mixed farming, which usually includes crop and animal production, is commonly carried out on the same land unit. The area is generally known for the predominant crops such as sorghum, millet, groundnut and teff, which are grown in rotation. In addition, fruits and vegetables are regularly grown in irrigated agriculture. The study area is characterized by a hot and semi-arid agro-climatic zone and has a well-drained sandy loam soil texture with typical Luvisols soil type. The rainfall regime is unimodal, with the main rainy season concentrated from June to September. During the research year, the average annual precipitation and average monthly temperature were 636 mm and 25.3°C, respectively [25]. The optimal rainfall depth and temperature for groundnut production are between 200 and 1000 mm and 20 and 30 °C, respectively [18]. Therefore, the current weather data in the study area was in a suitable range for groundnut cultivation.

### 2.2 Experimental Design and Treatments Set up

The experiment was conducted in the field of a selected farmer who had not grown groundnut in the previous growing season to determine the existing effects of the treatments on nodulation, BNF and net soil nitrogen balance of groundnut.

In this experiment, groundnut and sorghum crops were planted side by side in the same soil. Sorghum and groundnut crops were planted as non-nitrogen fixers and nitrogen fixers, respectively, in order to estimate BNF. The same treatments were applied to sorghum (a non-nitrogen fixer) and groundnut (a nitrogen fixer). The field trials included a total of 12 treatments, with three N levels at doses of 0, 15 and 30 kg ha<sup>-1</sup> and four P<sub>2</sub>O<sub>5</sub> levels at doses of 0, 23, 46 and 69 kg ha<sup>-1</sup>. These treatments were arranged in 3 × 4 factorial combinations and in RCBD with three replicates. The treatment with zero N and P fertilizer content was designated as the control treatment. All treatments were uniformly enriched with balanced fertilization containing potassium (K) at 30 kg K<sub>2</sub>O ha<sup>-1</sup>, sulfur (S) at 30 kg ha<sup>-1</sup>, zinc (Zn) at 2 kg ha<sup>-1</sup> and boron (B) at 0.5 kg ha<sup>-1</sup> to avoid BNF loss due to a limited supply of one of those nutrients from the soil. According to Cisse [26], a balanced fertilization schedule is important to avoid yield-limiting factors and nutrient balance should be considered beyond N, P and K and include secondary elements and micronutrients when the soil is facing nutrient deficiencies. The fertilizers N, P, K and S were mainly applied during the sowing period, while the microelements (Zn and B) were applied during the flowering period of groundnut. The source of these nutrients was urea for N, Trisuperphosphate (TSP) for P<sub>2</sub>O<sub>5</sub>, KCl for K<sub>2</sub>O, CaSO<sub>4</sub> for S, ZnSO<sub>4</sub> for Zn, and borax (Na<sub>2</sub>B<sub>4</sub>O<sub>7</sub>·10H<sub>2</sub>O) for B. A landrace Sedi groundnut variety was planted at a seed rate of 80 kg ha<sup>-1</sup> as a biological nitrogen-fixing plant. The *melkam* sorghum variety was planted as a non-nitrogen fixing plant at a seed rate of 15 kg ha<sup>-1</sup>. The plot size for growing groundnut (biological nitrogen fixer) was 3.6 m × 4 m (14.4 m<sup>2</sup>) with a net harvestable area of 9.6 m<sup>2</sup>, and the distance between blocks, plots, rows and plants for groundnut cultivation was set at 1.5, 1, 0.6 and 0.1 m, respectively. One plot had 6 rows of groundnut and a total stand of 240 groundnut plants. For sorghum cultivation, the distances between blocks, plots, rows and plants were 1.5 m, 1 m, 0.75 m and 0.2 m, respectively. The sorghum plot consisted of 5 rows and a total population of 100 plants.

### 2.3 Soil Sampling Procedures and Analysis

Before planting, a composite sample of the disturbed soil was collected with an auger sampler at a depth of 0–20 cm using the zigzag method according to the experimental setup.

This pre-planting composite sample was used to characterize the history of the site. Before planting, an additional sample of undisturbed soil was taken using a core sampler to determine the dry bulk density of the soil. After harvesting both crops, 36 soil samples were collected at similar depths for the sorghum and groundnut crops in each plot of the two crops. Post-harvest samples were deliberately used for the BNF determination. These disturbed composite soil samples were air-dried, ground and passed through a 2 mm sieve to analyze some chemical and physical properties of the soil. Soil samples collected after harvest were analyzed primarily to estimate residual soil N contents. The collected disturbed samples were analyzed for soil texture, pH, electrical conductivity (EC), organic carbon (OC), total N, available P, cation exchange capacity (CEC), and exchangeable cations (K, Na, Ca and Mg). Soil texture was determined

using the Bouyoucos hydrometric method [27]. The soil pH was measured in the supernatant suspension at a soil/water ratio of 1:2.5 using a pH meter [28]. The electrical conductivity of a soil solution in a saturated extract with a soil/water ratio of 1:2.5 was measured with an EC meter [29]. The percentage of soil organic carbon (SOC) was determined using the wet oxidation method proposed by Walkley and Black [30]. Total nitrogen was determined using the Kjeldahl method described by Bremner and Mulvaney [31]. Available P was determined using the Olsen method [32]. The cation exchange capacity and exchangeable cations of the soil were extracted by leaching an ammonium acetate solution, and the exchange cations were measured by atomic absorption spectrophotometry for Ca and Mg and a flame photometer for Na and K [33]. The soil parameters analyzed for the above parameters are listed in Table 2.

**Table 1. Total rainfall and temperature during experimental period**

Month	Rainfall	Temperature (°C)		
	Total rainfall (mm)	Maximum	Minimum	Mean
January	0	34.62	15.7	25.16
February	0	35.02	12.6	23.81
March	3.1	37.8	15.18	26.49
April	27	37.92	15.6	26.76
May	53.1	37.2	18	27.6
June	78.2	36.62	17.24	26.93
July	300.1	33.58	15.1	24.34
August	78.7	32.68	14.28	23.48
September	79	34.88	16	25.44
October	16.3	35.86	14.7	25.28
November	0	35.24	13.22	24.23
December	0	35.2	12.1	23.65
Total	635.5			

**Table 2. Physiochemical properties of experimental soil before sowing (NA-not applicable)**

Parameters	Values	Rating	References
%Clay	8	NA	
%Silt	39	NA	
%Sand	53	NA	
Textural Class	Sandy loam	NA	
Bulk density (g cm <sup>-3</sup> )	1.46	Moderate	[34]
pH	5.5	Moderately acid	[35]
EC(dS m <sup>-1</sup> )	2.94	Slightly saline	[34]
CEC (cmol(+) kg <sup>-1</sup> )	13.2	Moderate	[34]
OC (%)	0.61	Low	[34]
Total N (%)	0.053	Low	[34]
Available P (ppm)	6.92	Low	[34]
Exc.K (cmol(+) kg <sup>-1</sup> )	0.21	Low	[34]
Exc.Na (cmol(+) kg <sup>-1</sup> )	0.16	Low	[34]
Exc.Ca (cmol(+) kg <sup>-1</sup> )	5.8	Moderate	[34]
Exc.Mg (cmol(+) kg <sup>-1</sup> )	6.8	High	[34]

## 2.4 Collected Plant Data, Sampling Procedures and Analysis

Nodulation was assessed at the mid flowering of the groundnut. With destructive sampling, five randomly selected plants were taken from each plot, and a nodule count per plant was performed to determine effective nodules. The whole plant was carefully uprooted using a fork so as to obtain intact roots and nodules. From the uprooted plants, the number of nodules per plant was recorded by counting the number of nodules from five plants and averaging the number of nodule per plant. Nodules with a pink to dark-red color were considered effective nodules. Nodules with a green color were identified as non-effective nodules. At the time of harvesting of groundnut and sorghum crops, five plant samples were collected from 36 plots for each groundnut crop (haulms and grains) and sorghum crop (stovers and grains) to determine the N accumulated in the plants. The groundnut plants were dug up, depodded and the pods taken away. The five sorghum plant samples taken from the inner rows, were cut near the ground. Nitrogen determination in sorghum plant tissues was used as a non-nitrogen fixing plant to evaluate groundnut BNF. All nitrogen fixing and non-nitrogen fixing plant samples were thoroughly washed with distilled water to remove adhered contaminants and dust particles, then placed in a paper envelope and oven-dried at 60 °C for 68 h. After drying in the oven, haulms and grains of groundnut and stover and grains of sorghum were crushed and passed through a 0.5 mm sieve. The determination of the total N content of the plant samples was carried out by the Kjeldahl method using the treatment of plant materials and oxidation with concentrated sulfuric acid and distillation of an aliquot from the digest with NaOH, collection of the distillate in boric acid and titration with 0.01 N HCl to the endpoint of the mixed indicator [36].

## 2.5. Biological Nitrogen Fixation

The amount of biological nitrogen fixation by groundnut was calculated from the total nitrogen difference (TND). The TND method considers the cultivation of nitrogen-fixing legumes with adjacent non-nitrogen-fixing crops grown on the same land with the same soil fertility status and assumes that the difference in nitrogen in plant tissue at harvest between crops was the quantity of fixed N<sub>2</sub>. In the current experimental case, groundnut was the nitrogen-fixing crop, while

sorghum was the non-nitrogen-fixing crop. This method assumes that legumes and non-legumes absorb the same amount of N from the soil. The reason for choosing sorghum is due to its agroecological needs and root morphology, which is similar to that of groundnut. For each plot, the TND method was used based on the requirement for the same treatments. The BNF of groundnut was calculated according to the method described by Peoples et al [37] proposed formula used to quantify the amount of BNF by legumes in short-term cropping systems (Eq 1).

$$BNF = [N \text{ yield (legume)} - N \text{ yield (nonlegume)}] + [N \text{ soil (legume)} - N \text{ soil (nonlegume)}] \dots \dots \dots \text{Eq (1)}$$

All parameters included in the formula expressed by kg ha<sup>-1</sup>.

## 2.6 Proportion of Nitrogen Derived from the Atmosphere (%Ndfa)

The %Ndfa is the pure N which is contributed from the atmosphere by the N<sub>2</sub> fixing legume crop. The %Ndfa was calculated by adopting the formula of Rennie [38], which is used to estimate the %Ndfa (Eq 2).

$$\% Ndfa = \frac{[N \text{ yield (legume)} - N \text{ yield (nonlegume!)}]}{N \text{ yield (legume)}} \times 100 \dots \text{Eq (2)}$$

## 2.7 Net Nitrogen Balance (NNB)

Net nitrogen balance (NNB) is the difference between the biologically fixed nitrogen in legumes and the nitrogen removed from legume grains. This study assumed how much nitrogen would be added to the soil from biologically fixed nitrogen if groundnut stover residues were incorporated into the soil. Therefore, the NNB of the groundnut stover residues from biologically fixed nitrogen was calculated by subtracting the nitrogen removed by the groundnut grains from the biologically fixed nitrogen in legumes [39].

$$NNB \text{ (kg per ha)} = BNF \text{ (kg per ha)} - N \text{ removed by grain (kg per ha)} \dots \dots \dots \text{Eq (3)}$$

## 2.8 Statistics Analysis

The collected data were statistically analyzed using a two-way analysis of variance (ANOVA). Before data analysis, the collected data were tested for normality and homogeneity of variance at individual levels and in the combination of N

and P fertilizers. An analysis of variance was performed using the Genstat 18 edition. A significant difference between treatments was assessed using the Fisher's test for least significant difference (LSD) at a probability level of 0.05.

### 3. RESULTS AND DISCUSSION

#### 3.1 Before Planting Soil Physicochemical Properties of the Experimental Site

The result of the physicochemical analysis of the soil at the test site is shown in Table 2. The soil texture class of the test site was sandy loam, which was moderately suitable for plant growth. Therefore, sandy loam is suitable for growing groundnut because groundnut is usually grown on light-textured soils. The soil in the study area has a bulk density of  $1.46 \text{ g cm}^{-3}$ , which was classified as moderate ( $1.3\text{--}1.6 \text{ g cm}^{-3}$ ) [34]. The moderate bulk density of the soil with a sandy-loamy texture allows groundnut roots and pegs to easily penetrate the soil, absorb nutrients, absorb water and form pods. Furthermore, the pH value at the study site was 5.5, which corresponds to the moderately acidic category described by Tekalign [35]. As Murata and colleagues [40] studied, groundnut root and peg growth, pod formation, and overall plant development were best when soil pH was greater than or equal to five and declined rapidly when the pH value fell below this value. The low pH value of the soil also affects the nodulation ability of the groundnut [41]. The organic carbon content in the soil of the research area was 0.61% (Table 2). As stated by Hazelton and Murphy [34], a SOC value between 0.4 and 0.6% was considered low, and therefore the current value of soil organic carbon was low. Furthermore, the TN content in the research center was measured at 0.053% (Table 2), which is a low estimate [34]. The low SOC and TN content of the experimental site may have negatively affected groundnut production and it was necessary to add fertilizer to these soils. As shown in Table 2, the available soil phosphorus at the test site is 6.92 ppm. According to Hazelton and Murphy [34] review of soil test interpretations, the current result for soil available phosphorus is a low value (5–10 ppm). At the research area, soil EC was measured at  $2.94 \text{ dS m}^{-1}$  (Table 2), which belongs to slightly saline soils ( $2\text{--}4 \text{ dS m}^{-1}$ ) and at this level, saline soils can negatively affect the yield of salt-sensitive plants [34]. The content of exchangeable Na cations in the study area was

$0.16 \text{ cmol (+) kg}^{-1}$ , which is considered low ( $0.1\text{--}0.3 \text{ cmol (+) kg}^{-1}$ ) [34]. The exchangeable K content of the soil at the research site was  $0.21 \text{ cmol (+) kg}^{-1}$  (Table 2). According to the interpretation of the soil analysis by Hazelton and Murphy [34], the current exchangeable K result is in the low category ( $0.2\text{--}0.3 \text{ cmol (+) kg}^{-1}$ ). The exchangeable Ca and Mg contents in the soil at the study area were 5.8 and  $6.8 \text{ cmol (+) kg}^{-1}$ , respectively (Table 2). According to the category recommended by Hazelton and Murphy [34] for interpreting soil analysis results, the exchangeable Ca content in the experimental area is in the moderate category ( $5\text{--}10 \text{ cmol (+) kg}^{-1}$ ), while the value of the exchangeable Mg is high ( $3\text{--}8 \text{ cmol (+) kg}^{-1}$ ). Therefore, the Ca and Mg exchangeable cations of the experimental site are suitable for groundnut cultivation. The CEC of the research area was  $13.2 \text{ cmol (+) kg}^{-1}$ , which was considered moderate ( $12\text{--}25 \text{ cmol (+) kg}^{-1}$ ) [34]. The presence of moderate CEC in the soil creates a suitable soil environment for better groundnut growth.

#### 3.2 Nodulation

For the average number of nodules per plant, a significant effect was only observed for independently applied N fertilizer, while insignificant effects were observed for P fertilizer alone and the interaction of N fertilizers and P (Tables 3 and 4). The largest number of nodules per plant were observed in the plots treated with  $15 \text{ kg N ha}^{-1}$ , with no significant differences at the dose of  $30 \text{ kg N ha}^{-1}$ . The average number of nodules per plant decreased by 21.16% in the unfertilized plots compared to plots treated with  $15 \text{ kg N ha}^{-1}$ . Application of a higher dose of N fertilizer on groundnut plants can accelerate the disintegration of nodules, which is due to the inhibitory effect of nitrates on groundnut nodules and their development [42]. Application of high doses of N to legumes resulted in poor root growth and a reduction in flavonoid synthesis, thereby inhibiting rhizobia recognition and infection of legumes [43]. Confirming the current results, Li et al. [44] and Salvagiotti et al. [45] also observed a reduction in nodule number in faba beans and soybeans with increasing nitrogen rates. For nodule initiation of common bean,  $20 \text{ kg ha}^{-1}$  starter N was required [46]. Consistent observations from different studies have shown the negative effects of higher N doses on the nodulation and BNF of legumes [17].

**Table 3. Analysis of variance for groundnut nodulation, biological nitrogen fixation, Nitrogen derived from the atmosphere and net nitrogen balance as affected by starter N and P fertilizers and their interaction effects**

Source of variation	Groundnut traits	DF	SS	MS	P level	SE	CV (%)
N fertilizer	NN	2	2197.1	1098.6	0.05	17.78	17.8
	BNF	2	1699.2	849.6	0.093	17.97	24.9
	Ndfa	2	1250.61	625.30	<.001	7.81	21.2
	NNB	2	3866.5	1933.2	<.001	4.62	16.8
P fertilizer	NN	3	698.2	232.7	0.54	17.78	17.8
	BNF	3	9090.9	3030.3	<.001	17.97	24.9
	Ndfa	3	1939.37	646.46	<.001	7.81	21.2
	NNB	3	2146.7	715.6	<.001	4.62	16.8
N * P	NN	6	1137.8	189.6	0.73	17.78	17.8
	BNF	6	809.3	134.9	0.86	17.97	24.9
	Ndfa	6	443.93	73.99	0.33	7.81	21.2
	NNB	6	6586.7	1097.8	<.001	4.62	16.8
Residual	NN	24	7583.3	316.0			
	BNF	24	7747.2	322.8			
	Ndfa	24	1463.36	60.97			
	NNB	24	511.93	21.33			
Total	NN	35	11616.4				
	BNF	35	19346.6				
	Ndfa	35	5097.27				
	NNB	35	13111.73				

NN-number of nodules, DF- degree of freedom, SS- sum of squares, MS- mean squares, P level- provability level, SE- standard error and CV- coefficient of variation.

**Table 4. Groundnut nodulation, nitrogen derived from the atmosphere and net nitrogen balance affected by N fertilizer**

N (kg ha <sup>-1</sup> )	NN	Ndfa (%)	NNB (kg ha <sup>-1</sup> )
0	89.65 <sup>b</sup>	42.99 <sup>a</sup>	-17.34 <sup>a</sup>
15	108.62 <sup>a</sup>	38.72 <sup>a</sup>	-23.25 <sup>b</sup>
30	101.33 <sup>ab</sup>	28.91 <sup>b</sup>	-41.67 <sup>c</sup>
LSD (0.05)	14.98	6.58	3.89

Means followed by the same letters in the same column are not significantly different at ( $P \leq 0.05$ ), NN- number of nodules per plant, LSD-least significant difference.

**Table 5. Groundnut biological nitrogen fixation, nitrogen derived from the atmosphere and net nitrogen balance affected by N fertilizer**

P <sub>2</sub> O <sub>5</sub> (kg ha <sup>-1</sup> )	BNF (kg ha <sup>-1</sup> )	Ndfa (%)	NNB (kg ha <sup>-1</sup> )
0	71.30 <sup>a</sup>	40.02 <sup>a</sup>	-28.11 <sup>b</sup>
23	82.98 <sup>a</sup>	42.71 <sup>a</sup>	-15.64 <sup>a</sup>
46	87.75 <sup>a</sup>	40.48 <sup>a</sup>	-28.61 <sup>b</sup>
69	46.68 <sup>b</sup>	24.28 <sup>b</sup>	-37.31 <sup>c</sup>
LSD (0.05)	17.48	7.60	4.493

Means followed by the same letters in the same column are not significantly different at ( $P \leq 0.05$ ), LSD-least significant difference.

### 3.3 Biological Nitrogen Fixation (BNF)

No significant effect was observed on the interaction of N and P or on the separate effect of nitrogen fertilizer, while only P fertilizer had a

significant effect on the BNF of groundnut legume (Tables 3 and 5). When applying P<sub>2</sub>O<sub>5</sub> at a dose of 46 kg ha<sup>-1</sup>, a higher average BNF (87.75 kg ha<sup>-1</sup>) was observed, which was not significantly different from the application of P<sub>2</sub>O<sub>5</sub>

(0.23 kg ha<sup>-1</sup>). Mean BNF consistency increased for plots receiving 46 kg P<sub>2</sub>O<sub>5</sub> but decreased significantly at 69 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup>. The average BNF achieved after the application of 46 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> was 87.98% and higher than the P<sub>2</sub>O<sub>5</sub> applied at a dose of 69 kg ha<sup>-1</sup>. In fact, legumes can capture nitrogen from the atmosphere through a mutualistic symbiosis with some bacteria. However, to take advantage of this role, legumes require optimal phosphorus for energy conversion in the nodules. In addition, P plays an important role in the development of a healthy root, thereby improving plants' water and nutrient absorption. Confirming the current result, Hernandez et al. [47] reported that P significantly influences the growth and metabolism of legumes. Tropical legumes enriched with phosphate fertilizers have a higher BNF [46]. Although BNF from legumes has played an undeniably important role in global food security, its maximum utilization is limited by P availability [11]. Under conditions of low soil P availability, the effectiveness of BNF in legumes is reduced [48–53]. Accordingly, Ssali and Keya [54] found increased nitrogen and BNF yields in beans after P application.

### 3.4 Proportion of Nitrogen Derived from the Atmosphere (%Ndfa)

A significant influence on the %Ndfa was found when N and P fertilizers were applied separately, but not when N and P fertilizers were applied in

combination (Tables 3, 4 and 5). As the N fertilizer dose increased, the %Ndfa continuously decreased. A significantly higher amount of %Ndfa was found in plots not fertilized with N compared to plots receiving 30 kg N ha<sup>-1</sup>; however, no significant difference with plots receiving 15 kg N ha<sup>-1</sup>. Plots treated with 30 kg N ha<sup>-1</sup> were characterized by a lower Ndfa of 48.70 and 33.93% compared to unfertilized plots and plots treated with 15 kg N ha<sup>-1</sup>, respectively. Based on the results, it was observed that the N fixing capacity of groundnut plants decreases when fertilized with a higher dose of N and P fertilizers because the nutrient requirement of groundnut plants depends on the fertilizer applied. Therefore, Table 7 shows that BNF and Ndfa have a positive connection. Legumes sequester more N from the atmosphere when grown with little or no N fertilizer. The amount of %Ndfa can be negatively affected by increasing fertilizer rates, although groundnut kernel yield was higher with higher N fertilizer rates [9]. The BNF of alfalfa was reduced and delayed due to high N input [53]. The inclusion of legumes in agricultural production, such as crop rotation and green manure, can reduce the dependence on the use of N fertilizers as they have the potential to fix atmospheric N [54]. Legumes fix more N from the atmosphere when the soil is nutrient poor [1,55]. Legumes grown on N-rich soils or fertilized with large amounts of N fertilizer inhibited the growth of legume nodules and thus reduced BNF [1,45].

**Table 6. Interaction effect of N and P fertilizer rates on net nitrogen balance of groundnut**

N (kg ha <sup>-1</sup> )	P <sub>2</sub> O <sub>5</sub> (kg ha <sup>-1</sup> )			
	0	23	46	69
	<b>Net nitrogen balance</b>			
0	-9.18 <sup>a</sup>	-8.33 <sup>a</sup>	-19.32 <sup>b</sup>	-32.53 <sup>c</sup>
15	-4.04 <sup>a</sup>	-6.32 <sup>a</sup>	-35.73 <sup>c</sup>	-46.90 <sup>d</sup>
30	-71.12 <sup>e</sup>	-32.28 <sup>c</sup>	-30.79 <sup>c</sup>	-32.51 <sup>c</sup>
LSD (0.05)	7.78			

Means followed by the same letters across all columns and rows are not significantly different at (P≤0.05), LSD-least significant difference.

**Table 7. Correlation between groundnut nodulation, biological nitrogen fixation, nitrogen derived from the atmosphere and net nitrogen balance**

	NN	BNF	Ndfa	NNB
NN	1			
BNF	0.18 <sup>NS</sup>	1		
Ndfa	0.07 <sup>NS</sup>	0.80 <sup>***</sup>	1	
NNB	0.08 <sup>NS</sup>	0.39 <sup>*</sup>	0.73 <sup>***</sup>	1

Stars indicate significant effects \*p (probability level) ≤ 0.05, \*\*p ≤ 0.01, \*\*\*p ≤ 0.001, NS- non-significant, NN-number of nodules per plant.



Likewise, a separate application of P fertilizer had a significant impact on Ndfa percentage. Unfertilized plots and plots with 23 and 46 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> remained significantly similar but were characterized by a higher Ndfa of 64.83, 75.91 and 66.72%, respectively, compared to plots with 69 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup>. When soil P deficiency was alleviated by adding chemical P fertilizers, the %Ndfa in legumes improved [56]. Phosphorus nutrient significantly influences the growth of legumes and their symbiotic relationship with soil rhizobia due to the specific effect of P on the growth of rhizobia in soil, their survival and their nodulation ability [19].

### 3.5 Net N Balance (NNB)

It was noted that applying N and P fertilizers separately and combining them resulted in a significant change in the groundnut's net soil nitrogen balance (Tables 3, 4, 5 and 6). It is important to note that after applying 15 kg ha<sup>-1</sup> of nitrogen at a zero P level, a decrease in N export from groundnut seeds was observed. This dosage was then followed by a combined application of 15 kg N ha<sup>-1</sup> and 23 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup>, but was still very close to the combined application of zero N with four P<sub>2</sub>O<sub>5</sub> values (0, 23, 46 and 69 kg ha<sup>-1</sup>) and a combined application of 30 kg N ha<sup>-1</sup> with three P<sub>2</sub>O<sub>5</sub> levels (23, 46 and 69 kg ha<sup>-1</sup>). The greatest removal of N by groundnut kernels was observed at fertilization of 30 kg N ha<sup>-1</sup> at a constant P zero level. The current result shows that the application of high N fertilizers to groundnut plants reduced the amount of biologically fixed N since legumes rely on the chemical fertilizer used. The smaller negative numbers suggested that groundnut plants used little N from the soil and that most of the N requirement came from biologically fixed N in the atmosphere. Positive values provide a useful indication of the potential input of atmospheric nitrogen into the system, but negative values indicate the removal of N from the soil. Current results show that all biologically fixed N and some soil N are exported by the groundnut grains. A positive net addition of N to the soil by legumes can be expected if the Ndfa percentage is above the nitrogen harvest index (NHI) by grains. Opposing the current results, Kermah et al. [1] reported that groundnut contributes an average of 2 kg ha<sup>-1</sup> of net nitrogen to the soil when only grains are exported. The negative N balance in legume soil reflects a lower Ndfa percentage than NHI [57]. Peoples and Craswell [58] found that although the amount of BNF produced by legumes may be

high, it is still not sufficient to compensate for the N exported by the removed seeds. Higher values often mean that while the groundnut kernel removed substantial amounts of N from the soil has been absorbed rather than contributed to the soil system.

Similar to the interactions between N and P fertilizers, N fertilizer alone had a significant influence on the net N balance of groundnut (Tables 3 and 4). Lower N removal by groundnut kernels was observed in the unfertilized groundnut plot and was significantly similar to that applied at a rate of 15 kg N ha<sup>-1</sup>. The net N balance measured in plots not treated with N fertilizer increased by 140% compared to plots fertilized with 30 kg N ha<sup>-1</sup>. The decreasing trend in net N balance compared to groundnut with increasing N fertilization could be because the high N content in the root zone of groundnut kills the nodules, thus reducing the BNF, and the crop also depends on the N fertilizer applied for growth. A small nitrogen fertilizer application rate boosted plant growth, nitrogenase activity, nodule development, and BNF [59]. The overuse of chemical fertilizers on legume crops has caused a serious disruption in the nitrogen cycle, leading to nitrate accumulation in soil and water and nitrogen oxide pollution of the atmosphere [60]. The soil's NNB was significantly impacted by P's main effect. Plots receiving 23 kg ha<sup>-1</sup> P<sub>2</sub>O<sub>5</sub> showed better soil NNB, but plots receiving 69 kg ha<sup>-1</sup> P<sub>2</sub>O<sub>5</sub>, which is the higher rate, demonstrated low soil NNB. P availability in the legume crops' rhizosphere is essential for increasing BNF, which improves soil NNB [61-63]. Low soil N mining occurred from the stimulation of groundnut BNF by P fertilizer applied at 23 kg ha<sup>-1</sup>. Table 7 displays the positive correlation between BNF and NNB.

## 4. CONCLUSION

In conclusion, based on the results obtained, it was observed that groundnut nodulation, BNF, %Ndfa and net N balance improved with little or no fertilizer application. Adding large amounts of nutrients to groundnut likely reduced the potential ability to fix atmospheric nitrogen, due to groundnut growth is dependent on applied fertilizers. Therefore, legumes can reduce N fertilizer costs for smallholder farmers in developing countries. Furthermore, the risk of environmental pollution is low because legumes are included in the agricultural system due to low demand for external inputs. Therefore, overfertilizing groundnut crop excessively results

in environmental contamination and incurs extra costs.

### DISCLAIMER (ARTIFICIAL INTELLIGENCE)

The author declared that no generative AI tools, such as text-to-image generators or large language models (ChatGPT, COPILOT, etc.), were used in the writing or editing of the paper.

### DATA AVAILABILITY

The author states that, depending on the request, he can submit the data whenever it is in need. Upon a reasonable request, the corresponding author will provide the available datasets used and analyzed during the study.

### ACKNOWLEDGEMENTS

The author would like to thank his employer, Tigray Agricultural Research Institute (TARI), for financial support in conducting this study. I want to express my gratitude to Gebeyehu Taye, my advisor, for everything he has done to support the success of my thesis. I also extend my special thanks to the Shire Soil Research Center and the Mekelle Soil Research Center for providing laboratory facilities for soil and plant analysis, and to Mekelle University for enabling Mr. Kinfe to pursue the Master's degree program in Soil Sciences. Finally, I would like to thank local farmers for their support in providing experimental land and for their labor contributions.

### COMPETING INTERESTS

Author has declared that no competing interests exist.

### REFERENCES

1. Kermah, M, Franke, AC, Adjei-Nsiah, S, Ahiabor, BD, Abaidoo, RC, Giller, KE. N<sub>2</sub>-fixation and N contribution by grain legumes under different soil fertility status and cropping systems in the Guinea Savanna of Northern Ghana. *Agriculture, ecosystems & environment*. 2018;261: 201-210.
2. Woźniak, A. Effect of crop rotation and cereal monoculture on the yield and quality of winter wheat grain and on crop infestation with weeds and soil properties. *International Journal of Plant Production*. 2019; 13(3): 177-182.
3. Bationo, A., Ntare, BR. Rotation and nitrogen fertilizer effects on pearl millet, cowpea and groundnut yield and soil chemical properties in a sandy soil in the semi-arid tropics, West Africa. *the Journal of Agricultural Science*. 2000;134(3): 277-284.
4. Boincean, B, Dent, D, Boincean, B, Dent, D. Crop rotation. farming the black earth: sustainable and climate-smart management of chernozem soils. 2019;89:124.
5. Stevenson FJ. Origin and distribution of nitrogen in soil. *Soil nitrogen*. 1965 Jan 1;10:1-42.
6. Kebede E. Grain legumes production and productivity in Ethiopian smallholder agricultural system, contribution to livelihoods and the way forward. *Cogent Food & Agriculture*. 2020 Jan 1;6(1):1722353
7. Walley, FL, Clayton, GW, Miller, PR, Carr, PM, Lafond, GP. Nitrogen economy of pulse crop production in the northern great plains. *Agron. J*. 2007;99: 1710–1718.
8. O'Donovan, JT, Grant, CA, Blackshaw, RE, Harker, KN, Johnson, E, Gan, Y, Lafond, GP, May, WE, Turkington, TK, Lupwayi, NZ, Stevenson, FC. Rotational effects of legumes and non-legumes on hybrid canola and malting barley. *Agronomy Journal*. 2014; 106(6):1921-1932.
9. Ncube, B, Twomlow, SJ, Van Wijk, MT, Dimes, JP, Giller, KE. Productivity and residual benefits of grain legumes to sorghum under semi-arid conditions in southwestern Zimbabwe. *Plant and Soil*. 2007; 299(1-2):1-15.
10. Siddique KH, Johansen C, Rao JV, Ali M. Legumes in sustainable cropping systems. 2005;787-819.
11. Tekulu K, Taye G, Assefa D. Effect of starter nitrogen and phosphorus fertilizer rates on yield and yield components, grain protein content of groundnut (*Arachis Hypogaea* L.) and residual soil nitrogen content in a semiarid north Ethiopia. *Heliyon*. 2020;6(10):1-12
12. Kaewpradit, W, Toomsan, B, Cadisch, G, Vityakon, P, Limpinuntana, V, Saenjan, P, Jogloy, S Patanothai, A mixing groundnut residues and rice straw to improve rice yield and n use efficiency. *Field Crops Research*. 2009;110(2):130-138.
13. Vance CP, Graham PH, Allan DL. Biological nitrogen fixation: phosphorus-a

- critical future need? Nitrogen fixation: From Molecules to Crop Productivity. 2000;509-514.
14. Bado BV, Bationo A, Lompo F, Traore K, Sedogo MP, Cescas MP. Long term effects of crop rotations with fallow or groundnut on soil fertility and succeeding sorghum yields in the Guinea Savannah of West Africa. Lessons learned from long-term soil fertility management experiments in Africa. 2012;27-40.
  15. Toomsan, B, McDonagh, JF, Limpinuntana, VJHA, Giller KE. Nitrogen fixation by groundnut and soya bean and residual nitrogen benefits to rice in farmers' fields in northeast Thailand. Plant and Soil. 1995;175(1):45-56.
  16. Yakubu,H, Kwari, JD, Sandabe, MK. Effect of phosphorus fertilizer on nitrogen fixation by some grain legume varieties in sudano-sahelian zone of north eastern Nigeria. Nigerian Journal of Basic and Applied Sciences. 2010;18(1):44-49.
  17. Bado, BV, Bationo, A, Cescas, MP. Assessment of cowpea and groundnut contributions to soil fertility and succeeding sorghum yields in the guinean savannah zone of Burkina Faso (West Africa). Biology and Fertility of Soils. 2006;43(2):171-176.
  18. Van Kessel, C, Hartley, C. Agricultural management of grain legumes: Has it led to an increase in nitrogen fixation? *Field Crops Research*. 2000;65(2-3):165-181.
  19. Huang J, Keshavarz Afshar R, Tao A, Chen C. Efficacy of starter n fertilizer and rhizobia inoculant in dry pea (*Pisum sativum* Linn.) production in a semi-arid temperate environment. Soil Science and Plant Nutrition. 2017;63(3):248-253.
  20. Prasad PV, Kakani VG, Upadhyaya HD. Growth and production of groundnut. UNESCO Encyclopedia. 2010;1-26.
  21. McLaughlin MJ, Malik KA, Memon KS, Idris M. The role of phosphorus in nitrogen fixation in upland crops. 1990:295-305.
  22. Mitran T, Meena RS, Lal R, Layek J, Kumar S, Datta R. Role of soil phosphorus on legume production. Legumes for soil health and sustainable management. 2018;487-510.
  23. Hussain, RM. The effect of phosphorus in nitrogen fixation in legumes. Agricultural Research & Technology. 2017;5(1):12–14.
  24. Angus, JF, Kirkegaard JA, Hunt, JR, Ryan, MH, Ohlander, L, Peoples, MB. Break crops and rotations for wheat. Crop and Pasture Science. 2015;66(6):523-552.
  25. Peterson, TA, Russelle, MP. Alfalfa and the nitrogen cycle in the corn belt. Journal of Soil and Water Conservation. 1991;46(3):229-235.
  26. Graham, PH, Vance, CP. Legumes: Importance and constraints to greater use. Plant physiology. 2003;131(3):872-877, 2003.
  27. National Meteorological Agency of Federal Democratic Republic of Ethiopia (NMA). Mekelle; Tigray; Ethiopia; 2019.
  28. Cissé L. Balanced fertilization for sustainable use of plant nutrients. Fertilizer Best Management Practices. 2007;33.
  29. Bouyoucos GJ. Hydrometer method improved for making particle size analyses of soils 1. Agronomy Journal. 1962;54(5):464-5.
  30. Rhoades JD, Chanduvi F, Lesch SM. Soil salinity assessment: Methods and interpretation of electrical conductivity measurements. Food & Agriculture Org.; 1999.
  31. Richards LA, editor. Diagnosis and improvement of saline and alkali soils. US Government Printing Office; 1954.
  32. Walkley A, Black IA. An examination of the Degtjareff method for determining soil organic matter, and a proposed modification of the chromic acid titration method. Soil science. 1934;37(1):29-38.
  33. Bremner JM, Mulvaney CS. Nitrogen—total. Methods of soil analysis: part 2 chemical and microbiological properties. 1982;9:595-624..
  34. Olsen SR. Estimation of available phosphorus in soils by extraction with sodium bicarbonate. US Department of Agriculture; 1954.
  35. Polemio M, Rhoades JD. Determining cation exchange capacity: A new procedure for calcareous and gypsiferous soils. Soil Science Society of America Journal. 1977;41(3):524-8.
  36. Hazelton P, Murphy B. Interpreting soil test results: What do all the numbers mean?. CSIRO publishing; 2016.
  37. Tadesse T, Haque I, Aduayi EA. Soil, plant, water, fertilizer, animal manure and compost analysis manual. 1991.
  38. Plank CO. Reference plant analysis procedures for the southern region of the United States. Southern Co-operative Series Bulletin. 1992;368:71-73.

39. Peoples MB, Faizah AW, Rerkasem B, Herridge DF. Methods for evaluating nitrogen fixation by nodulated legumes in the field. 1989.
40. Rennie RJ. Comparison of N Balance and <sup>15</sup>N Isotope Dilution to Quantify N<sub>2</sub> Fixation in Field-Grown Legumes 1. *agronomy Journal*. 1984;76(5):785-790.
41. Peoples MB. 3.3. Nitrogen balance in cropping systems that include legumes. *Guidelines on Nitrogen Management in Agricultural Systems*. 2008;160.
42. Murata, MR, Zharare, GE, Hammes, PS. pH of the pod-zone affects reproductive growth of groundnut. *Journal of plant nutrition*. 2007;31(1):69-79.
43. Uguru, MI, Ezeh, NE. Growth, nodulation and yield of bambara groundnut (*Vigna subterranea* (L) Verdc) on selected Nigerian soils. *Journal of the Science of Food and Agriculture*. 1997;73(3):377-382.
44. Daimon H, Yoshioka, M. Responses of root nodule formation and nitrogen fixation activity to nitrate in a split-root system in peanut. *Journal of Agronomy and Crop Science*. 2001;187(2):89-95.
45. Basu M, Bhadoria, PBS, Mahapatra SC. Growth, nitrogen fixation, yield and kernel quality of peanut in response to lime, organic and inorganic fertilizer levels. *Bioresource technology*. 2008;99(11):4675-4683.
46. Li YY, Yu CB, Cheng X, Li CJ, Sun JH, Zhang FS, Lambers H, Li L. Intercropping alleviates the inhibitory effect of N fertilization on nodulation and symbiotic N<sub>2</sub> fixation of faba bean. *Plant and Soil*. 2009;323(1-2):295-308.
47. Salvagiotti F, Cassman KG, Specht JE, Walters DT, Weiss A, Dobermann A. Nitrogen uptake, fixation and response to fertilizer N in soybeans: A review. *Field Crops Research*. 2008;108(1):1-3.
48. Daba, S, Haile, M. Effects of rhizobial inoculant and nitrogen fertilizer on yield and nodulation of common bean. *Journal of plant nutrition*. 2000; 23 (5): 581-591.
49. Hernández G, Ramírez M, Valdés-López O, Tesfaye M, Graham MA, Czechowski T, Schlereth A, Wandrey M, Erban A, Cheung F, Wu HC. Phosphorus stress in common bean: root transcript and metabolic responses. *Plant physiology*. 2007;144(2): 752-767.
50. Zheng M, Li D, Lu X, Zhu X, Zhang W, Huang J, Fu S, Lu X, Mo J. Effects of phosphorus addition with and without nitrogen addition on biological nitrogen fixation in tropical legume and non-legume tree plantations. *Biogeochemistry*. 2016;13(1-2):65-76.
51. Bonilla I, Bolanos L. Mineral nutrition for legume-rhizobia symbiosis: B, Ca, N, P, S, K, Fe, Mo, Co, and Ni: A review. *Organic Farming, Pest Control and Remediation of Soil Pollutants: Organic farming, pest control and remediation of soil pollutants*. 2010;253-274.
52. Kleinert A, Thuynsma R, Magadlela A, Benedito VA, Valentine AJ. Metabolism and transport of carbon in legume nodules under phosphorus deficiency. *Legume Nitrogen Fixation in Soils with Low Phosphorus Availability: Adaptation and Regulatory Implication*. 2017:77-95.
53. Püschel D, Janoušková M, Voříšková A, Gryndlerová H, Vosátka M, Jansa J. Arbuscular mycorrhiza stimulates biological nitrogen fixation in two medicago spp. through improved phosphorus acquisition. *Frontiers in plant science*. 2017;8:390.
54. Ssali H, Keya SO. The effect of phosphorus on nodulation, growth and dinitrogen fixation by beans. *Biological Agriculture & Horticulture*. 1983;1(2):135-144.
55. Xie KY, LI XL, Feng HE, Zhang YJ, Wan LO, David BH, Dong Wang, Yan QIN, Gamal MF. Effect of nitrogen fertilization on yield, n content, and nitrogen fixation of alfalfa and smooth brome grass grown alone or in mixture in greenhouse pots. *Journal of Integrative Agriculture*. 2015;14(9):1864-1876.
56. Jensen ES, Peoples MB, Boddey RM, Gresshoff PM, Hauggaard-Nielsen H, Alves BJ, Morrison MJ. Legumes for mitigation of climate change and the provision of feedstock for biofuels and biorefineries. A review. *Agronomy for sustainable development*. 2012;32(2):329-364.
57. Giller KE. *Nitrogen fixation in tropical cropping systems*. Cabi; 2001.
58. Cadisch G, Sylvester-Bradley R, Boller BC, Nösberger J. Effects of phosphorus and potassium on N<sub>2</sub> fixation (<sup>15</sup>N-dilution) of field-grown *Centrosema acutifolium* and *C. macrocarpum*. *Field Crops Research*. 1993;31(3-4):329-40.
59. Kermah M, Franke AC, Ahiabor BD, Adjei-Nsiah S, Abaidoo RC, Giller KE. Legume–

- maize rotation or relay? Options for ecological intensification of smallholder farms in the Guinea savanna of northern Ghana. *Experimental Agriculture*. 2019;55(5):673-91.
60. Peoples MB, Craswell ET. Biological nitrogen fixation: investments, expectations and actual contributions to agriculture. *Plant and soil*. 1992;141:13-39.
61. Tsai SM, Bonetti R, Agbala SM, Rossetto R. Minimizing the effect of mineral nitrogen on biological nitrogen fixation in common bean by increasing nutrient levels. *Enhancement of Biological Nitrogen Fixation of Common Bean in Latin America: Results from an FAO/IAEA Co-ordinated Research Programme*. 1986–1991, 1993;131-8.
62. Raza A, Zahra N, Hafeez MB, Ahmad M, Iqbal S, Shaukat K, Ahmad G. Nitrogen fixation of legumes: Biology and Physiology. *The plant family Fabaceae: biology and physiological responses to environmental stresses*. 2020;43-74.
63. Janati W, Benmrid B, Elhaissoufi W, Zeroual Y, Nasielski J, Bargaz A. Will phosphate bio-solubilization stimulate biological nitrogen fixation in grain legumes? *Frontiers in Agronomy*. 2021;3: 637196.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*

<https://www.sdiarticle5.com/review-history/120966>