



Effect of Combined Inoculation of Phosphate Solubilizing Bacteria and Endomycorrhizae Fungi on Microbial Population and NPK Uptake by Wheat (*Triticum aestivum* L.) Crop

Pooja Rani¹, Mahendra Singh^{1*} and Manoj Kumar Dwivedi¹

¹*Department of Soil Science and Agricultural Chemistry, Bihar Agricultural University, Sabour, Bhagalpur-813210, Bihar, India.*

Authors' contributions

This work was carried out in collaboration between all authors. Author PR designed the study, performed the statistical analysis, wrote the protocol and wrote the first draft of the manuscript. Authors MS and MKD managed the analyses of the study. Author MS managed the literature searches. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/CJAST/2019/46747

Editor(s):

(1) Dr. Yahya Elshimali, Professor, Department of Internal Medicine, Charles Drew University of Medicine and Science, USA.

Reviewers:

(1) P. Saravana Kumari, Rathnavel Subramaniam College of Arts and Science, India.
(2) Martín María Silva Rossi, Argentina.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/46747>

Original Research Article

Received 23 October 2018
Accepted 14 January 2019
Published 11 February 2019

ABSTRACT

Aim: To study the response of co-inoculation of Phosphate Solubilizing Bacteria (PSB) and Arbuscular Mycorrhizal (AM) fungi to microbial population of soil and nutrient uptake by wheat crop .
Study Design: The used design was completely randomized design with three replications.
Place of Study: The pot experiment was conducted during *Rabi* season of 2017 taking wheat (cv. HD 2967) as test crop in experimental unit of the department of Soil Science and Agricultural Chemistry, Bihar Agricultural College, Sabour, Bhagalpur (Bihar).
Methodology: The present study includes eight treatments with three replications. N P K uptake by wheat crop was measured from each treatment and microbial populations of soil were determined from the rhizospheric soils collected from each treatment by using standard protocol.
Results: Maximum microbial population viz., Bacteria, Actinomycetes, Fungi (39.00, 21.33, 24.66

*Corresponding author: E-mail: m.singh30648@gmail.com;

CFU $\times 10^5$ g⁻¹ oven dry soil) were recorded under treatment T₄ (T₁+PSB @ 20 g kg⁻¹ seed+AM fungi @ 5.0 g pot⁻¹) for bacteria and actinomycetes and T₃ (T₁+AM fungi @ 5.0 g pot⁻¹) for fungi, at flowering stage and similar trend was followed at harvesting stage. The treatment T₄ significantly increased available nitrogen, phosphorus and potassium in soil as well as also contributed to comparatively better plant growth and higher uptake of N, P and K by grain and shoot. The maximum N, P and K content of wheat was also recorded under treatment T₄ which was found to be most effective in modifying soil microbial population, microbial community structure and grain yield of wheat crop.

Conclusion: Application of co-inoculation of Phosphorus solubilizing bacteria and arbuscular mycorrhizal fungi enhance the microbial population and N P K uptake from soil by wheat crop.

Keywords: AM fungi; microbial population; phosphorus solubilizing bacteria; wheat.

1. INTRODUCTION

Phosphorus is critical element for plant growth and their development, and is a component of the nucleic acid structure of plants and biomembrane. Consequently, it is important in cell division and tissue development. Phosphorus is also involved in the energy metabolism of cells and is required for the biosynthesis of primary and secondary metabolites in plants. Therefore, plants have evolved a range of strategies to increase phosphorus uptake and mobility [1], the most common among which are phosphate solubilizing bacteria (PSB) and Arbuscular mycorrhiza (AM) fungi symbiosis.

Most of the applied phosphorus in the soils is not taken up by the crop, but it is retained in insoluble forms or fixed as mineral forms in the farms even as high as 90% or more. Soil phosphorus can exist in various inorganic (Pi) and organic forms (Po). Specific determination of Pi can be obtained by fractionation methods. Generally, plants take up P as the primary (H₂PO₄⁻) and secondary orthophosphate (HPO₄⁻) ions. They are easily retained in most soils when added, and in many cases this retention is so high that the element becomes largely unavailable to the plants. The phosphorus retained by the soil is generally considered as fixed P, although a part of it can be utilized. The rate of release of P from fixed form to replenish the immediately available soil P is the most important factor in determining the P supplying capacity of the soil because the quantity of P present in soil solution is not sufficient to meet the crop requirements. The release of fixed soil P depends upon the nature of its fixation and the extracting power of the crop or reagent used to determine availability of soil P. The P is generally fixed as Fe-P and Al-P in acidic soils and Ca-P in alkaline soils.

P-solubilization ability of the microorganisms is considered to be one of the most important traits associated with plant phosphorus (P) nutrition. Recently, phosphate solubilizing microorganisms have attracted the attention of agriculturists as soil inoculums to improve the plant growth and yield. Conversion of the insoluble forms of P to the form which is available to plants (ortho-phosphate) is an important characteristics of phosphate- solubilizing bacteria (PSB) and Arbuscular Mycorrhizal Fungi (AMF). Bacteria such as PSB and AM fungi are usually effective on phosphate solubility due to different mechanism such as production and secretion of organic acids and by their co-inoculation they make phosphorus available to plant for different metabolic functions [2]. Release of phosphorus by PSB from insoluble and fixed or adsorbed forms is an important aspect regarding P availability in soils. There are strong evidences that soil bacteria are capable of transforming soil P to the forms available to plant. AM fungi are species of fungi that intimately associate with plant roots forming asymbiotic relationship, with the plant providing sugars for the fungi and the fungi providing nutrients such as phosphorus, to the plants. Mycorrhizal fungi can absorb, accumulate and transport large quantities of phosphate within their hyphae and release to plant cells in root tissue. Arbuscular mycorrhizal (AM) fungus plant relationships are usually described as mutually beneficial, because fungi supply mineral nutrients, especially phosphorus (P) to their host plants in return for photosynthates. The contribution of AM fungi to P uptake by positively responsive plants can be easily identified by comparison of P uptake in AM plants and non-mycorrhizal (NM) plants. The beneficial effect of arbuscular mycorrhiza on plant growth is mainly attributed to higher P uptake by plants [3].

This is the fact that phosphorus solubilizing bacteria and mycorrhizal fungi increase the availability of phosphorus. There is a great scope to make more P availability in soil by adopting new agronomic practices like use of co-inoculation of PSB and AMF with its appropriate dose, proper combinations and inorganic fertilizer etc. Considering above mentioned facts a pot experiment on response of Co-inoculation of Phosphorus solubilizing bacteria (PSB) and Arbuscular Mycorrhizal (AM) Fungi on phosphorus availability under wheat rhizosphere has been conducted.

2. MATERIALS AND METHODS

The present study was undertaken in pot to evaluate the response of co-inoculation of Phosphate Solubilizing Bacteria (PSB) and Arbuscular Mycorrhizal (AM) fungi to soil microbial population and availability of phosphorus under wheat rhizosphere during the *Rabi* season of 2017 with a promising var. HD 2967, at the experimental site of department of Soil Science and Agricultural Chemistry, Bihar Agricultural University, Sabour, Bhagalpur, India.

The microbial inoculums viz., PSB- *Burkholderia cepacia* and AM fungi- *Glomus mosseae* were procured from biofertilizer production unit, Bihar Agricultural University, Sabour, Bhagalpur, and Bihar. The soil used for the pot experiment was Ustochrept clayey in texture, having a pH of 7.78 and EC of 0.20 ds m⁻¹. The organic carbon content of the soil was 0.47%, and the available nitrogen (N), phosphorus (P) and potassium (K) content was found 150.42, 12.51 and 192.66 kg ha⁻¹, respectively.

The following treatments with three replications were formulated for the study: T₁-RDF (120:60:40), T₂- T₁+PSB @ 20 g kg⁻¹ of seed, T₃- T₁+AM fungi @ 5.0 g pot⁻¹, T₄-T₁+PSB@20 g kg⁻¹ seed+AM fungi @ 5.0g pot⁻¹, T₅-75% RDF of Phosphorus +PSB@ 20 g kg⁻¹ seed, T₆-75% RDF of Phosphorus +AM fungi @ 5.0 g pot⁻¹, T₇-75% RDF of Phosphorus +PSB @ 20 g kg⁻¹ of seed+AM fungi@5.0 g pot⁻¹, and T₈-50% RDF of Phosphorus +PSB @ 20 g/kg seed+AM fungi @5.0 g Pot-1. Earthen pots of 15 cm height and 30 cm diameter were filled with 10 kg of soil. The seed treatment with PSB@ 20 kg⁻¹ seed was done and AM fungi@ 5 g Inoculum pot⁻¹ were applied 2 cm below the seed at the time of sowing. In each pot, 10 seeds of wheat (var. HD-2967) were planted. Nitrogenous, phosphatic and potassic fertilizers were applied just before the

sowing according to the treatments. Using urea, single super phosphate and murate of potash as source of nitrogen, phosphorus and potash, respectively and mixed in the soil uniformly by working with spade. The irrigation was applied as and when required the crops. The plants were thin to maintain eight plants in all pots.

Rhizosphere samples were drawn from the soil adhering to the roots. The 10 g of soil samples were placed in an Erlenmeyer flask containing 90 ml of sterilized distilled water, and shaken for 30 min. Ten-fold series dilutions were prepared, and appropriate dilutions were plated in specific media. For the isolation of bacteria, fungi and actinomycetes, the Plate Count Agar, Czapek-Dox Agar [4] and Kenknight and Munaier's Medium, respectively were used. The numbers of colony forming cells were determined in each plot by serial dilution pour plate method [5]. Phosphorus concentration in straw and grain were determined by employing the vanadomolybdate yellow color method given by [6] and the distribution of P in soil was determined by using a modified version of the Hedley and Stewart [7]. The obtained data from the pot experiment were analyzed with completely Randomized Design (CRD). The analysis of variance (ANOVA) was performed as described by Gome and Gomez [8] to determine the effects of various treatments. Critical difference (CD) at 5% level of probability and P values was used to examine differences among treatment means.

3. RESULTS AND DISCUSSION

3.1 Effect on Bacterial Population

It is evident from presented data that the microbial population resulted highest in the flowering stage of wheat plant growth. This might be due to accumulation of various root exudates and which in turn, established a strong and well defined root-microbe interaction [9]. The inoculation with treatment T₄(T₁+PSB @20 g kg⁻¹ seed + AM fungi @ 5.0g pot⁻¹) having significantly more bacterial population, when compared with applied treatments. It is clearly shown that all applied treatments have given maximum bacterial count at flowering stage when compared with harvesting stage. The application of treatment T₄(T₁+PSB @ 20 g kg⁻¹ seed + AM fungi @ 5.0 g pot⁻¹) also produced significantly higher bacterial count by 22.23%, 4.28%, 9.41%, than application of T₁ (RDF 120:60:40), T₂ (T₁ + PSB@20g kg⁻¹ seed), T₃ (T₁ + AM fungi @ 5.0 g pot⁻¹) respectively. The

similar trends were observed at harvesting stage. At harvesting stage application of treatment T₆ (75% RDF of P + AM fungi @ 5.0g pot⁻¹) and treatment T₇ (75% RDF of P + PSB @ 20 g kg⁻¹ seed + AM fungi @5.0 g pot⁻¹) given significantly higher colonization over all the applied treatments. Similar trend followed at harvesting stage. It might be due to the AM fungi are probably the most abundant fungi in agricultural soil accounting for somewhere between 5 and 50% of biomass of soil microbes live on carbohydrates obtained from the root cells. They alter root exudation considerably and are therefore expected to influence rhizosphere populations as well [10]. Numerous studies have shown conclusively that AM is having synergistic interaction with other beneficial soil micro-organism such as N fixers and P solubilizers. AM fungi affect the composition of bacterial communities either directly by changing host plant physiology or indirectly by changing the pattern of root exudation. The number of both rhizospheric bacteria and actinomycetes enhanced when plant formed mycorrhizae, along with the inoculation of PSB [11]. There may be two pathways for AM fungi to change microbe community structure, the first one is that the AM fungal hyphae secretion directly impacts microbe community structures; the another one is that both AM fungi in roots and on the roots alter plant physiological and biochemical processes, then directly or indirectly change the plant root secretion, thus alter those structures [12].

3.2 Effect on Actinomycetes Population

Same trend of enhancement of actinomycetes count has been resulted under the experimentation as compared as obtained in case of bacteria. The growth pattern was resulted synonymous to bacterial counterpart. The highest individual treatment effect was observed under the treatment T₄ (T₁ + PSB@20 g kg⁻¹ seed + AM fungi @ 5.0 g pot⁻¹) in flowering stage, produced maximum population of actinomycetes (21.33 CFU×10⁶ g⁻¹ soil). The treatment was significantly affected actinomycetes population in all the applied treatments except treatment T₂ (T₁+PSB @ 20g/kg of seed). Similar trends were followed at harvesting stage. The similar results were obtained by Sah et al. [13] who conducted a field experiment and found that the maximum bacterial population (71.66 CFU × 10⁵ g⁻¹ soil and 40.00 CFU × 10⁵ g⁻¹ soil), fungi population (27.33 CFU × 10⁴ g⁻¹ soil and 22.66 CFU × 10⁴

g⁻¹ soil) and actinomycetes population (57.66 CFU × 10⁵ g⁻¹ soil and 46.33 CFU × 10⁵ g⁻¹ soil) are observed in the treatment T₅ {(75 % N (FYM) basal + 25% N (V/C) at 25 DAT + *Azospirillum* @ 5 kg ha⁻¹ + PSB @ 5 kg ha⁻¹ + KSB @ 5 kg ha⁻¹)} at both the panicle and harvesting stage in rice.

3.3 Effect on Fungal Population

It is depicted from the data that the fungal count resulted highest in the flowering stage of growth as compared to harvesting stage. The application of treatment T₃ (T₁ + AM fungi @ 5.0 g pot⁻¹) given maximum population of fungi (24.66 propagules ×10³ g⁻¹ soil). This treatment also given significantly higher fungal population by 17.02%, 40.53%,16.21%, 44.59%, 12.16%, 16.21%, over application of treatment T₁ (RDF (120:60:40), T₂ (T₁+PSB@ 20 g kg⁻¹ seed), T₄ (T₁+PSB @ 20 g kg⁻¹ seed + AM fungi @ 5.0g pot⁻¹), T₅ (75% RDF of P + PSB @ 20g kg⁻¹ seed), T₆ (75% RDF of P + AM fungi @ 5.0g pot⁻¹) and T₇ (75% RDF of P + PSB @ 20g kg⁻¹ seed + AM fungi@5.0 g pot⁻¹) respectively. Similar trend observed at harvesting stage. At harvesting stage the maximum number of fungal population (15.667 propagules ×10³ g⁻¹ soil) was recorded with the application of treatment T₃ (T₁ + AM fungi @ 5.0 g pot⁻¹). It might be due accumulation of various root exudates and which in turn, established a strong and well defined root-microbe interaction [14] also as compared to bacteria and fungi. The similar results were found by the Singh and Kumar [15].

3.4 Effect on Available Nitrogen (kg ha⁻¹)

The maximum available nitrogen (210.24 kg ha⁻¹) was recorded under the application of treatment T₄ (T₁+PSB @ 20 g kg⁻¹ seed +AM fungi @ 5.0 g Pot⁻¹). It has been clearly observed that the available nitrogen in soil was significantly higher than all the applied treatments. It might be due to the production of more shoot biomass and root biomass by the application of given treatments and might be due to the structural changes in the microbial community. These changes in the microbial community may alter the nutrients dynamics in the rhizosphere.

3.5 Available Phosphorus (kg ha⁻¹)

The data regarding available phosphorus in soil revealed that the maximum phosphorus (19.25 kg ha⁻¹) was found by the application of treatment T₄ (T₁+PSB @ 20 g kg⁻¹ seed + AM fungi @ 5.0 g Pot⁻¹). The application of

treatment T_4 given significantly higher available phosphorus by 31.68%, 15.37%, 19.16%, 10.38% and 5.66% when compared with the treatment T_1 {(RDF (120:60:40)}, T_2 (T_1 +PSB @ 20 g kg^{-1} of seed), T_3 (T_1 +AM fungi @ 5.0 g Pot^{-1}), T_5 (75% RDF of P +PSB @ 20 g kg^{-1} seed), T_6 (75% RDF of P +AM fungi @ 5.0 g Pot^{-1}), and numerically the least value of available phosphorus was found under the treatment T_1 (RDF (120:60:40)). Recorded data shows that the inoculation with T_7 (75% RDF of P +PSB @ 20 g kg^{-1} of seed + AM fungi @ 5.0 g Pot^{-1}) having also more significantly availability of phosphorus by 28.72%, 11.70%, 15.66%, 6.50% and 22.76% over the treatments T_1 , T_2 , T_3 , T_5 and T_8 respectively. It may be due to the *Glomus mosseae* had pronounced effect for phosphorus acquisition in soil inoculated with PSB have a great result. [16] who conducted an experiment on *Coriander sativum* L. to study the effect of arbuscular mycorrhizal fungus *Glomus mosseae* and phosphorus application on plant growth rate, essential oil content and composition of coriander, and found that the mycorrhizal inoculation significantly increased growth responses and P and N plant nutrients in shoot and root tissue, also after inoculation of arbuscular mycorrhizal fungi in to coriander plant is a feasible alternative to increase growth, nutrition, essential oil production and reduce the use of P fertilizers required to obtain economic production of coriander under phosphorus deficient soil condition.

3.6 Available Potassium (kg ha⁻¹)

The maximum available potassium (210.07 kg ha^{-1}) in soil was recorded by the application of T_4 (T_1 +PSB @ 20 g kg^{-1} seed + AM fungi @ 5.0 g Pot^{-1}). This treatment also gave significantly higher amount of available potassium in soil by 13.66%, 11.68%, 12.81%, 12.25%, 12.13%, 4.72% over the treatments T_1 {(RDF (120:60:40)}, T_2 (T_1 + PSB @ 20 g kg^{-1} seed), T_5 (75% RDF of P +PSB @ 20 g kg^{-1} seed), T_6 (75% RDF of P +AM fungi @ 5.0 g Pot^{-1}), T_7 (75% RDF of P +PSB @ 20 g kg^{-1} of seed + AM fungi @ 5.0 g Pot^{-1}) and T_8 (50% RDF of P +PSB @ 20 g kg^{-1} seed + AM fungi @ 5.0 g Pot^{-1}) while the treatment T_3 (T_1 + AM fungi @ 5.0 g Pot^{-1}) given significantly zero value. It has been shown that in co-inoculated treatment of PSB and AM fungi having more amount of available potassium present with respect to un-inoculated condition like T_1 , T_2 , T_3 etc. Another treatment which is inoculated with T_8 given significantly higher

availability of potassium except T_3 and T_4 but another by 9.37%, 7.30%, 8.49% and 7.77% than T_1 , T_2 , T_5 , T_7 respectively.

3.7 Soil Organic Carbon (%)

The data revealed with percent organic carbon in soil showed that the application of all treatments were non-significant. The data regarding available organic carbon in a soil maximum (0.54%) was recorded by the application of treatment T_2 (T_1 + PSB @ 20g/kg of seed), T_3 (T_1 +AM fungi @5.0 g Pot^{-1}), T_4 (T_1 +PSB @20 g kg^{-1} seed +AM fungi @ 5.0 g Pot^{-1}), T_5 (75% RDF of P +PSB @ 20 g kg^{-1} seed), T_6 (75% RDF of P +AM fungi @ 5.0 g Pot^{-1}), T_7 (75% RDF of P +PSB @ 20 g kg^{-1} of seed + AM fungi@5.0 g Pot^{-1}) and T_8 (50% RDF of P + PSB @ 20 g kg^{-1} seed + AM fungi @5.0 g Pot^{-1}). It might be due to the AM colonization produced more root biomass and plant biomass.

3.8 Effect of Microbial Inoculants on N P K Content and Uptake by Wheat

The data regarding N P K content and uptake in shoot and grain of wheat clearly depicted that the application of treatment T_4 (T_1 +PSB @ 20 g kg^{-1} seed + AM fungi @ 5.0 g Pot^{-1}) given maximum N P K content and their uptake over all the applied treatments, which is found to be significantly higher over the applied treatment T_1 {(RDF (120:60:40)}, T_2 (T_1 + PSB @ 20 g kg^{-1} seed), T_5 (75% RDF of P +PSB @ 20 g kg^{-1} seed), T_6 (75% RDF of P +AM fungi @ 5.0 g Pot^{-1}), T_7 (75% RDF of P +PSB @ 20 g kg^{-1} seed + AM fungi @ 5.0 g Pot^{-1}) and T_8 (50% RDF of P + PSB @ 20 g kg^{-1} seed + AM fungi @ 5.0 g Pot^{-1}).

The data pertaining N content (%) has been clearly observed that the nitrogen content percentage in shoot was maximum (0.149%) with the treatment T_4 (T_1 +PSB@20 g kg^{-1} seed+AM fungi @ 5.0g pot^{-1}), which is significantly zero over the applied treatment T_1 {(RDF (120:60:40)}, T_2 (T_1 +PSB @ 20 g kg^{-1} of seed), T_3 (T_1 +AM fungi@5.0 g pot^{-1}), T_5 (75% RDF of Phosphorus +PSB @ 20g kg^{-1} seed), T_6 (75% RDF OF Phosphorus +AM fungi @ 5.0g pot^{-1}) and T_7 (75% RDF of Phosphorus +PSB @ 20g kg^{-1} of seed+AM fungi@5.0 g pot^{-1}) The lowest nitrogen content (0.140%) was found under the treatment T_1 . While the treatment T_3 given significantly higher value by 2.75%, 2.06%, 2.7% over the applied treatment T_1 , T_2 and T_5 respectively while T_4 and T_8 are at par values. In the same way treatment T_8 (50% RDF of

Phosphorus +PSB @ 20g kg⁻¹ seed+AM fungi @ 5.0 g pot⁻¹) have given significantly higher value by 5.40%, 4.05%, 2.02%, 4.72%, 2.70%, over the applied treatment T₁, T₂, T₃, T₅ and T₇ while the treatment T₄ have at par values.

The data pertaining P content (%) has been clearly observed that the phosphorus content percentage was maximum (0.245%) with the treatment T₄(T₁+PSB @ 20 g kg⁻¹ seed+AM fungi @ 5.0 g pot⁻¹), which is found to be significantly higher over the applied treatment T₁{(RDF (120:60:40)), T₂(T₁+PSB @ 20 g kg⁻¹ of seed) T₃(T₁+AM fungi @ 5.0 g pot⁻¹), T₅(75% RDF of Phosphorus +PSB @ 20 g kg⁻¹ seed), T₆(75% RDF OF Phosphorus +AM fungi @ 5.0 g pot⁻¹), T₇(75% RDF of Phosphorus +PSB @ 20 g kg⁻¹ of seed+AM fungi @ 5.0 g pot⁻¹) and T₈(50% RDF of Phosphorus +PSB @ 20g kg⁻¹ seed+AM fungi @ 5.0 g pot⁻¹) by 22.85%, 1.22%, 2.04%, 6.93%, 4.48%, 19.18% respectively. While the treatment T₂ have given significantly higher value by 21.90%, 5.71%, 17.95% over the applied treatment T₁, T₅ and T₈ respectively, while the values of treatment T₂, T₃, T₇ are at par.

The data pertaining k content (%) has been clearly observed that the potassium content percentage was maximum (1.16%) with the treatment T₄(T₁+PSB @ 20 g kg⁻¹ seed+AM fungi @ 5.0 g pot⁻¹), which is found to be significantly higher over the applied treatment T₁{(RDF (120:60:40)), T₂(T₁+PSB @ 20 g kg⁻¹ of seed),

T₅(75% RDF of Phosphorus +PSB @ 20 g kg⁻¹ seed), T₆(75% RDF OF Phosphorus +AM fungi @ 5.0 g pot⁻¹), T₇(75% RDF of Phosphorus +PSB @ 20 g kg⁻¹ of seed+AM fungi @ 5.0 g pot⁻¹) and T₈(50% RDF of Phosphorus +PSB @ 20g kg⁻¹ seed+AMfungi @ 5.0 g pot⁻¹) by 13.79%, 9.48%, 11.20%, 10.34%, 9.48%, 7.75% respectively while the value of T₄ and T₇ are at par. The treatment T₃ have given significantly higher value than 13.04%, 1.904% over the applied treatments T₁ and T₂, while the values of T₃ and T₄ are at par. The data (Fig. 1) pertaining nitrogen uptake (g pot⁻¹), has been clearly observed that the nitrogen uptake in shoot was maximum (0.0460%) with the treatment T₄ (T₁+PSB @ 20 g kg⁻¹ seed+AM fungi @ 5.0 g pot-1) of table in which the nitrogen uptake was maximum which is significantly higher than applied treatment T₁{(RDF (120:60:40)), T₂(T₁+PSB @ 20 g kg⁻¹ of seed), T₃(T₁+AM fungi @ 5.0 g pot⁻¹), T₅(75% RDF of Phosphorus +PSB @ 20 g kg⁻¹ seed), T₆(75% RDF Of Phosphorus +AM fungi @ 5.0 g pot⁻¹) T₇(75% RDF of Phosphorus +PSB @ 20 g kg⁻¹ of seed+AM fungi @ 5.0 g pot⁻¹) and T₈(50% RDF of Phosphorus +PSB @ 20 g kg⁻¹ seed+AM fungi @ 5.0 g pot⁻¹) by 21.73%, 10.86%, 17.39%, 13.47%, 13.26%, 17.39% and 14.78% respectively. 6.57%, 6.44%, 17.10%, 13.55% respectively. While the treatment T₃ have given significantly higher value 5.26% over the applied treatment T₁ while the value T₂, T₅ and T₈ are at par.

Table 1. Effect of Co-inoculation of PSB and AM fungi on microbial population under wheat rhizosphere

Treatment	Bacterial count (CFU 10 ⁶ g ⁻¹ soil)		Actinomycetes count (CFU 10 ⁶ g ⁻¹ soil)		Fungi count (CFU 10 ⁴ g ⁻¹ soil)	
	Flowering stage	Harvesting stage	Flowering stage	Harvesting stage	Flowering stage	Harvesting stage
T ₁	30.33	21.00	12.66	5.66	20.46	10.66
T ₂	37.33	28.66	20.67	8.67	14.66	8.67
T ₃	35.33	27.66	18.66	17.67	24.66	15.66
T ₄	39.00	29.33	21.33	11.67	20.67	9.67
T ₅	30.66	29.00	10.67	7.66	13.67	7.66
T ₆	31.66	31.00	9.67	8.66	21.66	13.67
T ₇	32.66	31.00	13.67	11.66	20.66	11.66
T ₈	29.33	20.00	8.66	6.67	22.66	11.67
C.D.(P=0.05)	1.671	1.51	1.671	1.008	1.008	1.008
C.V.	2.876	3.18	2.876	5.896	2.899	5.17

Note: T₁-RDF (120:60:40), T₂- T₁+PSB @ 20 g kg⁻¹ of seed, T₃-T₁+AM fungi @ 5.0 g pot⁻¹, T₄-T₁+PSB @ 20 g kg⁻¹ seed+AM fungi @ 5.0 g pot⁻¹, T₅-75% RDF of P +PSB @ 20 g kg⁻¹ seed, T₆-75% RDF of P +AM fungi @ 5.0 g pot⁻¹, T₇-75% RDF of P +PSB @ 20 g kg⁻¹ of seed + AM fungi @5.0 g pot⁻¹, and T₈-50% RDF of P +PSB @ 20 g kg⁻¹ seed + AM fungi @ 5.0 g pot⁻¹

Table 2. Effect of co-inoculation of PSB and AM fungi on availability of N P K (kg ha⁻¹) and organic carbon in soil after wheat harvest

Treatment	Available N, P K (kg ha ⁻¹ soil)			O C (%)
	Nitrogen	Phosphorus	Potassium	
T ₁	188.19	13.15	181.37	0.53
T ₂	199.95	16.29	185.52	0.54
T ₃	198.36	15.56	208.18	0.54
T ₄	210.24	19.25	210.07	0.54
T ₅	197.24	17.25	183.14	0.54
T ₆	204.18	18.16	184.33	0.54
T ₇	204.52	18.45	184.57	0.54
T ₈	202.13	14.25	200.14	0.54
C.D.(P=0.05)	0.422	0.209	0.232	NS
C.V.	0.12	0.722	0.069	-

Note: T₁-RDF (120:60:40), T₂- T₁+PSB @ 20 g kg⁻¹ of seed, T₃-T₁+AM fungi @ 5.0 g pot⁻¹, T₄-T₁+PSB @ 20 g kg⁻¹ seed+AM fungi @ 5.0 g pot⁻¹, T₅-75% RDF of P +PSB@ 20 g kg⁻¹ seed, T₆-75% RDF of P +AM fungi @ 5.0 g pot⁻¹, T₇-75% RDF of P +PSB @ 20 g kg⁻¹ of seed+AM fungi @ 5.0 g pot⁻¹, and T₈-50% RDF of P +PSB @ 20 g kg⁻¹ seed+AM fungi @ 5.0 g pot⁻¹.

Table 3. Effect of co-inoculation of PSB and AM fungi on N P K content (%) and N, P, K uptake (g pot⁻¹) by wheat plant

Treatment	N, P, K content (%) in plant		
	N	P	K
T ₁	0.140	0.189	1.00
T ₂	0.142	0.242	1.05
T ₃	0.145	0.240	1.15
T ₄	0.149	0.245	1.16
T ₅	0.141	0.228	1.03
T ₆	0.142	0.234	1.04
T ₇	0.144	0.235	1.05
T ₈	0.148	0.198	1.07
C.D.(P=0.05)	0.001	0.008	0.008
C.V.	0.402	0.936	0.426

Note: T₁-RDF (120:60:40), T₂- T₁+PSB @ 20 g kg⁻¹ of seed, T₃-T₁+AM fungi @ 5.0 g pot⁻¹, T₄-T₁+PSB @ 20 g kg⁻¹ seed+AM fungi @ 5.0 g pot⁻¹, T₅-75% RDF of P +PSB@ 20 g kg⁻¹ seed, T₆-75% RDF of P +AM fungi @ 5.0 g pot⁻¹, T₇-75% RDF of P +PSB @ 20 g kg⁻¹ of seed+AM fungi @ 5.0 g pot⁻¹, and T₈-50% RDF of P +PSB @ 20 g kg⁻¹ seed+AM fungi @ 5.0 g pot⁻¹.

Also from the data (Fig. 1) pertaining P uptake (g pot⁻¹) clearly observed that the phosphorus uptake in shoot was maximum (0.0760 g pot⁻¹) with the treatment T₄ (T₁+PSB @ 20 g kg⁻¹ seed+AM fungi @ 5.0 g pot⁻¹) of table in which the phosphorus uptake was maximum which is significantly higher than applied treatment T₁{(RDF (120:60:40)), T₂(T₁+PSB @ 20 g kg⁻¹ of seed), T₃(T₁+AM fungi @ 5.0 g pot⁻¹), T₅(75% RDF of Phosphorus +PSB @ 20 g kg⁻¹ seed), T₆(75% RDF of Phosphorus +AM fungi @ 5.0 g pot⁻¹) by 35.52%, 6.57%, 6.44%, 17.10%, 13.55%, 17.63%, 29.73% respectively. The similar results reported by [15].

The data (Fig 1) pertaining K uptake (g pot⁻¹) has been clearly observed that the potassium uptake in shoot was maximum (0.362%) with the

application of treatment T₄ (T₁+PSB @ 20 g kg⁻¹ seed+AM fungi @ 5.0 g pot⁻¹) of table in which the potassium uptake was maximum which is significantly higher than applied treatment T₁{(RDF (120:60:40)), T₂(T₁+PSB @ 20 g kg⁻¹ of seed), T₃(T₁+AM fungi @ 5.0 g pot⁻¹), T₅(75% RDF of Phosphorus +PSB @ 20 g kg⁻¹ seed), T₆(75% RDF OF Phosphorous +AM fungi @ 5.0 g/pot) by 27.07% and 14.91%, 6.077%, 21.27%, 19.33%, respectively. While the treatment T₃ have given significantly higher value by 22.35%, 9.41%, 16.17%, 14.11%, 17.94%, 15% over the applied treatment T₁, T₂, T₅, T₆, T₇ and T₈ respectively with the value T₄ is at par. It clearly indicate that the potassium uptake was more and more occur in inoculated species with respect to uninoculated treatment. It might be due to the co-inoculation of PSB and AMF along




Fig. 1. Effect of co-inoculation of PSB and AM fungi on NPK uptake by wheat plant

Note: T1-RDF (120:60:40), T2- T1+PSB @ 20 g kg⁻¹ of seed, T3-T1+AM fungi @ 5.0 g pot⁻¹, T4-T1+PSB @ 20 g kg⁻¹ seed+AM fungi @ 5.0 g pot⁻¹, T5-75% RDF of P +PSB @ 20 g kg⁻¹ seed, T6-75% RDF of P +AM fungi @ 5.0 g pot⁻¹, T7-75% RDF of P +PSB @ 20 g kg⁻¹ of seed

with full dose of fertilizers, PSB secrete organic acids of lower molecular weight and AM fungi make it more available to plants through its hyphae and thus nutrient content and uptake concentration increases. Similar results were obtained by Sandhya et al. [17], who conducted an experiment on Co-inoculation studies of vesicular Arbuscular Mycorrhizal fungi (VAM) and Phosphate solubilizing bacteria (PSB) on nutrient uptake of *Marsdenia volubilis* (T. Cooke) and found an excellent improvements in uptake of nutrients like N, P, K, Ca, Mg, Fe, Mn and Zn concentrations in *Marsdenia volubilis* than single application.

4. CONCLUSION

The results of the study revealed that the maximum microbial population viz., Bacteria, Actinomycetes, Fungi were found co-inoculation of PSB and AM fungi along with recommended dose of phosphorus. The Co-inoculation of phosphate solubilizing bacteria and arbuscular mycorrhizal fungi along with 100% RDF gave significantly higher available nitrogen, phosphorus and potassium in soil as well as their uptake by the wheat plant.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Marschner M. Mineral nutrition of higher plants. Plant and Soil. 1996;159:89-102.
- Gyaneshwar P, Kumar N, Parekh LJ, Poole PS. Role of soil microorganisms in improving P nutrition of plants. Plant and Soil. 2002;245:83-93.
- Sakurai K, Tanaka N, Iwasaki K, Tanaka S. Effect of arbuscular mycorrhizal application on the distribution Azcon-Aguilar C, Barea JM. Applying mycorrhizal biotechnology to horticulture; 2001.
- Thom C, Raper KB. A manual of the *Aspergilli*. Williams and Wilkins Co. Baltimore, U. S. A; 1945.
- Subba RNS. *Rhizobium* and root nodulation. In: Soil microorganisms and plant growth. Oxford IBH New Delhi; 1986.
- Jackson ML. Soil chemical analysis prentice-hall inc. Englewood Cliffs, New Jersey; 1973.
- Hedley M, Stewart J. Method to measure microbial phosphate in soils. Soil Biol. Biochem. 1982;14:377-385.
- Gomez KA, Gomez AA. Statistical procedures in agricultural research, by New York, Chichester, etc. Wiley 2nd edition, paperback. 1984;680.
- Marschner P, Crowley DE, Higashi M. Root exudation and physiological status of

- a root colonizing fluorescent pseudomonas in mycorrhizal and non-mycorrhizal pepper (*Capsicum annuum*). Plant Soil. 1997;189:11-20.
10. Hayman DS. The physiology of vesicular-arbuscular mycorrhizal symbiosis. Canadian Journal of Botany. 1983;61:944-963.
 11. Singh M, Kumar N, Kumar S, Mehi lal M. Effect of co-inoculation of *B. Japonicum*, psb and am fungi on microbial biomass carbon, nutrient uptake and yield of soybean (*Glycine max* L. merril). Agriways. 2015;3(1):14-18. [ISSN: 2321-8614]
 12. Zhu HH, Long LK, Yang SZ. Influence of AM fungus on *Ralstonia Solanacearum* Population and bacterial community structure in rhizospheren mycosystema. 2005;24(11):137-142.
 13. Sah AK, Singh M, Prasad S, Bahera KS, Chattopadhaya N, Kiran K. Effect of organic fertilizer and microbial inoculants on soil biological properties and yield of scented rice (*Oryza sativa* L.). Journal of Applied and Natural Science. 2018;10(3): 995–1002.
 14. Rovira AD. Interaction between plant roots and soil microorganisms. Ann. Rev. Microbiol. 1965;19:241-246.
 15. Singh M, Kumar N. Effect of FYM, Vermicompost, Vermiwash and NPK on Growth, Microbial Biomass C and Yield of Soybean. Soybean Res. 2012;10:60-66.
 16. Salem M, Al-Amri, Khalid M, Elhindi Ahmed F, Sharaf El-Din. Effect of arbuscular mycorrhizal fungus *Glomus mosseae* and phosphorus application on plant growth rate, essential oil content and composition of coriander (*Coriander sativum* L.). Progress in Nutrition. 2016; 10(4):443-454.
 17. Sandhya A, Vijaya T, Sridevi A, Narasimha G. Annals of plant sciences co-inoculation studies of vesicular arbuscular mycorrhizal fungi (VAM) and phosphate solubilizing bacteria (PSB) on nutrient uptake of *Marsdenia volubilis* (T. Cooke); 2014.

© 2019 Rani et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<http://www.sdiarticle3.com/review-history/46747>