



Effect of Methanolic Extract of *Ocimum gratissimum* on Blood Pressure, Some Electrolytes, Renal and Cardiac Biomarkers in 8% NaCl Induced Hypertensive Male Wistar Rats

**O. B. Onyema-Iloh^{1*}, S. C. Meludu², E. O. Iloh³, C. E. Dioka⁴, O. S. Usman⁴,
O. A. Onyegbule⁴ and E. B. Olugbenga⁵**

¹Department of Chemical Pathology, Nnamdi Azikiwe University Teaching Hospital, Nnewi, Nigeria.

²Department of Basic Medical Sciences, Nnamdi Azikiwe University, Awka (Nnewi Campus), Nigeria.

³Department of Industrial Chemistry, Chukwuemeka Odumegwu Ojukwu University, Uli Campus, Nigeria.

⁴Department of Chemical Pathology, Nnamdi Azikiwe University, Awka (Nnewi Campus), Nigeria.

⁵Department of Chemical Pathology, University of Port Harcourt, River State, Nigeria.

Authors' contributions

This work was carried out in collaboration between all authors. Author OBOI wrote the protocol and carried out sample analysis. Author EOI carried out phytochemical and mineral analysis. Authors SCM and CED assisted in study design and protocol. Author OSU managed the statistical analysis. Authors OAO and EBO managed the literature searches. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/EJMP/2018/45379

Editor(s):

- (1) Dr. Patrizia Diana, Professor, Department of Molecular and Biomolecular Sciences and Technologies, University of Palermo, Palermo, Italy.
(2) Dr. Shanfa Lu, Professor, Institute of Medicinal Plant Development, Chinese Academy of Medical Sciences & Peking Union Medical College, China.
(3) Dr. Marcello Iriti, Professor, Plant Biology and Pathology, Department of Agricultural and Environmental Sciences, Milan State University, Italy.

Reviewers:

- (1) Ugwu Melvin Nnaemeka, Cross River University of Technology, Okuku Campus, Nigeria.
(2) Sharafudeen D. Abubakar, College of Health Sciences, Katsina, Nigeria.
(3) Md Afjalus Siraj, University of Hawaii at Hilo, USA.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/45379>

Original Research Article

Received 14 October 2018
Accepted 27 December 2018
Published 21 January 2019

ABSTRACT

Background: *Ocimum gratissimum* (OG) is a shrub belonging to the family of Lamiaceae. It is commonly called scent leaf or clove basil and it is found in many tropical countries. Studies have shown that the leaf extract of *Ocimum gratissimum* possess medicinal properties.

*Corresponding author: E-mail: onyemailohoby@yahoo.com;

Aim: The effect of methanolic extract of *Ocimum gratissimum* on blood pressure, electrolytes, renal and cardiac biomarkers in 8% NaCl- induced hypertensive male Wistar rats.

Methodology: Forty Wistar rats (120-160) g were assigned to 5 groups of eight rats each. Group 1, 2, 3, 4 and 5 constitute the normal, hypertensive group, OG (200 mg/kg bwt) group, OG (400 mg/kg bwt) group and reference drugs (lisinopril, 30 mg/kg) group respectively. Group 3, 4 and 5 were given the extract and reference drug through oral gavage. All groups except group 1 were induced with 8% NaCl from 0-4weeks before treatment with OG and reference drug from 5-8 weeks. Electrolytes and other biochemical parameters were assayed using standard methods.

Results: The phytochemical results revealed the presence of phenol, flavonoids, alkaloids, phytate, tannis and saponin. At 4 weeks (after induction), systolic blood pressure (SBP), diastolic blood pressure (DBP), serum sodium, chloride, urea, and creatinine significantly ($p < 0.05$) increased while serum potassium significantly ($p < 0.05$) decreased in all the groups except group 1. At 8 weeks, after treatment with OG (200 mg/kgbwt), OG (400 mg/kgbwt) and lisinopril (30 mg/kg), SBP, DBP, serum sodium, chloride, urea, and creatinine significantly ($p < 0.05$) decreased while serum potassium significantly ($p < 0.05$) increased. Creatine kinase (CK) and CK-MB however, were not significantly altered after the 4th and 8th week.

Conclusion: OG extract possesses an antihypertensive effect and enhances the proper functioning of the kidney. It may also be useful in hypertensive condition due to its nephroprotective effect at 200mg/kgbwt and 400 mg/kgbwt.

Keywords: *Ocimum gratissimum* (OG); phytochemical analysis; blood pressure; electrolytes; renal markers; cardiac markers; NaCl hypertension.

1. INTRODUCTION

Ocimum gratissimum (OG) is an edible plant belonging to the family of Lamiaceae. It is commonly called scent leaf or clove basil and it is found in many tropical countries. The genus, *Ocimum* (Lamiaceae), comprises 65 aromatic species, distributed in tropical and subtropical regions of the world including Africa and Asia [1]. Various *Ocimum* species, including *Ocimum basilicum* and *Ocimum gratissimum*, have important culinary and pharmacological uses. Their culinary and pharmacological values are attributed mainly to their aromatic compounds [2]. Hence, the leaves are used as a fragrance and flavoring agent in a variety of products including food, beverages, condiments, and oral care products [3]. *Ocimum gratissimum* leaves extract possesses anti-arthritis [4], hypoglycemic [1], anticonvulsant and anxiolytic activities [5]. The leaf extract of *Ocimum gratissimum* showed anti-diabetic properties [6,7] and antibacterial activities [8,9]. It has mosquito repellent and mosquitocidal potential [10], as well as hepatoprotective effect [11,12]. Hypertension is a serious public health problem due to increasing incidence and prevalence. It is a significant risk factor for heart disease, stroke and other cardiovascular diseases [13]. Although antihypertensive drug are used in the management of hypertension, some of these drugs are faced with the problem of fake and

adulterated drugs, drug abuse, high cost of drugs and drug side effects such as erectile dysfunction, extra urination, weakness and asthma symptoms. There is paucity of literature on the use of this plant as antihypertensive. Also, natural plant products are increasingly investigated for their therapeutic potentials. Hence, the need to look at our indigenous plant *Ocimum gratissimum* which are natural, safe, readily available at almost no cost when compared with conventional drugs for their possible nutraceutical benefit and therapeutic properties that may aid in the prevention and management of hypertension when taken as a supplement.

The aim of the study is to determine the antihypertensive effect of *Ocimum gratissimum* that may possibly be used in the management and prevention of hypertension.

- 1) To determine the phytochemical and mineral contents of methanolic leaf extract of *Ocimum gratissimum*(OG).
- 2) To determine the effect of OG on blood pressure, electrolytes, urea, creatinine, Creatine kinase (CK) and Creatine kinase-MB in 8% NaCl induced hypertensive male Wistar rats in order to ascertain its antihypertensive effect when compared with the reference drug (lisinopril).

2. MATERIALS AND METHODS

2.1 Plant Extraction (Maceration Method) [14]

Fresh leaves of *Ocimum gratissimum* (scent leaf) were air-dried at room temperature. Air-dried leaves of the plant were milled into powder. The powdered leaves were weighed and macerated into methanol (500 g of the plant material to 2.5 liters of methanol in a stopper) for 5 days with occasional shaking to soften and break the plant cell wall to ensure sufficient extraction of the active photochemical. At the end of five days, the methanolic extract was filtered using whatman No.1 filter paper and the filtrate concentrated to dryness under reduced pressure at $60 \pm 1^\circ\text{C}$ in a rotary evaporator at 45°C , weighed and stored frozen until used. The exact weight of dried extract from 500g powder was 87.8g and gave a percentage yield of 17.56%. The extract was dissolved in tween 80 and was given to the animals at graded doses of 200mg/kgbw and 400 mg/kgbw.

2.2 Procurement and Care of Animals

Forty male Wistar rats weighing 120-160 g were obtained from university of Nigeria Nsukka and housed in cages in the animal facility. They were fed with clean water and rat chow *ad libitum*. The rats were allowed to acclimatize for 2 weeks during which the rat local restrainer was included in their cages to prepare the rat for blood pressure measurement. The rats were maintained under good laboratory conditions at a temperature of $22 \pm 2^\circ\text{C}$, relative humidity of $50 \pm 5\%$ and photoperiod of 12 hr (12 h-dark and 12h-light cycle).

2.3 Experimental Design

The animal study protocol was designed into two stages. The first stage is the induction period while the second stage is the treatment period.

The animals were grouped into five different groups with each group assigned different numbers (1 – 5). Each group (Group 1, 2, 3, 4 and 5) contains 8 rats and were fed with normal rat chow and clean water *ad libitum* for the period of the study.

Group 1 (negative control) - Normal rat chow and water *ad libitum*

Group 2 (Untreated hypertensive) -Induction with 8% NaCl for 4weeks and subsequently no treatment

Group 3 (200 mg/kgbw OG) -Induction with 8% NaCl for 4 weeks and subsequent treatment with 200 mg/kgbw OG for another 4 weeks.

Group 4(400 mg/kgbw OG) -Induction with 8% NaCl for 4weeks and subsequent treatment with 400 mg/kgbw OG for another 4 weeks

Group 5(30 mg/kg Lisinopril) -Induction with 8% NaCl for 4 weeks and subsequent treatment with lisinopril for another 4 weeks.

2.4 Blood Pressure Measurement in Rats Using Kent Scientific CODA Machine

Volume pressure recording (VPR) method of blood pressure measurement was used as suggested by the American Heart Association [15]. Blood pressure was measured using VPR non-invasive blood pressure monitoring system (CODA-6) Kent Scientific, Torrington, CT).

Blood pressure measurement procedure: The method of measurement of VPR is based on tail volume. At the start of the measurement cycle, blood is pushed from the tail by the VPR cuff and then the occlusion cuff inflates to prevent blood flow back into the tail. When the occlusion cuff deflates, blood begins to flow back into the tail, increasing the tail volume. The occlusion cuff pressure at which the tail volume increases is the SBP. The tail volume will continue to increase as the occlusion cuff deflates until blood flow into and out of the tail equalizes; the occlusion cuff pressure at this point is the DBP.

2.5 LD 50: Acute Toxicity Studies of Methanolic Extract of *Ocimum gratissimum* (MEOG)

The acute toxicity of MEOG was determined by Lorke's method [16] as described by Enevide [17]. Rats were divided into two phases. In the first phase of the study, 9 rats were divided into 3 groups of 3 rats each and they were treated with MEOG through oral gavage at the doses of 10, 100 and 1000 mg/kg respectively. The rats were observed for behavioural change and lethality in the first 24 hrs.

In the second stage, 3 rats were divided into 3 groups of 1 rat each and they were treated with MEOG through oral gavage at the doses of 1600,

2900 and 5000 mg/kg. The general behaviour of the animals were observed continuously for 1 hr after treatment and then intermittently for 4 hrs, then hourly for the next 24 hrs. The LD₅₀ was determined using the formular below

$$LD_{50} = \sqrt{a \times b}$$

Where a = minimal lethal dose
b = maximal survival dose

$$OG LD_{50} = \sqrt{a \times b} = \sqrt{1600 \times 5000} \Rightarrow \sqrt{8000000} = 2828$$

2.6 Biochemical methods/assay

Determination of Alkaloids: Alkaloids were determined using the method of Harborne [18].

Procedure:

Five grams (5 g) of the sample was weighed into a 250 ml beaker and 200 ml of 20% acetic acid in ethanol was added and covered and allowed to stand for 4 hours at 25°C. This was filtered with filter paper No. 42 and the filtrate was concentrated using a water bath (Memmert) to one quarter of the original volume. Concentrated ammonium hydroxide was added drop wise to the extract until the precipitate was complete. The whole solution was allowed to settle and the precipitate was collected and washed with dilute NH₄OH (1% ammonia solution). Then, filter with pre- weighed filter paper. The residue on the filter paper is the alkaloid, which is dried in the oven (precision electrothermal model BNP 9052 England) at 80°C. The alkaloid content was calculated and expressed as a percentage of the weight of the sample analyzed.

Calculation:

%weight of alkaloid = (weight of filter paper with residue - weight of filter paper)/ Weight of sample analyzed x 100

Determination of Flavonoids: Flavonoids were determined using the method of Bohm and Kocipai-Abyazan [19].

Procedure:

10 g of the plant sample was extracted repeatedly with 100 ml of 80% aqueous methanol at room temperature. The whole solution was filtered through whatman filter paper No. 42 (125 mm).

The filtrate was later transferred into a crucible and evaporated into dryness over a waterbath and weighed to a constant weight.

Calculation:

%flavonoids = {(weight of crucible + residue) - (weight of crucible)}/ Weight of sample analyzed x 100

Determination of Saponin: Saponin was determined using the method of Obadoni and Ochuko [20].

Procedure:

20 g of each grounded sample was put into a conical flask and 100 cm³ of 20% aqueous ethanol was added. Then the flask was heated on a hot water bath for 4 hrs with constant stirring at about 55°C. The mixture was then filtered and the residue was again extracted with another 200 ml 20% ethanol. The combined extract was reduced to 40 ml on a hot water bath at about 90°C. The concentrate was transferred into a 250 ml separatory funnel, added 20 ml diethyl ether in it followed by vigorous shaking. The aqueous layer was recovered while the ether layer was discarded. The purification process was repeated. 60 ml of n-butanol was added. The combined n-butanol extracts were washed twice with 10 ml of 5% aqueous sodium chloride. The remaining solution was heated in a water bath. After evaporation the samples were dried in oven, weighed and saponin content was calculated as percentage.

Calculation:

% saponin content = {(weight of filter paper + residue) - (weight of filter paper)}/ Weight of sample analyzed x 100

Determination of Tannin: Tannin content of the sample was determined by Follins Dennis titration method as described by Pearson [21].

Procedure:

The follins Dennis titrating method as described by Pearson (1974) was used. To 20 g of the crushed sample in a conical flask was added 100 ml of petroleum ether and covered for 24 hours. The sample was then filtered and allowed to stand for 15 minutes allowing petroleum ether to evaporate. It was then re-extracted by soaking in 100ml of 10% acetic acid in ethanol for 4hrs. The

sample was then filtered and the filtrate collected. Then 25 ml of NH₄OH were added to the filtrate to precipitate the alkaloids. The alkaloids were heated with electric hot plate to remove some of the NH₄OH still in solution. The remaining volume was measured to be 33 ml. 5 ml of this was taken and 20 ml of ethanol was added to it. It was titrated with 0.1 M NaOH using phenolphthalein indicator until a pink end point is reached.

Calculation:

Tannin content was then calculated in % ($C_1V_1 = C_2V_2$) molarity.

Where:

C_1 = conc. of Tannic Acid

C_2 = conc. Of Base

V_1 = Volume of Tannic acid

V_2 = Volume of Base

$$\text{Therefore } C_1 = \frac{C_2 V_2}{V_1}$$

% of tannic acid content =

$$\frac{C_1 \times 100}{\text{Weight of sample analyzed}}$$

Determination of Total Phenols: Total phenols were determined by spectrophotometric method as described by Khan et al. [22].

Procedure:

Two grams (2 g) of each plant sample was defatted with the help of 100 ml of diethyl ether using a soxhlet apparatus for 2 h. The fat free sample was boiled with 50 ml of ether for 15 min for the extraction of phenolic component. 5 ml of the extract was pipetted into a 50 ml flask and 10 ml distilled water was added. 2 ml of ammonium hydroxide solution and 5 ml of concentrated amylalcohol were also added in it. The samples were made up to mark and left to react for 30 min. Colour was developed and its absorbance was measured at 505 nm.

Determination of Phytate Contents: Phytate contents were determined using the method of Lucas and Markakes [23].

Procedure:

0.2g of each samples was weighed into different 250ml conical flasks. Each sample was soaked

in 100 ml of 2% concentrated HCl for 3 hours. The samples were then filtered. 50 ml of each filtrate was placed in 250ml beaker and 100ml distilled water added to each sample. 10 ml of 0.3% ammonium thiocyanate solution was added as indicator and titrated with standard iron (III) chloride solution which contained 0.00195g iron per l ml.

Calculation:

$$\text{Phytic acid} = \frac{\text{Titre value} \times 0.00195 \times 1.19 \times 100}{\text{Wt of sample}}$$

Determination of mineral content of *Ocimum gratissimum*: Mineral content was determined

by Association of Official Analytical Chemists methods [24] using the flame system of the atomic absorption spectrophotometry (AAS), (Varian Spectr AA240, USA).

Procedure:

Ocimum gratissimum was ashed at 550°C overnight and the ash was dissolved in concentrated nitric acid and filtered, diluted to 50 ml with deionized water and the absorbance of the samples was read directly on the AAS. Working standard solutions of potassium, calcium, magnesium, phosphorus and sodium were prepared from stock standard solution (1000 ppm), in 2 N HNO₃ and absorbance was noted for standard solution of each element and samples using atomic absorption spectrophotometer (AAS). Calibration curve for each metal was prepared by plotting the absorbance of standards versus their concentrations. A blank reading was also taken and necessary corrections were made during the calculation of concentration of various elements.

Determination of urea level: Blood urea was determined by the method described by Kassirer [25].

Urea is decomposed by urease to form ammonia and carbon dioxide. Ammonia combines with 2-oxo-glutarate in presence of glutamate dehydrogenase and NADH to form L-Glutamate and NAD. The rate of NAD formation measured at 340 nm is directly proportional to the amount of blood urea.

Procedure:

Test tubes were labelled: reagent blank, Standard, Control and Sample. 1.0 ml of enzyme

reagent (urease, ADP), followed by 0.01 ml of sample/standard/control in their respective tubes and mixed thoroughly. These solutions were allowed to stand at 37°C for one minute. 0.250 ml of 2-oxoglutarate and NADH reagent was added to all the tubes and mixed. The initial absorbance was measured after 30 seconds (A1) and the second reading taken at exactly one minute (A2) measured against the reagent blank at 340 nm.

Calculation:


The mean change in absorbance readings were calculated thus ($\Delta A2-A1/min$)

Urea (mmol/l) = ($\Delta A_{sample}/\Delta A_{std}$) X Concentration of standard

3.5.13 Determination of Creatinine Level

Serum creatinine was estimated by Jaffe's method as described by Laron [26].

Serum creatinine in alkaline medium reacts with Picric acid to produce orange colour that absorbs light at 492 nm. The rate of increasing absorption is directly proportional to the amount of creatinine in the sample.





Procedure:

Test tubes were labelled: Blank, Standard, Control, and Sample. One (1.0) ml of reagent was added into a test tube and allowed to equilibrate at 37°C. The spectrophotometer was zeroed with the reagent blank at 510 nm. 0.05 ml of sample or standard or control was added to the working reagent, mixed and incubated at 37°C. The first reading was taken after 30 seconds and the absorbance recorded as A1. At exactly sixty seconds after the A1 reading, the solution was read again and the absorbance recorded as A2. The change in absorbance ($\Delta A/min$) was calculated by subtracting (A2 - A1).

Calculation:

Creatinine ($\mu\text{mol/l}$) = ($\Delta\text{Abs. of unknown}/ \Delta\text{Abs. of standard}$) x Concentration of standard

Determination of CK Activity: CK activity was assayed using the methods of Szasz [27] as recommended by the IFCC.



The rate of NADPH formation is directly proportional to the catalytic CK activity. It is determined by measuring the increase in absorbance at 340 nm.

Creatine kinase-MB determination:

Immunoinhibition method was used in estimation of CK-MB [28]. The sample was incubated in the CK-MB reagent which includes the CK-M antibody. The activity of the non-inhibited CK-B was then determined using the following series of reaction. CK-B catalyses the reversible phosphorylation of ADP, in the presence of creatine phosphate to form ATP and creatinine. The auxiliary enzyme hexokinase (HK) catalyses the phosphorylation of glucose by ATP format, to produce ADP and glucose-6-phosphate is oxidized to 6-phosphogluconate with concomitant production of NADH. The rate of NADH formation, measured at 340 nm, is directly proportional to serum CK-B activity.

2.7 Statistical Analysis

The version 23 of Statistical Package for Social Sciences (SPSS) was used in statistical analysis. The variables were expressed as mean±SD. The independent student t-test was used to assess significant mean difference between two independent groups, while paired t-test was used to assess the mean difference between two related groups. Analysis of Variance (ANOVA) was also used and POST HOC was used to determine the significant difference within the groups. The level of significance was considered at P<0.05.

3. RESULTS

Phytochemical result revealed the presence of phenol, steroid, alkaloids, flavonoids, saponin, tannis and phytate while the mineral content revealed the presence of potassium, phosphorus, calcium, magnesium and sodium (Table 1).

Systolic blood pressure, diastolic blood pressure, serum sodium, chloride, urea and creatinine were significantly increased (p<0.05) while

serum potassium and bicarbonate were significantly decreased ($p < 0.05$) after 4 weeks induction of hypertension. After administration of methanolic extract of *Vernonia amygdalina* and *Ocimum gratissimum* at 200 mg/kgbw and 400mg/kgbw systolic blood pressure, diastolic blood pressure, serum sodium, chloride, urea and creatinine were significantly decreased ($p < 0.05$) while serum potassium significantly increased ($p < 0.05$) after 4 weeks induction of hypertension in a dose-dependent manner (Tables 2-9). Serum creatine and creatine-kinase did not differ significantly at 4 weeks and 8 weeks respectively (Tables 10-11).

Table 1. Phytochemical and mineral component of methanolic extract of *Ocimum gratissimum* (OG)

| Phytochemical (mg/100 g) | <i>Ocimum gratissimum</i> (OG) |
|---------------------------------|--------------------------------|
| Phenol | 0.52 |
| Steroid | 0.24 |
| Alkaloids | 1.04 |
| Flavonoids | 3.02 |
| Saponins | 3.35 |
| Tannis | 2.16 |
| Phytate | 4.24 |
| Mineral components (ppm) | |
| Potassium | 42.60 |
| Calcium | 32.60 |
| Magnesium | 60.20 |
| Phosphorus | 12.35 |
| Sodium | 15.80 |

Table 2. Effect of methanolic extract of *Ocimum gratissimum* (OG) on systolic blood pressure (SBP) at 0 week (baseline), 4 weeks (after induction of hypertension) and 8 weeks (after treatment with OG)

| SBP (n=8) | 0 week | 4 weeks | 8 weeks |
|------------------------|-------------|---------------------------|------------------------------|
| Neg Control | 81.00±11.23 | 82.66±23.71 | 81.00±11.21 |
| Untreated group | 83.50±10.32 | 162.33±6.12 ^a | 190.17±5.49 ^{b,*} |
| OG 200mg/kg bwt | 84.83±11.35 | 178.16±13.15 ^a | 123.83±3.48 ^{b,c,*} |
| OG 400mg/kg bwt | 82.66±11.20 | 174.17±13.39 ^a | 92.66±6.94 ^{b,c,*} |
| Lisinopril 30mg/kg bwt | 82.23±19.66 | 168.66±6.88 ^a | 84.66±4.63 ^b |

Values expressed as mean± S.D. a= Statistically significant at $P < 0.05$ when compared between 0 week and 4 weeks, b= when compared between 4 weeks and 8 weeks and c= when compared between 0 week and 8 weeks and * when compared within the groups and control

Table 3. Effect of methanolic extract of *Ocimum gratissimum* (OG) on Diastolic blood pressure (DBP) at 0 week (baseline), 4 weeks (after induction of hypertension) and 8 weeks (after treatment with OG)

| DBP (n=8) | 0 week | 4 weeks | 8 weeks |
|------------------------|-------------|---------------------------|-------------------------------|
| Neg Control | 60.66±11.91 | 60.67±11.91 | 60.67±11.91 |
| Untreated group | 61.50±8.57 | 124.66±16.35 ^a | 132.83±10.92 ^{b,c,*} |
| OG 200mg/kg bwt | 61.16±8.70 | 127.66±13.4 ^a | 74.00±8.62 ^{b,*} |
| OG 400mg/kg bwt | 63.83±5.60 | 126.67±13.78 ^a | 63.00±8.83 ^b |
| Lisinopril 30mg/kg bwt | 63.80±9.57 | 126.66±13.41 ^a | 74.66±4.63 ^{b,*} |

Values expressed as mean± S.D. a= Statistically significant at $P < 0.05$ when compared between 0 week and 4 weeks, b= when compared between 4 weeks and 8 weeks and c= when compared between 0 week and 8 weeks and * when compared within the groups and control

Table 4. Effect of methanolic extract of *Ocimum gratissimum* (OG) on serum sodium (Na) level at 0 week (baseline), 4 weeks (after induction of hypertension) and 8 weeks (after treatment with OG)

| Na (n=8) | 0 week | 4 weeks | 8 weeks |
|-------------------------|-------------|---------------------------|------------------------------|
| Neg Control | 136.67±1.36 | 136.62±1.32 | 136.67±1.36 |
| Untreated group | 138.16±1.94 | 168.50±3.45 ^a | 174.83±6.49 ^{b,c,*} |
| OG 200 mg/kg bwt | 136.67±1.36 | 182.33±4.68 ^a | 155.67±8.56 ^{b,c,*} |
| OG 400 mg/kg bwt | 138.17±1.94 | 180.83±17.51 ^a | 143.33±8.21 ^{b,c,*} |
| Lisinopril 30 mg/kg bwt | 138.67±1.50 | 181.10±7.51 ^a | 149.50±7.28 ^{b,c,*} |

Values expressed as mean± S.D. a= Statistically significant at P<0.05 when compared between 0 week and 4 weeks, b= when compared between 4 weeks and 8 weeks and c= when compared between 0 week and 8 weeks and * when compared within the groups and control

Table 5. Effect of methanolic extract of *Ocimum gratissimum* (OG) on serum Potassium (K) level at 0 week (baseline), 4 weeks (after induction of hypertension) and 8 weeks (after treatment with OG)

| K (n=8) | 0 week | 4 weeks | 8 weeks |
|-------------------------|-----------|------------------------|--------------------------|
| Neg control | 5.53±0.16 | 5.50±0.12 | 5.53±0.16 |
| Untreated group | 5.53±0.16 | 3.40±0.36 ^a | 3.20±0.32 ^{c,*} |
| OG 200 mg/kg bwt | 5.43±0.07 | 4.98±0.63 ^a | 5.78±0.29 ^b |
| OG 400 mg/kg bwt | 5.45±0.09 | 5.00±0.95 ^a | 5.72±0.34 ^b |
| Lisinopril 30 mg/kg bwt | 5.47±0.08 | 4.83±0.84 ^a | 4.82±0.84 ^{c,*} |

Values expressed as mean± S.D. a= statistically significant at P<0.05 when compared between 0 week and 4 weeks, b= when compared between 4 weeks and 8 weeks and c= when compared between 0 week and 8 weeks and * when compared within the groups and control

Table 6. Effect of methanolic extract of *Ocimum gratissimum* (OG) on serum chloride (Cl) level at 0 week (baseline), 4 weeks (after induction of hypertension) and 8 weeks (after treatment with OG)

| Cl(mmol) n=8 | 0 week | 4 weeks | 8 weeks |
|-------------------------|---------------|----------------------------|--------------------------------|
| Neg control | 105.33 ± 2.41 | 105.31 ± 2.42 | 105.32 ± 2.42 |
| Untreated group | 105.83 ± 0.75 | 125.00 ± 9.65 ^a | 130.33 ± 3.50 ^{b,c,*} |
| OG 200 mg/kg bwt | 105.83 ± 1.17 | 121.33 ± 3.72 ^a | 104.83 ± 3.76 ^b |
| OG 400 mg/kg bwt | 105.33 ± 2.42 | 123.00 ± 4.24 ^a | 105.50 ± 3.94 ^b |
| Lisinopril 30 mg/kg bwt | 103.67 ± 2.58 | 128.00 ± 3.63 ^a | 128.00 ± 3.63 ^{c,*} |

Values expressed as mean± S.D. a= Statistically significant at P<0.05 when compared between 0 week and 4 weeks, b= when compared between 4 weeks and 8 weeks and c= when compared between 0 week and 8 weeks and * when compared within the groups and control

Table 7. Effect of methanolic extract of *Ocimum gratissimum* (OG) on serum bicarbonate (HCO₃) level at 0 week (baseline), 4 weeks (after induction of hypertension) and 8 weeks (after treatment with OG)

| HCO ₃ (mmol/l) n=8 | 0week | 4 weeks | 8 weeks |
|-------------------------------|--------------|---------------------------|-----------------------------|
| Neg control (G1) | 25.16 ± 1.72 | 25.13 ± 1.82 | 25.17 ± 1.78 |
| Untreated group (G2) | 26.50 ± 1.64 | 24.50 ± 0.54 ^a | 23.17 ± 1.69 ^c |
| OG 200 mg/kg bwt(G5) | 25.50 ± 1.64 | 17.17 ± 2.23 ^a | 17.17 ± 2.24 ^{c,*} |
| OG 400 mg/kg bwt(G6) | 25.16 ± 1.72 | 17.16 ± 3.31 ^a | 16.63 ± 2.73 ^{c,*} |
| Lisinopril 30 mg/kg bwt(G7) | 26.00 ± 1.41 | 20.33 ± 1.86 ^a | 20.33 ± 1.86 ^{c,*} |

Values expressed as mean± S.D. a= Statistically significant at P<0.05 when compared between 0 week and 4 weeks, b= when compared between 4 weeks and 8 weeks and c= when compared between 0 week and 8 weeks and * when compared within the groups and control

Table 8. Effect of methanolic extract of *Ocimum gratissimum* (OG) on serum urea level at 0 week (baseline), 4 weeks (after induction of hypertension) and 8weeks (after treatment with OG)

| Urea(mmol/l) n=8 | 0 week | 4 weeks | 8 weeks |
|-----------------------------|-------------|--------------------------|----------------------------|
| Neg control (G1) | 5.63 ± 0.03 | 5.61 ± 0.01 | 5.62 ± 0.02 |
| Untreated group (G2) | 5.62 ± 0.02 | 7.33 ± 0.67 ^a | 7.82 ± 0.46 ^{c,*} |
| OG 200 mg/kg bwt(G5) | 5.62 ± 0.01 | 6.86 ± 0.52 ^a | 5.78 ± 0.48 ^b |
| OG 400 mg/kg bwt(G6) | 5.63 ± 0.01 | 7.11 ± 0.41 ^a | 5.68 ± 0.33 ^b |
| Lisinopril 30 mg/kg bwt(G7) | 5.63 ± 0.01 | 6.85 ± 0.57 ^a | 6.85 ± 0.56 ^{c,*} |

Values expressed as mean ± S.D. a= Statistically significant at P<0.05 when compared between 0 week and 4 weeks, b= when compared between 4 weeks and 8 weeks and c= when compared between 0 week and 8 weeks and * when compared within the groups and control

Table 9. Effect of methanolic extract of *Ocimum gratissimum* (OG) on serum creatinine level at 0 week (baseline), 4 weeks (after induction of hypertension) and 8weeks (after treatment with OG)

| Creatinine (µmol/l)n=8 | 0 week | 4 weeks | 8 weeks |
|-------------------------|--------------|---------------------------|--------------------------------|
| Neg control | 47.18 ± 1.49 | 47.17 ± 1.47 | 47.16 ± 1.47 |
| Untreated group | 47.67 ± 1.21 | 63.35 ± 3.77 ^a | 61.17 ± 4.95 ^{b,*} |
| OG 200 mg/kg bwt | 47.16 ± 1.47 | 62.00 ± 5.32 ^a | 51.33 ± 5.71 ^{b,c,*} |
| OG 400 mg/kg bwt | 47.33 ± 1.21 | 63.33 ± 1.03 ^a | 49.50 ± 3.72 ^b |
| Lisinopril 30 mg/kg bwt | 47.00 ± 0.89 | 66.00 ± 2.75 ^a | 68.50 ± 13.40 ^{b,c,*} |

Values expressed as mean ± S.D. a= Statistically significant at P<0.05 when compared between 0 week and 4 weeks, b= when compared between 4 weeks and 8 weeks and c= when compared between 0 week and 8 weeks and * when compared within the groups and control

Table 10. Effect of methanolic extract of *Ocimum gratissimum* (OG) on serum creatine kinase(CK) level at 0 week (baseline), 4 weeks (after induction of hypertension) and 8weeks (after treatment with OG)

| CK(U/L) n=8 | 0 week | 4 weeks | 8 weeks |
|-------------------------|------------|------------|------------|
| Neg control | 45.36±0.09 | 44.63±1.27 | 44.49±1.83 |
| Untreated group | 45.10±0.17 | 44.95±1.75 | 46.99±1.54 |
| OG 200 mg/kg bwt | 45.37±0.07 | 44.39±0.92 | 43.58±1.21 |
| OG 400 mg/kg bwt | 45.13±0.24 | 43.92±0.95 | 43.72±0.66 |
| Lisinopril 30 mg/kg bwt | 44.97±0.94 | 43.82±1.08 | 43.48±1.36 |

Values expressed as mean ± S.D. a= Statistically significant at P<0.05 when compared between 0 week and 4 weeks, b= when compared between 4 weeks and 8 weeks and c= when compared between 0 week and 8 weeks and * when compared within the groups and control

Table 11. Effect of methanolic extract of *Ocimum gratissimum* (OG) on serum CK-MB level at 0 week (baseline), 4 weeks (after induction of hypertension) and 8weeks (after treatment with OG)

| CK-MB(U/L) n=8 | 0 week | 4 weeks | 8 weeks |
|-------------------------|------------|------------|------------|
| Neg control | 14.47±0.03 | 14.48±0.03 | 14.48±0.03 |
| Untreated group | 14.48±0.03 | 15.28±0.05 | 15.15±1.00 |
| OG 200 mg/kg bwt | 14.47±0.03 | 15.14±0.39 | 14.33±0.03 |
| OG 400 mg/kg bwt | 14.49±0.01 | 15.19±0.14 | 14.32±0.08 |
| Lisinopril 30 mg/kg bwt | 14.48±0.02 | 15.39±0.08 | 15.18±0.18 |

Values expressed as mean ± S.D. a= Statistically significant at P<0.05 when compared between 0 week and 4 weeks, b= when compared between 4 weeks and 8 weeks and c= when compared between 0 week and 8 weeks and * when compared within the groups and control

4. DISCUSSION

Hypertension was induced in male Wistar rats to look at the possible prevention and management of hypertension using methanolic extract of *Ocimum gratissimum*(OG) in comparison with a reference drug, lisinopril. Induction of hypertension was achieved through oral gavage administration of 8% NaCl in water for a period of 4 weeks. The induction process was in agreement with work done by Rini [29]. The oral dosing of 8% NaCl was performed without anaesthesia. The NaCl induced hypertension may have been achieved through increased sympathetic nerve activity, a major trigger of vasoconstriction as well as the activation of renin angiotension aldosterone system [30]. Phytochemical analysis revealed the presence of steroid, flavonoids, phenol, alkaloids, saponin, tannis and phytate while mineral content revealed the presence of potassium, magnesium, phosphorus, calcium and sodium. The phytochemical result obtained was in accordance with Udochukwu [31]. After 4 weeks, systolic and diastolic blood pressures were significantly elevated in all the groups except group 1 which is the control group. On treatment with 200 mg/kg OG, 400 mg/kgOG and 30 mg/kg lisinopril, the elevated SBP and DBP significantly reduced ($P<0.05$). The significant reduction in SBP and DBP may be related to the high concentration of potassium and magnesium in methanolic extract of *Ocimum gratissimum*. Potassium and magnesium are important for muscle function, which includes relaxing the walls of blood vessels. Normal potassium levels are important for the conduction of electrical signals in the nervous system and in the heart. This protects against an irregular heartbeat. Potassium counters the effect of sodium which was significantly increased after 4 weeks induction with NaCl. At 8 weeks, serum sodium level was significantly reduced and potassium was significantly increased. Urea and creatinine were significantly increased at 4weeks suggesting impairment of renal function or loss of structural integrity of the kidney cell membrane but at 8weeks, the elevated renal parameters were reduced significantly. The observed changes is in accordance with the report of Ogundipe [32] who reported on the effects of two weeks administration of *Ocimum gratissimum* leaf on feeding pattern and markers of renal function in rats treated with gentamicin. Ogundipe observed a decrease in urea and creatinine although it was for a shorter period of 2 weeks. Reduction in the renal markers suggests that methanolic extract of

Ocimum gratissimum supplementation may lead to increase in glomerular filtration rate hence maybe helpful in hypertensive nephropathy. Methanolic extract of *Ocimum gratissimum* did not significantly affect the serum levels of creatine kinase(CK)and creatine kinase-MB which makes the cardioprotective effect of the extract uncertain in this study.

5. CONCLUSION

Ocimum gratissimum possesses antihypertensive effect and enhances the proper functioning of the kidney. It may also be useful in hypertensive condition due to its nephro-protective effect at 200 mg/kgbw and 400 mg/kgbw.

CONSENT

It is not applicable.

ETHICAL APPROVAL

The ethical approval for this study was obtained from the Ethics Committee of Nnamdi Azikiwe University Teaching Hospital, Nnewi with approval number: NAUTH/CS/66/VOL.9/145/2016/119.

ACKNOWLEDGEMENT

I wish to acknowledge Iloh EO for his financial support during the practical aspect of the work. Also Prof Meludu and Prof Dioka in the manuscript preparation.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Casanova LM, da Silva D, Sola-Penna M, Camargo LM, CelestriniDde M, Tinoco LW, et al. Identification of chicoric acid as a hypoglycemic agent from *Ocimum gratissimum* leaf extract in a biomonitoring *in vivo* study. *Fitoterapia*. 2014;93:132–41.
2. Bhuvaneshwari K, Gokulanathan A, Jayanthi M, Govindasamy V, Milella L, Lee S, et al. Can *Ocimum basilicum* L. and *Ocimum tenuiflorum* L. *In vitro* culture be a potential source of secondary metabolites? *Food Chemistry*. 2016;194:55–60.

3. Al Abbasy DW, Pathare N, Al-Sabahi JN, Khan SA. Chemical composition and antibacterial activity of essential oil isolated from Omani basil (*Ocimum basilicum* Linn.) Asian Pacific Journal of Tropical Diseases. 2015;5:645–9.
4. Madhu KD, Harindran J. Antiarthritic potential of *Ocimum gratissimum* L. in collagen induced arthritic Sprague-Dawley rats. Biomedicine and Aging Pathology. 2014;4:191–6.
5. Okoli CO, Ezike AC, Agwagah OC, Akah PA. Anticonvulsant and anxiolytic evaluation of leaf extracts of *Ocimum gratissimum* a culinary herb. Pharmacognosy Research. 2010;2:36–40.
6. Stanley IO, Isamila AU, Dorcas BJ, Hajiya MI. Antidiabetic potential of *Ocimum gratissimum* leaf fractions in fortified Diet-fed streptozotocin treated rat model of type-2-diabetes. Medicine (Basel). 2017; 4(4):73.
7. Ikpesu TO, Ezenwaka CO. *In vivo* screening of the extract of *Solenostemon monstachyus* and *Ocimum gratissimum* for the management of type-2-diabetes. Current Research in Diabetes and Obesity Journal. 2018;7(5):555723.
8. Oboh G. Tropical green leafy vegetables prevent garlic-induced hepatotoxicity in rat Journal of Medicinal Food. 2009;9:545-551.
9. Omodamiro OD, Jimoh MA. Antioxidant and antibacterial effect of *Ocimum gratissimum*. American Journal of Phyto-medicine and Clinical Therapeutics. 2015; 3(10):010-019.
10. Oparaocha ET, Iwu I, Ahaneke JE. Preliminary study on mosquito repellent and mosquitocidal activities of *Ocimum gratissimum* (L) grown in eastern Nigeria. Journal of Vector Borne Disease. 2010; 47(1):45-50.
11. Arhoghro EM, Ekpo KT, Ibeh GO. Effect of aqueous extract of scent leaf (*Ocimum gratissimum*) on carbon tetrachloride(CCl₄) induced liver damage in albino wistar rats. African Journal of Pharmacy and Pharmacology. 2009;3(11):562-563.
12. Surana V, Jain D. Hepatoprotective effect of *Ocimum gratissimum* against carbon tetrachloride induced hepatic damage model in rats. Pharmacology Online. 2010; 2:1111-1119.
13. Delacroix S, Chokka RC, Worthley SG. Hypertension: Pathophysiology and treatment. Journal of Neurology and Neurophysiology. 2014;5:250.
14. Lolodi O, Eriyamemu GE. Effect of methanolic extract of *Vernonia amygdalina* (common Bitter leaf) on lipid peroxidation and antioxidant enzymes in rats exposed to cycasin. Pakistan Journal of Biological Science. 2013;16:642-646.
15. Kurtz TW, Griffin KA, Pridani AK, Davisson RL, Hall JE. Recommendation for blood pressure measurement in humans and experimental animals – Part 2: Blood pressure measurement in experimental animals – A statement of professionals from the subcommittee of professional and public education of the American Heart Association Council on High Blood Pressure Research. Hypertension. 2005; 45:229-310.
16. Lorke D. A new approach to practical acute toxicity testing. Archive Toxicology. 1983; 54(275-287).
17. Enevide C, David A, Fidelis SA. A new method for determining acute toxicity in animal models. Toxicology International. 2013;20(3):224-226.
18. Harborne JB. Methods of plant analysis. In: Phytochemical methods. Chapman and Hall, London; 1973.
19. Bohm BA, Kocipai-Abyazan R. Flavonoid and condensed tannis from the leaves of *Vaccinium raticulation* and *Vaccinium calcyimium*. Pacific Science. 1994;48:458-463.
20. Obadoni BO, Ochuko PO. Phytochemical studies and comparative efficacy of the crude extract of some homeostatic plants in Edo and Delta states of Nigeria. Global Journal of Pure and Applied Science. 2001;8:203-208.
21. Pearson D. Chemical analysis of food. 8th Edition. Churchill Livingstone. 1974;5-63.
22. Khan AM, Qureshi RA, Ullah F, Gilani SA, Nosheen A, Sahreen S, Laghari MY, Shafiq-Ur-Rehman, Hussain I, Murad W. Phytochemical analysis of selected medicinal plants of Margalla Hills and surroundings. Journal of Medicinal plant Research. 2011;5:6017-6023.
23. Lucas GM, Markakes P. Phytic acid and other phosphorus compounds of nevy bean (*Phaseolous vulgaris*). Journal of agricultural and food chemistry. 1975;23: 13-15.
24. AOAC. AOAC, Official Methods of analysis. Gaithersburg, 15th ed. MD, USA,

- association of Official Analytical chemists, Arlington, VA; 1990.
25. Kassirer JP. Clinical evaluation of kidney function, glomerular function. *New England Journal of Medicine*. 1971;9:285-385.
 26. Laron K. Creatinine assay by reaction kinetic approach. *Clinica Chemica Acta*. 1972;41:209-217.
 27. Szasz G, Gerhardt W, Gruber W. Creatine Kinase in serum: 3. Further study of adenylate Kinase inhibitors. *Clinical Chemistry*. 1977;23:1888.
 28. Gerhardt. Immunoinhibition method of CK-MB determination. *Clinica Chemistry Acta*. 1977;79:29.
 29. Rini S, Elin YS, Tommy A, Joseph IS. The effect of coconut water (*Cocos nucifera* L.) and an isotonic drink on the change of heart rate frequency in the rats induced hypertension. *Procedia Chemistry*. 2014; 13:177-180.
 30. Blaustein MP, Zhang J, Chen L, Hanolton BP. How does salt retention raise blood pressure? *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology*. 2006;290(3):514-523.
 31. Udochukwu U, Omeje FI, Uloma IS, Oseiwe FD. Phytochemical analysis of *Vernonia amygdalina* and *Ocimum gratissimum* extracts and their antibacterial activity on something resistant bacteria. *American Journal of Research Communication*. 2015;3(5):228-235
 32. Ogunidipe DJ, Akomolafe RO, Sanusi AA, Imafidon CE, Olukiran OS, Oladele AA Effects of two weeks administration of *Ocimum gratissimum* leaf on feeding pattern and markers of renal function in rats treated with gentamicin. *Egyptian Journal of Basic and Applied Sciences*. 2016;3(3):219-231.

© 2018 Onyema-Iloh et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<http://www.sdiarticle3.com/review-history/45379>