

Ethnobotany of Aromatic Plants used by Phytotherapists in the District of Glazoue, Ouesse and Save in Benin

Assongba Yédjanlognon Faustin ^{a,b*}, Konko Cossi Jacques ^a, Yoka Joseph ^c,
Kpogniwe K. Joel ^a, Djego G. M. Julien ^b and Brice A. Sinsin ^b

^a *Laboratory of Plant Biology and Traditional Pharmacopoeia (LaBVephaT), Ecole Nationale Supérieure des Biosciences et Biotechnologies Appliquée (ENSBBA), Université Nationale, des Sciences, Technologies, Ingénierie et Mathématiques (UNSTIM) Abomey, Benin.*

^b *Laboratoire d'Ecologie Appliquée, Faculté des Sciences Agronomiques, Université d'Abomey-Calavi, Benin.*

^c *Laboratory of Biodiversity, Management of Ecosystems and Environment, Faculty of Sciences and Techniques, Université Marien NGOUABI, Brazzaville, BP: 69, Congo.*

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/ARRB/2021/v36i1130452

Editor(s):

(1) Prof. Md. Aminur Rahman, Jashore University of Science and Technology, Bangladesh.

Reviewers:

(1) Emmanuel AKINTIMEHIN, Olusegun Agagu University of Science and Technology, Nigeria.

(2) Mohammad Manjur Shah, Yusuf Maitama Sule University, Nigeria.

(3) Irene Esi Donkoh, University of Cape Coast, Ghana.

Complete Peer review History, details of the editor(s), Reviewers and additional Reviewers are available here:
<https://www.sdiarticle5.com/review-history/77411>

Original Research Article

Received 15 September 2021

Accepted 25 November 2021

Published 04 December 2021

ABSTRACT

Aromatic and medicinal plants (AMP) are flora with a fragrant essence. The aim of this study is to identify the aromatic plants used by herbalists in three districts of the Département des Collines. The method of data collection is based on ethnobotanical surveys of herbalists using the Dagnelli formula and pre-established survey questionnaires. The results of the survey of 300 herbalists identified 22 most commonly used aromatic plants, divided into 17 families and 21 genera. The most represented families are Lamiaceae (15%); Annonaceae (15%). Among the MAPs inventoried, the most cited and used are: *Ocimum canum* L. *Ocimum gratissimum* L.; *Chenopodium ambrosioides* L.; *Hyptis suaveolens* L.; *Xylopia aethiopica* (Dunal) A. Rich; *Securidaca longepedunculata* Fresen. MAPs treat 163 conditions identified in this study. Some of the AMP are used alone or in

*Corresponding author: E-mail: yedjanlognon@gamil.com, yedjanlognon@yahoo.fr;

combination with other plants. AMP drugs are: leaves (64%), roots (23%). The ailments treated by the majority of AMP are: fever; tooth decay; conjunctivitis; constipation; haemorrhoids; hernia; inflammation; skin lesion; stomach ache; snakebite; causes of malaria; sores and painful periods. In view of the proven importance and different uses of AMP, it is necessary to study them further (ecology, distribution, and conservation status) in order to better prevent or cure human health.

Keywords: Aromatic plants; useful plants; diseases; Healing and Benin.

1. INTRODUCTION

The floristic resource constitutes a huge reservoir that is very important for health needs [1-2]. The vast majority of the Beninese population remains dependent on the plant for the satisfaction of daily needs [3]. Plants, since then, have been used as medicines and even under certain conditions as food [4]. According to the World Health Organization (WHO), about 65-80% of the world's population, use medicinal plants in primary health. According to Agbangnan et al, [5], 25-50% of conventional medicines nowadays have the bioactive molecules of medicinal plants as active ingredients. The high cost of modern medicine drugs makes people resort to traditional medicine (medicinal plants) for treatment. Medicinal plants are an inevitable source of medicine [6]. Modern medicine has turned to ethnobotanical studies to have a wide range of varieties of molecules from medicinal plants to prevent and fight against diseases [7]. In Benin, traditional medicine (medicinal plants) has become increasingly important in the treatment of diseases. The promotion of aromatic and medicinal plants is a major concern of many researchers and institutional leaders [8-9]. Aromatic and medicinal plants (AMPs) useful to humans can be categorized according to uses: herbal tea plant; hygienic and amenity drink; plant for cosmetic use; plant for aromatic and condiment use; plant for food use; plant for industrial use; medicinal plant. Indeed, aromatic and medicinal plants are used in phytomedicines to bring healing and even preventive [10]. Enough authors have demonstrated the therapeutic aspect of aromatic plants [8,11]. Some of the MAPs are used in the agricultural field to control crop pests to promote good agricultural yield [12]. MAPs are generally used in food for the purpose of flavoring sauce and drink. The recognition of the clinical, pharmaceutical, and economic value of MAP-based medicines continues to grow. MAPs are considered a commodity chain in their own right [10]. MAPs were at the origin of the first botanical gardens [13,10,14]. Different ethnobotanical works have been carried out in Benin to document and perpetuate the traditional

knowledge of MAPs. These are: Adjanohoun [15], Houessou [16], Adomou et al. [8] and Assongba [1]. Beninese plant formations are dwarfed in quality by aromatic and medicinal plants [17]. A single plant species is used in different ways and in different areas, in different localities where it exists according to the needs and knowledge knew about this plant [9]. Despite the numerous studies on medicinal plants in Africa [18] and elsewhere in the world [19-20] it should be noted that enough studies carried out on aromatic plants are not yet done in Benin. As for the one in the hills department, it is the first study of its kind. This is one of the reasons why this study on aromatic and medicinal plants used by phytotherapists in three districts of the department of the hills was chosen in order to list these aromatic and medicinal plants with a view to their perpetuation.

2. MATERIALS AND METHODS

2.1 Study Environment

The study took place in the Collines Department, which has an area of 13,931 km². This Department is composed of six (6) District (Dassa-Zounmè, Glazoué, Ouèssè, Savalou, Savè and Bantè). The district of this department is subjected to the influences of the humid tropical climate of the Sudano-Guinean type with two rainy seasons with hazards. They have vegetation of wooded forests and shrubby savannahs. The Ouémé River runs through the study districts. This study took into account three (3) districts of the six (6) of the department, which are: the district of Glazoué, Savè, and Ouesse. The districts are selected because of their diversity of languages, plant formations, and high human population.

2.2 Study Materials

The biological material (plant and human) is made up of the phytotherapists of the three districts (Glazoué, Savè, and Ouèssè) in which the survey was carried out, as well as all the aromatic and medicinal plants that are listed.




Fig. 1. Location of the Study Area

2.3 Ethnobotanical Survey

The choice of villages was made in the dynamics of having representativeness of the sociolinguistic groups of the study area. The

choice of study villages was based on the existence of herbalists who use medicinal and aromatic plants [1,6] A total of 71 villages were visited and 300 herbalists were surveyed by using Dagnilli's (1998) formula at a rate of 100

per district. The herbalists questioned (men and women) were those who had knowledge of plant aromas. The information concerning the symptoms or causes of diseases was confirmed with local doctors, nurses (health agents) and completed by the bibliographical review (Adjanohoun et al., 1989). For practical exploitation of the data and harmonization with the international system.

2.4 Data Collection

In each district surveyed, two types of data were collected: the first related to the identification of the respondent (ethnicity, sex, age, level of education, religion, marital status, etc.) and the second related to the knowledge of aromatic and medicinal plants provided by the herbalists (vernacular names of the plants, characteristic features of the aromatic plant, the difference between aromatic and medicinal plants, plant drugs used). During this phase of field investigation, the various discussions were conducted in an open manner and without time limits to allow the herbalists to freely express their knowledge.

2.5 Data Processing

The collected data are entered in the Excel spreadsheet to have a database. These data were used to calculate means, variances, frequencies to build tables. R 643.4.0 software is used for linear regression, multivariate analysis etc. The Relative Citation Frequency (RCF) of species [21] is calculated as follows

FRC = FC/N; FC: Frequencies of quotations; FRC: Relative Frequencies of Quotation; N: Number of respondents. Jaccard Similarity Index is calculated to compare sociolinguistic groups. This index measures the similarity of the sociolinguistic groups taken two by two with regard to the knowledge of aromatic plants. It was calculated for each group of species according to the following formula:

$J = C / (A + B - C)$ with A: number of species of aromatic plants cited by sociolinguistic group A; B: number of species of aromatic plants cited by sociolinguistic group B; C: number of species of aromatic plants common to both groups A and B. If $J \leq 40\%$; the ethnic groups have different knowledge about aromatic plants. If on the other hand $J > 40\%$; the ethnic groups have similar knowledge about aromatic plants.

3. RESULTS

3.1 Endogenous Knowledge of Aromatic Medicinal Plants

The surveys carried out among the phytotherapists made it possible to note that 98% of the phytotherapists surveyed in the various district know and use aromatic and medicinal plants. Thus, 23 species of aromatic plants are listed and used. They are divided into 17 families and 21 genera (Table 1). The most represented families are Lamiaceae (15%), Annonaceae (15%), Alliaceae (11%), and Rutaceae (7%). According to the results, 100% of five (5) ethnic groups (Idaasha, Mahi, Nago, Fon and Pheul) know and use more plants. The vernacular names of the identified aromatic plants and their frequency of citation are summarized in Table 1. The plants with a frequency of citation (FRC) of more than 50% are: *Ocimum canum*; *Ocimum gratissimum*; *Momordica Charantia*; *Chenopodium ambrosioides*; *Hyptis suaveolens*; *Lippia multiflora Moldenke*; *Lippia rugosa*; *Xylopi aethiopica*; *Heliotropium indicum*; *Securidaca longepedunculata*. These plants are the most frequently used in the treatment of ailments.

3.2 Jaccard Similarity Index (J)

The analysis of Table 2 allows us to say that the different sociolinguistic groups have a similar knowledge about aromatic plants ($J \geq 40$), only that the similarity differs according to each sociolinguistic group.

3.3 Diseases Treated with Aromatic and Medicinal Plants

Table 3 shows the aromatic and medicinal plants with the different ailments treated, the drugs used in the phytomedicines, and the modes of preparation of phytomedicines. A total of 24 medicinal plants were identified in the study area. These plants are used daily by herbalists and the general population to season sauces and to treat ailments.

3.4 Diseases and Ailments Treated by the Inventoried Plants

Table 3 presents the different diseases treated by the 24 plants inventoried with the methods of preparation and use. In total, 20 disease symptoms or conditions were identified. The

ailments treated by the majority of the plants are Fever, attacks, and seizures; tooth decay, conjunctivitis, constipation, tastes and their odor, aches, arousals, digestion facilitator, internal and external hemorrhoids, hernia, hypotension and hypertension, inflammation, skin lesion, stomach

ache, snakebite, malaria, wounds, painful period, intestinal worms and vomiting. The frequency of plant drugs used is as follows: leaf (64%), root (23%), bark (8%), bulb (3%), stem and seeds/fruits are weakly used. Fugue 2 reflects the frequency of drug use.

Table 1. Vernacular names and citation frequencies of aromatic plants

Scientific names	Idaash	Nago	Mahi	FRC
<i>Ocimum canum</i> L.	Ina-éwé/Hisihisi	Owofi	Akeman	0,88
<i>Ocimum gratissimum</i> L.	Alibara	Simonba	Tchao	0,84
<i>Momordica Charantia</i> L.	Kpalari	Kpalari	Yinssikin	0,82
<i>Chenopodium ambrosioides</i> L.	Odanmaira	Orèdjougou	Godo	0,67
<i>Hyptis suaveolens</i> L.	Ewéidjako	Couloubi	Sansoukpèman	0,66
<i>Lippia multiflora</i> Moldenke	-	-	akalala	0,65
<i>Lippia rugosa</i> L. A.Chev.	Kanhoun	Kinhoun-kinhoun	Aglala	0,62
<i>Xylopi aethiopica</i> (Dunal) A. Rich.	Oroun	Aroun	Kpejelekou	0,58
<i>Heliotropium indicum</i> L.	Ogbolarouko	Igbéako	Koclosoudèkpadja	0,55
<i>Annona senegalensis</i> Pers. Pers.	Otribobo	Ambo	Nyiglwé	0,52
<i>Allium ascalonicum</i> auct. non Strand	Mansa Kpikpa	Amansa elewe	Masa winiwini	0,5
<i>Securidaca longepedunculata</i> Fresen.	Kpatalè	Ikpata	Atakpa Wanlwi	0,49
<i>Ocimum basilicum</i> L.	Akohoun	Akohoun	Kohoun	0,43
<i>Zanthoxylum zanthoxyloides</i> (Lam.)	Eguiatta	Eguiatta	Hetti	0,36
<i>Allium sativum</i> L.	Ayo	Ayu	Ayo	0,35
<i>Cymbopogon citratus</i> (DC.) Stapf	Ofrin	Ewéiti	Tcha	0,35
<i>Zingiber officinale</i> Griff.	Atalè	Atalè	Dotan	0,3
<i>Crassocephalum crepidioides</i> (Benth)	Gbolo	Gbolo	Kogbo	0,3
<i>Piper guineense</i> Schum. et Thonn.	Idjayé	Ata iyere	Linlinkou	0,27
<i>Allium cepa</i> L.	Masa	Amansa	Masa	0,25
<i>Olax subscorpiodea</i> Oliv.	Miitin	Akpatcha	Kiviman	0,21
<i>Monodora myristica</i> (Geartn).	Ariwo	Ariwo	Sasalikouin	0,19
<i>Uvaria chamae</i> P.Beauv.	Yaha	Yaha	Yaha	0,17

Table 2. Jaccard similarity indices

Comparative ethnicities	Idaasha and Mahi	Idaasha and Nago	Mahi and Nago
Index J	86%	90%	95%




Fig. 2. Frequency of drug use

Table 3. Plants identified, drugs used, preparations made and conditions treated

Species	Families	Org	Preparation	Diseases and conditions
<i>Allium ascalonicum</i> auct. non Strand	Alliaceae	Bu/Fe	Sauce (gold)	Flavouring, Angina, Aid digestion
<i>Allium cepa</i> L.	Alliaceae	Bu/Fe	Maceration (Bu/Fe) (Spread) Sauce (gold)	Anti-snake, Infection, Angina
<i>Allium sativum</i> L.	Alliaceae	Bu	Sauce (gold)	Aromatizing, Cough, aids digestion
<i>Annona senegalensis</i> Pers.	Annonaceae	R/Fe/E	Leaves (pocket) Decoction (Bath, Gold)	Anti-snake, dizziness, stress
<i>Monodora myristica</i> (Gearn).	Annonaceae	R/ Fr	Decoction (gold) Market	Intestinal worms, infection, cyst
<i>Uvaria chamae</i> P.Beauv.	Annonaceae	R/Fe/E	Decoction (bath, gold)	Fever, body aches, kidney pain
<i>Xylopia aethiopica</i> (Dunal) A. Rich.	Annonaceae	Fr/Fe	Sauce (gold) Maceration (gold)	Flavouring, a recipe activator
<i>Crassocephalum crepidioides</i> (Benth)	Asteraceae	Fe/R	Sauce (gold) Maceration (gold)	Flavouring, Used by women alert
<i>Heliotropium indicum</i> L.	Boraginaceae	Fe/FI	Decoction or trituration of leaves (gold)	Sexual weakness, Coughing in babies, Hypotension
<i>Chenopodium ambrosioides</i> L.	Chenopodiaceae	Fe/Gr	Trituration (past) Grain (gold)	Skin lesion, abscesses Deworming of poultry,
<i>Combretum collinum</i> Fresen.	Combretaceae	R/F/E	Decoction (bath, gold)	Awakening the Baby, Jaundice, Fatigue
<i>Lippia multiflora</i> Moldenke	Verbenaceae	Fe / Fr	Sauce / Decoction	High blood pressure
<i>Momordica Charantia</i> L.	Cururbitaceae	Fe/Fr	Trituration or maceration	Stomach ache, Hernia, Itching
<i>Hyptis suaveolens</i> L.	Lamiaceae	Fe/T	Decoction (bath, gold)	Baby bath, insecticide, jaundice
<i>Ocimum basilicum</i> L.	Lamiaceae	Fe/R	Sauce Trituration	Aromatizer Meat preservative
<i>Ocimum canum</i> L.	Lamiaceae	Fe/R	Trituration (bath, past)	Purification and disinfectant
<i>Ocimum gratissimum</i> L.	Lamiaceae	Fe/Pe	Sauce or trituration	Flavouring, Digestion, Intestinal worms
<i>Khaya senegalensis</i> (Desr.) A. Juss. (Desr.) A. Juss.	Meliaceae	E/Fe/R	Decoction/Maceration (Gold/Bath)	Purification, Disinfectant, Baby's first bath
<i>Musa sapientum</i> auct. div.	Musaceae	Fr/FI	Powder (Gold)	Memory aid, Sexual weakness, Paralysis
<i>Musa sp</i> L.	Musaceae	Fr/Fe/R	Powder (Gold)	Sexual weakness, checklist
<i>Olax subscorpiodea</i> Oliv.	Olacaceae	R/Fe	Decoction (bath, gold)	Protection, Child development, Palu
<i>Piper guineense</i> Schum. et Thonn	Piperaceae	Gr	Decoction (gold)	Flavouring, Angina, Fibroma, Intestinal worms
<i>Cymbopogon citratus</i> (DC.) Stapf	Poaceae	Fe/Pe	Decoction or Maceration (Gold)	Flavouring, Convalescence, Lack of blood
<i>Securidaca longepedunculata</i> Fresen.	Polygalaceae	R/E	Racine at your fingertips Decoction (Bath, gold)	Antivenomous, Sexual weakness, Protection

Preparations: Gold: Oral; Dec: Decoction; Mac: Maceration

As regards the methods of preparation, trituration is the most used method (46%); decoction (28%). Maceration and powdering of plant drugs are in equal proportion (13%). Infusion is poorly represented.

Some plants like : *Ocimum canum*, *Ocimum basilicum*, *Ocimum gratissimum*, *Zingiber officinale*, *Lippia multiflora* Moldenke; *Lippia rugosa*, *Crassocephalum crepidioides*, *Allium sativum*, *Xylopiya aethiopic*a, *Allium Ascalonicum*, *Allium cepa*, *Piper guineense* are used to flavor sauce and food. *Ocimum canum* L. is used to purify places of worship; *Ocimum basilicum* L. is used by hunters and fishermen to preserve meat and fish respectively. Plants like *Allium sativum* L. *Xylopiya aethiopic*a (Dunal) A. Rich; *Allium ascalonicum* auct. non Strand; *Allium cepa* L and *Monodora myristica* (Geartn) are plants used by Phyto therapists to accelerate and precipitate the activity of phytomedicines during their use.

4. DISCUSSION

4.1 Strengths and Weaknesses of the Survey Methodology

The ethnobotanical survey method based on individual interviews allowed us to know the importance of the use of aromatic and medicinal plants in the treatment of ailments and diseases in the hills department. This method, which requires the memory of the interviewees, could cause biases related to the personal appreciation of the respondents [22,3]. The importance given to the use of the species is given by individuals who implicitly take into account a personal appreciation; which often refers to their preference. Nevertheless, these methods are widely used in ethnobotany by other authors and have the privilege to bring out quite relevant results, Nguenang [23] Dahouenon et al. [12]. As the frequencies of the different uses of these species for the different uses identified among herbalists are not precise, this could cause errors when assigning scores for the calculation of the various indices that can measure the importance of a species for a given sociolinguistic group or groups. It should also be noted that during the survey, exchanges were made in languages other than those mastered by the respondent, which required interpreters.

4.2 Level of Knowledge on Aromatic and Medicinal Plants

Ethnobotanical knowledge of aromatic species in the hills varies according to sociolinguistic groups

on the one hand and within sociolinguistic groups on the other hand, which means that a small group of community holds most of the knowledge about the plants. These results are similar to those of Quiroz et al. [24] Wedjangnon et al. [3]. Other studies have also shown that ethnobotanical knowledge related to plant species varies mostly with age, gender, and sociolinguistic group Ekué et al [18] Fandohan et al. [25] Gouwaknnou et al. [26] Assogbadjo et al. [27] and Assongba [1]. This study was conducted among herbalists and it should be noted that these people were reluctant on information, this is moreover the reason for taking a large enough sample to be able to mobilize enough information, this confirms the results of Adomou [8], on a study conducted among women plant sellers in the market of Cotonou. The information collected shows that plant species are of great wealth for humans. In general, therefore, it can be said that sociolinguistic groups know and use the plants around them according to their way of life. It is necessary to emphasize that any plant not found in the country is imported. This is similar to that of Adomou et al [8]. The botanical families most found are Lamiaceae (15%) and Annonaceae (15%), these results are similar to those of Salhi et al. [28] in a floristic and ethnobotanical study of medicinal plants in Morocco. Some species of aromatic plants such as *Allium sativum*; *Xylopiya aethiopic*a; *Allium Ascalonicum*; *Allium cepa* and *Monodaro myristica* have a particular character of accelerating the activity of recipes. This is confirmed by the findings of Adomou et al. [8] Dougnon et al. [29-30] which states that some drugs, especially fruits, seeds are used as auxiliary or adjuvant ingredients, enable to enhance the therapeutic action of the main components of the recipes and also enable to treat the secondary symptoms of the disease [31-32].

5. CONCLUSION

The ethnobotanical survey revealed the existence of numerous aromatic and medicinal plants used by traditional therapists who have medicinal and magico-spiritual knowledge and know well the use of plants as aliments. The study made it possible to constitute a database on the main aromatic species used by the phytotherapists of the hills in general and the district of Glazoué, Savè, and Ouesse in particular. A total of 23 species of aromatic plants were identified to treat 20 different symptoms of illness. The ethnic groups surveyed were

Idaasha, Mahi, Nago, Fon, and Peuhl. These sociolinguistic groups know and use the plants for various purposes, which reveals the importance that people give to natural plant resources in the localities. The data collected will be used for the promotion of aromatic and medicinal plants with a view to discovering new active ingredients for use in therapeutics, cosmetics, perfumery, and others.

CONSENT

As per international standard or university standard, respondents' written consent has been collected and preserved by the authors.

NOTE

The study highlights the efficacy of "traditional medicine" which is an ancient tradition, used in some parts of Benin. This ancient concept should be carefully evaluated in the light of modern medical science and can be utilized partially if found suitable.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Assongba F.Y. Ethnobotany, Ecology, Spatial distribution and conservation status of *Dialium guineense* Willd (Fabaceae) in Benin. Single doctoral dissertation. Aménagement et Gestion des Ressources Naturelles; University of Abomey-Calavi. 2014;245.
2. Akodewou A, Akpavi S, Dourma M, Batawila K, Amegnaglo KB, Atakpama W, Akpagama K. *Sorindeia warneckei* Engl. (Anacardiaceae), a multi-use species from the Lama depression in Togo. *Africa Science: International Journal of Science and Technology*. 2014;10(2).
3. Wedjangnon AA, Houetchegno T, Ouinsavi C. Ethnobotanical characteristics and socio-cultural importance of *Mansonia altissima* A. Chev. in Benin, West Africa Laboratoire d'Études et de Recherches Forestières (LERF), Faculté d'Agronomie, Université de Parakou, Parakou, Bénin. 2016;22:4678-469.
4. Din N, Mpondo E, Dibong SD, Kwin NF, Ngoye A. Inventory and identification of plants used in the treatment of diabetes in Douala town (Cameroon). *European Journal of Medicinal Plant*. 2011;1:60-73.
5. Agbangnan CPD, Noudogbessi JP, Chrostowska A, Tachon C, Fouquet E, Sohounhloue DCK. Phenolic compound of benin's red sorghum and their antioxidant properties. *Asian J Pharm Clin Res*. 2013;6(2):277-280.
6. Kpètèhoto WH, Hessou S, Dougnon VT, Christian R, Johnson GB, Houéto EE, Gbénou J. Ethnobotanical, phytochemical and ecotoxicological study of *Ocimum gratissimum* Linn (Lamiaceae) in Cotonou. *Journal of Applied Biosciences*. 2017;109:10609-10617.
7. Jiofack T, Fokunang C, Guedje N, Kemeuze V, Fongzossie E, Nkongmeneck BA, Tsabang N. Ethnobotanical uses of medicinal plants of two ethnoecological regions of Cameroon. *International Journal of Medicine and Medical Sciences*. 2010;2(3):60-79.
8. Adomou AC, Yedomonhan H, Djossa B, Legba SI, Oumorou M, Akoegninou A. Ethnobotanical study of medicinal plants sold in Abomey-Calavi market in Benin. *International Journal of Biological and Chemical Sciences*. 2012;6(2):745-772.
9. Ouinsavi C, Sokpon N, Alohou EC. Determinants of the fragmentation of the classified forest-sacred forest block in South Benin. *Journal of Applied Biosciences*. 2016;101(1):9618-9633.
10. Alfaiz C. (INRA-Edition). *Culture et Domestication des Plantes Aromatiques et Médicinales au Maroc*. 2015;10.
11. Mangambu MDD, Mushagalusa KF, Kadima NJ. Contribution to the photochemical study of some anti-diabetic medicinal plants from Bukavu city and its surroundings (South Kivu, DR Congo). *Journal of Applied Biosciences*. 2014;75(1):6211-6220.
12. Dahouenon-Ahoussi E, Kpatinvoh B, Adjou ES, Konfo TC, Atrevi B, Soumanou MM, Sohounhloue DC. Efficacy of essential oils of three aromatic plants against spoilage mycoflora of cowpea (*Vigna unguiculata* L., Walp) collected from sales shops in South Benin. *Journal of Applied Biosciences*. 2017;109(1):10680-10687.

13. Tardivon J, Mohamed C, Aromatic and medicinal plants, an example of human development in Morocco : The women's cooperative of Ben Karrich – Tetouan. 2012;19.
14. Neffati M, Sghaier M. Development and valorization of aromatic and medicinal plants (AMP) at the level of desert areas in the MENA region (Algeria, Egypt, Jordan, Morocco and Tunisia); 2014.
15. Adjanohoun EJ, Adjakidje V, Ahyi MRA. Aké assi LL, Akoegninou A, d'Almeida J, Apovo F, Boukef K, Chadare M, et al. 1999;55-63.
16. Houessou S. Effects of the reduction of floristic diversity on the health of rural populations in southern Benin. In Colloque international SIFEE, Paris; 2010.
17. Neuenschwander P, Milleret V, Simona B, Hall H. Tuning electrospinning parameters for production of 3D-fiber-fleeces with increased porosity for soft tissue engineering applications. Eur Cell Mater. 2011;21:286-303.
18. Ekué MR, Sinsin B, Eyog-Matig O, Finkeldey R. Uses, traditional management, perception of variation and preferences in ackee (*Blighia sapida* KD Koenig) fruit traits in Benin: implications for domestication and conservation. Journal of Ethnobiology and Ethnomedicine. 2010;6(1):12.
19. Pavani A, Naushad SM, Rupasree Y, Kumar TR, Malempati AR, Pinjala RK, Kutala VK. Optimization of warfarin dose by population-specific pharmacogenomic algorithm. The Pharmacogenomics Journal. 2012;12(4):306.
20. Thirumalai T, Beverly CD, Sathiyaraj K, Senthilkumar B, David E. Ethnobotanical Study of Anti-diabetic medicinal plants used by the local people in Javadhu hills Tamilnadu, India. Asian Pacific Journal of Tropical Biomedicine. 2012;2(2):S910-S913.
21. Tardío J, Pardo-de-Santayana M. Cultural importance indices: a comparative analysis based on the useful wild plants of Southern Cantabria (Northern Spain). Economic Botany. 2008 ;62(1):24-39.
22. Dossou ME, Houessou GL, Lougbégnon OT, Tenté AHB, Codjia JTC. Ethnobotanical study of woody forest resources of Agonvè swamp forest and related terroirs in Benin. Tropicultura. 2012;30(1):41-48.
23. Nguenang GM, Fedoung EF, Nkongmeneck BA. Importance of secondary forests for the collection of useful plants among the Badjoué of East Cameroon. Tropicultura,. 2010;28(4):238-245.
24. Quiroz D, Towns A, Legba SI, Swier J, Briere S, Sosef M, Van Andel T. Quantifying the domestic market in herbal medicine in Benin, West Africa. Journal of Ethnopharmacology. 2014;151(3):1100-1108.
25. Fandohan B, Assogbadjo AE, Kakaï RG, Kyndt T, De Caluwé E, Codjia JTC, Sinsin B. Women's traditional knowledge, use value, and the contribution of tamarind (*Tamarindus indica* L.) to rural households' cash income in Benin. Economic Botany. 2010;64(3):248-259.
26. Gouwakinnou GN, Lykke AM, Assogbadjo AE, Sinsin B. Local knowledge, pattern and diversity of use of *Sclerocarya birrea*. Journal of Ethnobiology and Ethnomedicine. 2011;7(1):8.
27. Assogbadjo AE, Glegrave R, Azihou AF, Kyndt T, Codjia JTC. Ethnic differences in use value and use patterns of the threatened multipurpose scrambling shrub (*Caesalpinia bonduc* L.) in Benin. Journal of Medicinal Plants Research. 2011;5(9):1549-1557.
28. Salhi S, Fadli M, Zidane L, Douira A. Floristic and ethnobotanical studies of medicinal plants in the city of Kenitra (Morocco). Lazaroa. 2010;31: 133.
29. Dougno TV, Agbankpé AJ, Bankole SH, Houngbégnon O, Dah-Nouvlessounon D. In vitro antibacterial effects of *Crateva adansonii*, *Vernonia amygdalina* and *Sesamum radiatum* used for the treatment of infectious diarrhoeas in Benin. Journal of Infectious Diseases & Therapy; 2016.
30. Diatta CD, Gueye M, Akpo LE. Medicinal plants used against dermatoses in the Bainouk pharmacopoeia of Djibonker, Ziguinchor region, Senegal. Journal of Applied Biosciences. 2013;70(1):5599-5607.
31. Ngoule CC, Ngene JP, Kidik CP, Ottou PM, Dibong SD, Mpondo EM. Importance in the traditional pharmacopoeia of

flavonoid plants sold in the markets of Douala East (Cameroon). Journal of Applied Biosciences. 2015;88(1):8194-8210.

32. WHO. Strengthening the role of traditional medicine in health systems: a strategy for the African region. Secretariat report. 2013;39.

© 2021 Faustin et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/77411>