



Prevalence of Dysmenorrhea and its Management among Undergraduate Students of the University of Port Harcourt, Nigeria

Faustina Chiamaka Irozulike ^{a*}, Nicholas Asiwe ^{a*},
Joy Wilberforce Ekokodje ^b, Masterson Sampson Wedeh ^a,
James Dibagha Izibeya ^c and Patrick Leo Filima ^a

^a Department of Anatomy, Faculty of Basic Medical Sciences, University of Port Harcourt, Port Harcourt, Nigeria.

^b Department of Human Anatomy, Faculty of Basic Medical Sciences, Rivers State University, Port Harcourt, Nigeria.

^c Department of Human Anatomy, Faculty of Basic Medical Science, Bayelsa Medical University, Yenagoa Bayelsa State, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJARR/2023/v17i12582

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/107877>

Original Research Article

Received: 19/08/2023

Accepted: 26/10/2023

Published: 06/11/2023

ABSTRACT

Background: Dysmenorrhea is a pain felt by women before or during menstruation and is accompanied by some symptoms like headache, dizziness, food cravings, mood swings, and fatigue. Many ways have been adopted for the management of dysmenorrhea. The study aims to evaluate the prevalence of dysmenorrhea and its management among university students.

*Corresponding author: Email: asiwenicholas@gmail.com;

Methods: A total of two hundred (200) female university students were involved in this study. A cross-sectional descriptive study design was used where students were sampled randomly a questionnaire was used to collect data and data obtained were analyzed using IBM SPSS version 26.

Results: The study found that a majority of the population experiences dysmenorrhea during menstruation (81.5%) and the majority of respondents experienced menstrual flow within 4-5 days with 37% experiencing severe pain. Symptoms include anxiety, mood swings, tender breasts, food cravings, fatigue, irritating mood, depression, dizziness, vomiting, diarrhea, and headache. Management of dysmenorrhea varies, with 23.5% using synthetic drugs and 1.5% using local and herbal practices.

Conclusion: There was a high prevalence of dysmenorrhea among the University students and presenting symptoms like mood swings, fatigue, and diarrhea, with most using local/herbal medication.

Keywords: Dysmenorrheal; women; headache; dizziness; university students.

1. INTRODUCTION

Menstruation is the monthly shedding of the lining of your uterus. Is a natural phenomenon that occurs throughout the reproductive years of every woman and most often it is accompanied by pain [1]. This pain varies in individuals but collectively the pain is termed dysmenorrhea. Dysmenorrhea is a painful menstruation [2]. It is a common condition experienced by many women and can range in severity from mild discomfort to debilitating pain which could have significant impacts on their daily activities and disturb their productivity at home, school, or workplace.

Dysmenorrhea is associated with many factors, including behavioral and psychological aspects [3], Symptoms include cramping or pain in the lower abdomen, low back pain, pain spreading down the legs, nausea, vomiting, diarrhea, fatigue, weakness, fainting, or headaches [4]. Dysmenorrhea is classified into two; primary and secondary dysmenorrhea where primary dysmenorrhea is menstrual pain that's not a symptom of an underlying gynecologic disorder but is related to the normal process of menstruation [5]. Primary dysmenorrhea is the most common type of dysmenorrhea, affecting more than 50% of women, and quite severe in about 10%. it is most common in late adolescence and the early 20s. while secondary dysmenorrhea is menstrual pain that is generally related to some kind of gynecologic disorder. Most of these disorders can be easily treated with medications or surgery. Secondary dysmenorrhea is more likely to affect women during adulthood [5].

Over the years, individuals have used either pharmacological or non-pharmacological

methods to relieve pain during the menstruation period [6]. The non-pharmacological methods used include fatty diet restriction, herbal remedies, exercise, and heat application while the medications used to relieve pain include nonsteroidal anti-inflammatory drugs (NSAIDs) as first-line therapy, combined contraceptives, antispasmodics, and vitamins [6]. The prevalence of either of these methods used by individuals has become the object of this study. Though studies on the prevalence of dysmenorrhea abound in our communities; 78.1% in Ogun State [7], 62% in Oyo State [8], 58% in Osun State [9], 60-64% in Ile-Ife [10,11], and 69.8% in Maiduguri [12]. Also, some studies that had been carried out in other countries have shown the prevalence of dysmenorrhea shown 85.6% in Kuwait [13], 79.4% in Saudi Arabia [14], 84.2% in India [15], 74.8% in Spanish [16] and the prevalence of primary dysmenorrhea was 80.9% in Lebanon [17] and 92% in Riyadh [18].

literature review on the Prevalence of dysmenorrhea varies greatly across different populations. Therefore, the present study aims to evaluate the prevalence of dysmenorrhea and the possible strategies or methods used to manage the pain among the female university Students of the University of Port Harcourt, Nigeria. To improve the quality of life and productivity of women experiencing dysmenorrhea.

2. MATERIALS AND METHODS

2.1 Study Design

The study adopted a cross-sectional study design that was used to generate data from university students, a total of two hundred (200)

female students were involved in the study, drawn from different levels of undergraduate programs. A multistage random sampling technique was used where the sample size was determined using the Cochran formula of a descriptive survey [19, 20].

$$Sample\ size = \frac{Z_{1-\alpha/2}^2 p(1-p)}{d^2}$$

Where;

$Z_{1-\alpha/2}$ = Standard normal variate (at 5% type 1 error) = 1.96

p = expected proportion of respondents (15%)

d = absolute error = 0.05 $Sample\ Size = \frac{1.96^2 \times 0.15(1-0.15)}{0.05^2} = 195.92$

For the study, the sample size was rounded up to 200. The study was carried out at the University of Port Harcourt, Port Harcourt, Nigeria. Among students within the designated age interval of 18-26 years.

2.2 Study Criteria

Only respondents who have declared their consent to participate in the study and of age (currently undergoing or understand what dysmenorrhea is all about) was allowed to participate in the study while respondent yet to experience menarche (less than designated age) were excluded from the study.

2.3 Method of Data Collection

For this study, data were collected using a descriptive questionnaire where it was divided

into sections A, B, C, and D; Section, A set to obtain the socio-demography of the respondents, section B; knowledge of dysmenorrhea, section C; possible symptoms experienced, and Section D; managerial procedure taken. All respondents declared their consent to participate, questionnaire was administered and retrieved.

2.4 Statistical Analysis

Data were retrieved from every respondent and entered into a Microsoft spreadsheet (excel package) and International Business Machine of Statistical Package for Social Science (IBM SPSS version 26). Results were presented as frequency and percentage and chi-square was used as an inferential statistic and probability less than 0.05 ($p < 0.05$) was considered statistically significant.

3. RESULTS

The demographic characteristics of the participants are presented in Figs. 1, 2, and 3 which show the age range, marital status, and level of the students. In the Table 1, the nulliparous were 96.5% while multiparous were 3.5%. Those who experienced dysmenorrhea were 81.5% while 18.5% of the total population does not experience dysmenorrhea during menstruation, 74.5% claim that dysmenorrhea abounds in their families. The majority of the respondents observed their menstrual flow within 4-5 days (72.5%), and it was followed by 2-3 days (13%), 6-7 days (12%) and only 2.5% observed it after 7 days. The study evaluated different categories of pain felt during




Fig. 1. Age range

menstruation, 37% of the total population had severe pain, 30% mild and 33% showed moderate pain though over 92% felt the pain within their waist/hip region. 83.5% of the total population experience dysmenorrhea during their menstruation and only 16.5% claim to experience dysmenorrhea a week or days before the menstrual period.

Table 2, shows the symptoms of dysmenorrhea and it was observed that anxiety 55%, mood swings 92%, tender breasts 70.5%, food craving 63.5%, fatigue 83.5%, irritating mood 84.5%, depression 40%, dizziness 54.5%, vomiting 15.5%, diarrhea 59.5%, and headache 50% of the total population respectively.

Table 3 illustrates the management of dysmenorrhea and how they manage dysmenorrhea and it shows that 23.5% of the population uses synthetic drugs and 1.5% uses local and herbal medical practices. Of respondents who manage dysmenorrhea using local and herbal medical practices, 37% claim they manage it by drinking warm water, 5% claim they do exercise, 8% manage dysmenorrhea by reducing sugary substance while those that use synthetic/therapeutic, 19% uses paracetamol, 8% uses felvin, 9.5% uses buscopan and 4% uses ibuprofen.




Fig. 2. Marital status




Fig. 3. Level of student

Table 1. Knowledge and prevalence of dysmenorrhea

Variable	Nulliparous		Multiparous		X ²	p-value	Inference
Do you have kids of your own	193 (96.5%)		7 (3.5%)		172.98	0.00	S
	Yes (%)		No (%)				
Do have pain during menstruation	163(81.5%)		37 (18.5%)		79.38	0.00	S
Do any family members also notice menstrual pain	149 (74.5%)		51 (25.5%)		48.02	0.00	S
	2-3 days	4-5 days	6-7 days	Above 7 days			
What is the duration of the menstrual period	26 (13%)	145 (72.5%)	24 (12%)	5 (2.5%)	246.04	0.00	S
	Severe	Mild		Moderate			
How can you categorize the pain during the menstrual period	74 (37%)	60 (30%)		66 (33%)	63.52	0.00	S
	Waist/Hip	Legs		Back			
Where do you feel the pain	184 (92%)	6 (3%)		10 (5%)	309.88	0.00	S
	DMP		W/DtMP				
When does the pain normally occur	167 (83.5%)		33 (16.5%)		89.78	0.00	S

X²= chi-square, S=significant, DMP=During menstrual period, W/DtMP= weeks/days to menstrual period,

Table 2. Symptoms of dysmenorrheal

	Yes	No	X ²	p-value	inference
Anxiety	110 (55%)	90 (45%)	2.00	0.15	NS
Mood swings	184 (92%)	16 (8%)	141.12	0.00	S
Tender breast	141 (70.5%)	59 (29.5%)	33.62	0.00	S
Food craving	127 (63.5%)	73 (36.5%)	14.58	0.00	S
Fatigue	167 (83.5%)	33 (16.5%)	89.78	0.00	S
Irritating mood	169 (84.5%)	31 (15.5%)	95.22	0.00	S
Depression	80 (40%)	120 (60%)	8.00	0.00	S
dizziness	109 (54.5%)	91 (45.5%)	1.62	0.20	NS
Vomiting	31 (15.5%)	169 (84.5%)	95.22	0.00	S
Diarrhea	119 (59.5%)	81 (40.5%)	7.22	0.00	S
Headache	100 (50%)	100 (50%)	0.00	1.00	S

X²= chi-square S=significant, NS= non-significant, (p<0.05)

Table 3. Management of Dysmenorrhea

Variables		None	SD	L/hMP	X²	p-value	inference
How do you manage the pain		150 (75%)	47 (23.5%)	3 (1.5%)	170.77	0.00	S
What L/hMp do use in the management of pain	None	95 (47.5%)			172.05	0.00	S
	Drinking warm water	74 (37%)					
	Physical exercise	10 (5%)					
	Reduce sugary substance	16 (8%)					
	Taking herbs/bitter substance	5 (2.5%)					
what synthetic/therapeutic practices do you use?	None	119 (59.5%)			207.15	0.00	S
	Paracetamol	38 (19%)					
	Felvin	16 (8%)					
	Bosco pan	19 (9.5%)					
	Ibuprofen	8 (4%)					

X²= chi-square *S*=significant, ($p < 0.05$) *SD*= synthetic drug, *L/hMP*=local/herbal medical practices

4. DISCUSSION

Dysmenorrhea, or painful menstrual cramps, is a prevalent problem among female university students. Although the prevalence varies, research reveals that a considerable number of female students suffer from dysmenorrhea during their menstrual periods. In this study, the frequency of dysmenorrhea was 81.5% among the respondents. Studies conducted among Northern Ghana university students by Ameade et al., [21], reported similar prevalence rates of 83.6%, Ezebialu et al., [22] also reported that the prevalence of dysmenorrhea was 82.2% among Nigerian undergraduate students. In Nigeria prevalence of dysmenorrhea has been reported at 78.1% in Ogun State [7], 62% in Oyo State [8], 58% in Osun State [9], 60-64% in Ile-Ife [10,11], and 69.8% in Maiduguri [12]. These studies have agreed with our study that dysmenorrhea is prevalent among women in their reproductive years though studies have also been conducted aside Nigeria, and it showed that 85.6% in Kuwait [13], 79.4% in Saudi Arabia [14], 84.2% in India [15], 74.8% in Spanish [16] 89.4% in Al-Asadi, [23] 84.2% in Lithuania [24], 88% in Australia [25], 85.1% in Palestine [26], 83.6% in Northern Ghana [27] and 84.1% in Italy [28], 68.7% in China [29], 74.8% in Spain [16], 94% in Oman [30], 98.4% in Iran [31] and 95% in Turkey [32], 80.9% in Lebanon [17] and 92% in Riyadh [18]. Our findings agree with the findings of these studies that dysmenorrhea abounds in women in their reproductive years.

The present study revealed that the prevalence of dysmenorrhea was higher among female students who had a family history of dysmenorrhea (74.5%). Similar results were disclosed in a study that reported inconsistent with 65.6% and 60.7% of the students had a family history of dysmenorrhea by El-Mawgod et al. [33] and Ali et al. [14] respectively. Though our study claims that 30% experienced mild pain, 33% had moderate pain, and 37% had severe pain. Our findings agree with Ali et al. [14], in their report, they showed that the majority of the study population experiences moderate pain followed by severe pain the students suffered from mild, moderate, and severe pain, respectively. Alsaleem, [34] also disagrees with our findings that only a few had severe pain. Even a study by Fernández-Martínez et al. [16], further agrees with the present study that moderate pain and severe pain are experienced more by women. This indicated that dysmenorrhea can be considered one of the

most common health problems that may affect the health, psychological status, and daily activities of female students at the university level.

Women who experienced dysmenorrhea during menstruation were accounted for in the present study (83.5%) and 16.5% experienced dysmenorrhea week/day to menstrual periods. Our findings agree with a similar result by Ali et al., [14], it was observed that some women highly experienced pain in their waist/hip at the rate of 92%, leg (3%), and back (5%), and this had a similar result with Al-Matouq et al., [13]. This indicates that most students may be incapacitated in one way or the other as a result of dysmenorrhea. To confirm this, the majority reported that dysmenorrhea has limitations on their activities. This agrees with the studies of Burnett et al. [35], Armour et al. [36], and Tu and Hellman, [37] that posited that symptoms of dysmenorrhea have a significant impact on the quality of life and limitation on activities which shows in case of this study that mood swings have the highest rate of 92%. 55% of anxiety, 70.5% of tender breasts, 63.5% of food cravings, 83.5% of fatigue, 84.5% of irritating mood, 40% of depression, 54.5% of dizziness, 15.5% of vomiting, 59.5% of diarrhea and 50% of headache and agreed with the Aktas [38] and Fernández-Martínez et al. [16].

The present study evaluates the different methods of managing dysmenorrhea, it was observed in this study that 37% of the population drink warm water, 5% manage dysmenorrhea through physical exercise, 8% reduce sugary substances and 2.5% takes herbs and bitter substance. Omidvar et al., [39] agree with the present study on the various ways of dysmenorrhea management. Of women that use synthetic/therapeutic, 19% use paracetamol, 8% use felvin, 9.5% of buscopan, and 4% of Ibuprofen. It shows a significant difference. The findings agree with Ali et al. [14] and Omidvar et al., [39].

The above-discussed study has shown some similarities and differences in the prevalence and management of dysmenorrhea among university students. The difference could be attributed to race, lifestyle and methodology.

5. CONCLUSION

The prevalence of dysmenorrhea is higher among the university students of Port Harcourt,

Nigeria and the symptoms range from mood swings, fatigue, irritating mood, tender breasts, food cravings, dizziness, diarrhea, and headache. The majority of the respondents use local/herbal medication to manage dysmenorrhea rather than taking synthetic drugs.

CONSENT

As per international standards or university standards, respondents' written consent has been collected and preserved by the authors

ETHICAL CONSIDERATION

This study was approved by the faculty research and ethical committee of Basic Medical Sciences, University of Port Harcourt, Nigeria.

ACKNOWLEDGEMENT

All authors wish to thank the study respondents for consenting to be part of the successful completion of this research and also acknowledge Dr. Oghenefego Michael Adheke, and Ikechukwu Williams Osuchukwu for their assistance in bringing the study to completion.

COMPETING INTERESTS

The authors have declared that no competing interests exist.

REFERENCES

1. Mao L, Xi S, Bai W, Yao C, Zhou Y, Chen X, Sun Y. Menstrual patterns and disorders among Chinese women of reproductive age: a cross-sectional study based on mobile application data. *Medicine*. 2021;100(16).
2. Seifalian A, Kenyon J, Khullar V. Dysmenorrhoea: Can Medicinal Cannabis Bring New Hope for a Collective Group of Women Suffering in Pain, Globally? *International Journal of Molecular Sciences*. 2022;23(24):16201.
3. Aolymat I, Khasawneh AI, Al-Tamimi M. COVID-19-associated mental health impact on menstrual function aspects: dysmenorrhea and premenstrual syndrome, and genitourinary tract health: a cross-sectional study among Jordanian medical students. *International journal of environmental research and public health*. 2022 Jan 27;19(3):1439.
4. Bolkar PE, Shelke PA, Sanap GS. Effect of medicinal herbs on dysmenorrhoea. *Journal of Pharmacognosy and Phytochemistry*. 2023;12(2):150-9.
5. Mizuta R, Maeda N, Tashiro T, Suzuki Y, Oda S, Komiya M, Urabe Y. Quality of life by dysmenorrhea severity in young and adult Japanese females: A web-based cross-sectional study. *Plos one*. 2023 Mar 16;18(3):e0283130.
6. Thakur P, Pathania AR. Relief of dysmenorrhea—A review of different types of pharmacological and non-pharmacological treatments. *Materials Today: Proceedings*. 2022 Jan 1;48:1157-62.
7. Farotimi AA, Esike J, Nwozichi CU, Ojediran TD, Ojewole FO. Knowledge, attitude, and healthcare-seeking behavior towards dysmenorrhea among female students of a private university in Ogun State, Nigeria. *Journal of basic and clinical reproductive sciences*. 2015;4(1):33-8
8. Busari AO. Menstrual knowledge and health care behavior among adolescent girls in rural Nigeria. *Int J Appl Sci Technol* 2012;2:149-54.
9. Ogunfowokan AA, Babatunde OA. Management of primary dysmenorrhea by school adolescents in Ile-Ife, Nigeria. *J School Nurs* 2010;26:131-6.
10. Titilayo A, Agunbiade OM, Banjo O, Lawani A. Menstrual discomfort and its influence on daily academic activities and psychosocial relationship among undergraduate females in Nigeria. *Tanzania J Health Res* 2009;11:181-8.
11. Esimai O, Esan GO. Awareness of menstrual abnormality amongst college students in urban area of Ile-Ife, Osun state, Nigeria. *Indian J Community Med* 2010;35:63-6.
12. Amaza DS, Sambo N, Ziraha JV, Dalori MB, Japhet H, Toyin H. Menstrual pattern among female medical students in University of Maiduguri Nigeria. *Br J Med Medical Res* 2012;2:327-37.
13. Al-Matouq S, Al-Mutairi H, Al-Mutairi O, Abdulaziz F, Al-Basri D, Al-Enzi M, Al-Taiar A. Dysmenorrhea among high-school students and its associated factors in Kuwait. *BMC pediatrics*. 2019 Dec;19:1-2.
14. Ali A, Ali A, Alotaibi NS, Alsufyani MS, Alotaibi AJ, Almutairi MM, Eldalo AS. Prevalence, impact, and management perception of dysmenorrhea among university students: A cross-sectional

- study. *Brazilian Journal of Pharmaceutical Sciences*. 2022 Dec 2;58.
15. Vilšinskaitė DS, Vaidokaitė G, Mačys Ž, Bumbulienė Ž. the risk factors of dysmenorrhea in young women czynniki ryzyka bolesnego miesiączkowania u młodych kobiet. *Wiad Lek*. 2019;72(6):1170-7.
 16. Fernández-Martínez E, Onieva-Zafra MD, Parra-Fernández ML. Lifestyle and prevalence of dysmenorrhea among Spanish female university students. *PLoS one*. 2018 Aug 10;13(8):e0201894.
 17. Karout S, Soubra L, Rahme D, Karout L, Khojah HM, Itani R. Prevalence, risk factors, and management practices of primary dysmenorrhea among young females. *BMC women's health*. 2021 Dec;21:1-4
 18. Dahlawi H, Bukhari I, Alshammari F, Althaqib G, Alhawasawi M, Bashiri R, Alruweili R. Effect of dysmenorrhea on the academic performance among students studying in princess nourah bint abdulrahman university, Riyadh. *International Journal of Medicine in Developing Countries*. 2021 Feb 11;5(2):588
 19. Adheke OM, Asiwe N, Uahomo PO, Ezerioha CE, Wariboko LI, Irozulike FC, Opara AI. (2023) Assessment of Determinants for Choosing Medical Sciences as a Career Path amongst Medical Students of University of Port-Harcourt, Nigeria. *Journal of Complementary and Alternative Medical Research*. 3;22(2):30-6.
 20. Asiwe N, Irozulike FC, Wariboko LI and Adheke OM, (2023) Health Assessment of the Ikwerres and Okrikas Ethnic Group of Rivers State, Nigeria: Using Body Mass Index and Waist-to-Hip Ratio, *Journal of Complementary and Alternative Medical Research*. 31;23(2):27-35.
 21. Ameade EP, Amalba A, Mohammed BS. Prevalence of dysmenorrhea among university students in Northern Ghana; its impact and management strategies. *BMC Women's Health*. 2018 Dec;18(1):1-9.
 22. Ezebialu IU, Ezenyeaku CC, Umeobika JC. Prevalence of dysmenorrhea and its contribution to school absenteeism among Nigerian undergraduate students. *Annals of Health Research*. 2021 Mar 3;7(1):59-66.
 23. Al Asadi J, Abdul Qadir R. Dysmenorrhea and its impact on daily activities among secondary school students in Basra, Iraq. *J Fac Med Baghdad*. 2013;55(4):339-44.
 24. Vilšinskaitė DS, Vaidokaitė G, Mačys Ž, Bumbulienė Ž. THE RISK FACTORS OF DYSMENORRHEA IN YOUNG WOMEN CZYNNIKI RYZYKA BOLESNEGO MIESIĄCZKOWANIA U MŁODYCH KOBIEĆ. *Wiad Lek*. 2019;72(6):1170-7.
 25. Abdel-Salam DM, Alnuman RW, Alrwuaili RM, Alrwuaili GA, Alrwuaili EM. Epidemiological aspects of dysmenorrhea among female students at Jouf University, Saudi Arabia. *Middle East Fertility Society Journal*. 2018 Dec 1;23(4):435-9.
 26. Abu Helwa, H. A., Mitaeb, A. A., Al-Hamshri, S., & Sweileh, W. M. (2018). Prevalence of dysmenorrhea and predictors of its pain intensity among Palestinian female university students. *BMC women's health*, 18(1), 1-11.
 27. Acheampong K, Baffour-Awuah D, Ganu D, Appiah S, Pan X, Kaminga A, Liu A. Prevalence and predictors of dysmenorrhea, its effect, and coping mechanisms among adolescents in Shai Osudoku District, Ghana. *Obstetrics and Gynecology International*. 2019 May 20;2019.
 28. Grandi G, Ferrari S, Xholli A, Cannoletta M, Palma F, Romani C, Volpe A, Cagnacci A. Prevalence of menstrual pain in young women: what is dysmenorrhea? *Journal of pain research*. 2012 Jun 20;169-74.
 29. Wong CL. Health-related quality of life among Chinese adolescent girls with Dysmenorrhoea. *Reproductive Health*. 2018 Dec;15(1):1-0.
 30. Oni TH, Tshitangano TG. Prevalence of menstrual disorders and its academic impact amongst Tshivenda speaking teenagers in rural South Africa. *Journal of Human Ecology*. 2015 Jul 1;51(1-2):214-9.
 31. Ghaderi F, Asghari Jafarabadi M, Mohseni Bandpei MA. Dysmenorrhea and self-care strategies in Iranian female students: a regression modeling of pain severity and underlying factors. *International Journal of Adolescent Medicine and Health*. 2016 Jul 18;29(6):20160017.
 32. Duman NB, Yıldırım F, Vural G. Risk factors for primary dysmenorrhea and the effect of complementary and alternative treatment methods: Sample from Corum, Turkey. *International journal of health sciences*. 2022 May;16(3):35.
 33. Abd El-Mawgod MM, Alshaibany AS, Al-Anazi AM. Epidemiology of dysmenorrhea

- among secondary-school students in Northern Saudi Arabia. *Journal of the Egyptian Public Health Association*. 2016 Sep 1;91(3):115-9.
34. Alsaleem MA. Dysmenorrhea, associated symptoms, and management among students at King Khalid University, Saudi Arabia: An exploratory study. *Journal of family medicine and primary care*. 2018 Jul;7(4):769.
35. Burnett MA, Antao V, Black A, Feldman K, Grenville A, Lea R, Lefebvre G, Pinsonneault O, Robert M. Prevalence of primary dysmenorrhea in Canada. *Journal of Obstetrics and Gynaecology Canada*. 2005 Aug 1;27(8):765-70.
36. Armour M, Hyman MS, Al-Dabbas M, Parry K, Ferfolja T, Curry C, MacMillan F, Smith CA, Holmes K. Menstrual health literacy and management strategies in young women in Australia: a National Online Survey of young women aged 13-25 Years. *Journal of Pediatric and Adolescent Gynecology*. 2021 Apr 1;34(2):135-43.
37. Tu F, Hellman K. Primary dysmenorrhea: diagnosis and therapy. *Obstetrics & Gynecology*. 2021 Apr 1;137(4):752.
38. Aktaş D. Prevalence and factors affecting dysmenorrhea in female university students: effect on general comfort level. *Pain Management Nursing*. 2015 Aug 1;16(4):534-43.
39. Omidvar S, Bakouei F, Amiri FN, Begum K. Primary dysmenorrhea and menstrual symptoms in Indian female students: prevalence, impact and management. *Global journal of health science*. 2016 Aug;8(8):135.

© 2023 Irozulike et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/107877>