



# **Role of Foliar Application of Micronutrients on Growth and Yield of Pulses: A Review**

**Shruti Pandey<sup>a++</sup>, Pawan Kumar Sharma<sup>b†</sup>, Shreya Roy<sup>c†</sup>,  
Dilip Choudhary<sup>d#</sup>, Rajendra Choudhary<sup>e†</sup>,  
Indar Raj Naga<sup>f#\*</sup>, Ravin Singh<sup>g#</sup> and Jigyasa Ninama<sup>h#</sup>**

<sup>a</sup> Department of Agronomy, SAGE, University, Indore, India.

<sup>b</sup> Department of Agrometeorology, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj, India.

<sup>c</sup> Department of Agronomy, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj, India.

<sup>d</sup> Department of Agronomy, Swami Keshwanand Rajasthan Agriculture University, Bikaner, India.

<sup>e</sup> Department of Agronomy, College of Agriculture, Naik Marathwada Krishi Vidyapeeth, Prabhani, Maharashtra, India.

<sup>f</sup> Department of Soil Science and Agricultural Chemistry, Jawaharlal Nehru Krishi Vishwa Vidyalaya Jabalpur, India.

<sup>g</sup> Department of Agronomy, Shri Khushal Das University, Hanumangarh, Rajasthan, India.

<sup>h</sup> Department of Agronomy, Swami Keshwanand Rajasthan Agriculture University, Bikaner, India.

## **Authors' contributions**

*This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.*

## **Article Information**

DOI: 10.9734/IJECC/2024/v14i13838

## **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/111562>

**Review Article**

**Received: 05/11/2023**

**Accepted: 11/01/2024**

**Published: 13/01/2024**

<sup>++</sup> Assistant Professor;

<sup>†</sup> M.Sc. Scholar;

<sup>#</sup> Ph.D. Scholar;

<sup>\*</sup>Corresponding author: E-mail: indrarajnaga0@gmail.com;

## ABSTRACT

Pulses respond positively to micronutrients application, particularly during the reproductive stages, their application is still uncommon. The current review suggest that foliar application of Zn, Fe or Mo had a favourable impact on growth indicators and grain yield. protein content, total and active nodule counts and yield components plant<sup>-1</sup>. Molybdenum is essential for the development of nodule tissue and an increase in N<sub>2</sub> fixation, both of which are necessary for Rhizobium to increase the nitrogen fixation process.

*Keywords: Iron; molybdenum; pulses; yield; zinc.*

## 1. INTRODUCTION

“Pulses are the main source of protein particularly for vegetarians and contribute about 14 per cent of the total protein of an Indian average diet” [1]. “The production and consumption of pulses, sometimes referred to as food legumes, is second only to cereals in India. The word pulse is derived from the Latin word Puls means pottage i.e., seed boiled to make porridge or thick soup. Pulse is a rich source of dietary protein (25%), energy, minerals and vitamins for the vegetarian population. As a vital source of both human and animal food and animal feed, it is also crucial for maintaining soil fertility” [2].

In addition to crop productivity, micronutrients also improve the amount of crude protein, amino acids, total lipids, energy value, and other nutrients in chickpea, soybean, black gramme, etc. Foliar fertilisation is the most efficient way to raise output, and research has shown that it can increase yield by 12 to 25% when compared to conventional fertilisation [3].

“Iron is an essential key enzymes Co-factor which performs different function in breakdown of hormone and also take part in many chemical reactions takes place in plants” [4]. Iron is required for the formation of chlorophyll even though it is not a component of it. As a result, a lack of chlorophyll causes iron deficiency and chlorotic plants [5].

“Zinc has been the micronutrient that crops, especially pulses, have required in sufficient amounts” [6]. “Zinc in plants required for biosynthesis of hormone” [7]. Zinc is essential for metabolism and participates in the production of nodules, which fix nitrogen [8]. Zinc is a crucial component of RNA polymers and gives ribosomes their structural integrity [6]. Since Zn is required for the synthesis of tryptophan [9,10],

Zinc is required for pollen function and fertilization [11].

“The availability of Molybdenum in soil has been shown to enhance the production of leguminous crops” [12]. Molybdenum, which is essential for several metabolic activities and regulates ascorbate-glutathione in plants. Plants deficient in molybdenum have lower quantities of chlorophyll and abnormal changes to their chloroplasts [13]. The synthesis of the nitrate reductase enzyme and the symbiotic nitrogen fixation in legumes depend on the crucial micronutrient molybdenum [14,15]. “Molybdenum increases plant height, number of branches and pods plant<sup>-1</sup>, number of seeds plant<sup>-1</sup> and nodule production in legume crops. It also enhances yield quality” [16].

## 2. EFFECT ON PLANT HEIGHT

Alam et al. [17] reported In an earlier trial, the longest plant was produced with a 2.5 kg Zn ha<sup>-1</sup> application. The height of the plants was strongly impacted by the application of 10 kg Zn ha<sup>-1</sup> (47.76 cm and 47.88 cm) in summer mungbean [18]. El Sayed Hameda et al. [19] reported that the By applying a combination of microelements to the leaves of pea plants, the yield components indicated by plant height were significantly increased. Zinc in auxin mixture [20]. The plant height of the pigeonpea crop was increased by applying 5 mg kg<sup>-1</sup> Zn added to the soil [21]. Some enzymes with significant roles in cell division and lengthening were activated by zinc, and as a result, the stem height increased [22]. Habbasha et al. [23] observed 0.2% ZnSO<sub>4</sub> application to chickpea at seed filling stage result in significantly higher plant height (81.43 cm). Zn and Fe may have positive benefits when given topically because of their importance in respiration, photosynthesis, and other biochemical and physiological processes as well as their function in increasing yields [24,25]. “The

increment in plant height of cowpea with zinc spray ( $1 \text{ g L}^{-1}$ ) has been documented” by Salehin and Rahman [26]. The application of molybdenum to the leaves has a considerable impact on plant parameters, including plant height and biological performance. The result was in conformity with those of Manjili et al. [27]. Hazra and Tripathi [28] observed that “molybdenum application at the rate of  $1.5 \text{ kg/ha}$  increased forage in calcareous soil”. Sahu et al. [29] reported that “the application of  $\text{FeSO}_4$  at  $2 \text{ kg ha}^{-1}$  significantly increased the growth characters over control in chickpea”. Thapu et al. [30] concluded that “the application of micronutrients like Fe (as ferrous sulphate at 0.4%), Mn, Cu and Zn significantly increased the growth attributes in pea”.

### 3. EFFECT ON NUMBER OF NODULES PLANT<sup>-1</sup>

Increased infection and rhizobial colonisation in the rhizosphere may be the source of the increased nodule count due to the increased availability of micronutrients [31]. Chauhan et al. [32] reported “91% higher nodulation in soybean under the application of Zn @  $5 \text{ kg ha}^{-1}$  in consecutive years of study”. Ram and Kattiyar [18] found that “the application of Zn @  $10 \text{ kg ha}^{-1}$  to mungbean significantly increase the nodulation”. Badar et al. [33] “Molybdenum is essential for the development of nodule tissue and an increase in  $\text{N}_2$  fixation, both of which are necessary for Rhizobium to increase the nitrogen fixation process. Every bacterium that fixes nitrogen requires molybdenum since it is an essential component of the nitrogenase enzyme reported by Hirpara et al. [34]. Iron and molybdenum significantly enhanced nodules number in legumes crops [35]. An agreement with Tantawy et al. [36] reported that “Fe and Mo significantly increased number of active nodules in broad bean”.

### 4. EFFECT ON DRY MATTER ACCUMULATION

The production of straw was enhanced by applying multiple foliar sprays of various fertiliser mixtures at distinct growth stages. As previously documented, foliar treatment of macro and micronutrients boosted biological yield by Piri Issa et al. [37]. “The improvement in nodulation might have resulted in a higher amount of nitrogen fixation and there by better vegetative growth and dry matter production” Balla et al. [38].

### 5. EFFECT ON PROTEIN CONTENT

Zinc treatment to the leaves resulted in higher protein content as a result of improved physiological properties and protease enzyme activity Nandan et al. [39], Usman et al. [40] and Lokhande et al. [41]. Singh and Singh [42] also reported 20.2% higher protein content in chickpea grain under the application of  $10 \text{ kg Zn ha}^{-1}$ . Khrogamy and Farnia [43] on zinc application in chickpea revealed the fact that, the application of  $\text{ZnSO}_4$  @  $20 \text{ kg ha}^{-1}$  improved the protein content (24.02%) in chickpea. Imsande [44] stated that iron had an important role in the synthesis of chlorophyll and also help in the absorption of other nutrients. Oguz [45] that “increasing Mo application up to  $6 \text{ g kg}^{-1}$  considerably affected the protein content in chickpea”. Roy et al. [46] reported that “the combined application of  $5.5 \text{ kg Zn ha}^{-1} + 0.1\% \text{ Zn}$  as foliar spray increased the seed crude protein by 26.9% over control”. Sharma et al. [47] reported that the treatment of chelated Fe ( $1$  or  $2 \text{ kg ha}^{-1}$ ) considerably enhanced the protein content in seed as well as all yield-contributing attributes in the pigeon pea crop.

### 6. EFFECT ON YIELD ATTRIBUTES

The beneficial effects of foliar nutrition of micro nutrients on mung bean were also reported by several researchers Kumawat et al. [48]. The micronutrients might have enhanced role in seed setting that resulted in improvement in no. of seeds pod<sup>-1</sup>.

### 7. EFFECT ON NUMBER OF PODS PLANT<sup>-1</sup>

Kumar et al. [49] reported that “seed inoculation with zinc @  $5 \text{ kg ha}^{-1}$  gave the significantly higher growth and yield attributes”. Togay et al. [16] reported that molybdenum increased pods plant<sup>-1</sup>. Valenciano et al. [50] showed pod number plant<sup>-1</sup> were greater with Zn application. Ram and Kattiyar [18] found that “the application of  $10 \text{ kg Zn ha}^{-1}$  to mungbean resulted in significant improvement in the pod and seed number”. In keeping with our findings, it was reported that micronutrient foliar spraying increased the number of pods plant<sup>-1</sup> in lentil [51]. Shah et al. [52] concluded that “the number of seeds pod<sup>-1</sup> and number of pods plant<sup>-1</sup> in pigeonpea significantly increase under the application of Zn @  $20 \text{ kg ha}^{-1}$ ”. Increase of pods plant<sup>-1</sup> using Zn spraying reported by Teixeira et al. [53] in common bean. A decrease in flower

shedding, a rise in pod formation, and the improvement of photosynthetic material distribution to these organs could all boost the pod plant<sup>-1</sup> [54].

## 8. EFFECT ON NUMBER OF SEEDS POD<sup>-1</sup>

Rizk and Abdo [55] reported that yield components showed a highly significant increase with the application of Zn (0.2 or 0.4 g/l) compared to the control. The reports of other researchers that indicated the foliar application of Zn increased seeds pod<sup>-1</sup> in common bean [53,22,56] and Faba bean [57]. Sarkar and Banik [58] observed the molybdenum application significantly increased seeds pod<sup>-1</sup>.

## 9. SEED YIELD

Zn application at the rate of 0.3% in combination with Fe increased seed yield and lead to the highest amounts of seed yield. Khrogamy and Farnia [43] It was noted that seed output and total biomass in chickpea cultivars were significantly affected by the treatment of ZnSO<sub>4</sub> @ 20 kg ha<sup>-1</sup>. Ramakrishna et al. [59] studied “the effects of improved production technologies involving application of 10 kg Zn ha<sup>-1</sup> and found increased pigeonpea yield”. Patil et al. [60] The application of ZnSO<sub>4</sub> @ 10 kg ha<sup>-1</sup> yielded considerably greater grain (29.55 q ha<sup>-1</sup>), stalk (95.81 t ha<sup>-1</sup>), and bhusa yields (20.58 q ha<sup>-1</sup>) when four different amounts of zinc (0, 10 and 20 kg ha<sup>-1</sup>) were investigated to see how they affected pigeonpea growth and yield. Ibrahim and Ramadan [61] indicated Zn spraying increased seed yield and growth of dry bean. Malik et al. [62] When the impact of zinc on mungbean growth and yield was examined, it was discovered that the treatment of 20 ppm zinc resulted in the highest seed yield plant<sup>-1</sup> (78.20 g), compared to the control. Shah et al. [52] demonstrated that the treatment with Zn @ 20 kg ha<sup>-1</sup> had the highest pigeonpea yield. We discovered that the increase in bean seed yield brought on by Fe and Zn spraying was comparable to the others [50,63,56] soybean [64] and Faba bean [57]. “The grain yield of chickpea was significantly increased with the application of iron and molybdenum” [65,66]. “Zinc fertilization had positive effect on the yield of pulses. Application of Zn @ 5 kg ha<sup>-1</sup> along with S @ 60 kg ha<sup>-1</sup> recorded 67% higher seed yield over control in mung bean” [67] Application of FeSO<sub>4</sub> @ 2 kg ha<sup>-1</sup> along with bio-fertilizer inoculation in chickpea led to higher grain and straw yields over control [29]. Thapu et al. [30] observed that

“the application of micronutrients like Fe (as FeSO<sub>4</sub> at 0.4%), Mn, Cu and Zn significantly increased the grain yield in pea”. Salam et al. [68] conducted “a field experiment at Raipur, Chhatisgarh and concluded that the seed yield of Urd bean under application of FeSO<sub>4</sub> at 2-20 kg Fe ha<sup>-1</sup> was maximum over control”. Kumawat et al. [69] observed that “the soil application of Fe at 25 kg FeSO<sub>4</sub> ha<sup>-1</sup> recorded significantly higher seed and straw yield of summer mung bean”. Kumar et al. [70] conducted “an experiment at Kanpur and results revealed that the application of 10 kg Fe ha<sup>-1</sup> increase the grain yield of chickpea by 17.3% over control”.

## 10. CONCLUSION

It is concluded that Molybdenum is essential for the development of nodule tissue and an increase in N<sub>2</sub> fixation, both of which are necessary for Rhizobium to increase the nitrogen fixation process.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Meena KK, Meena RS, Kumawat SM. Effect of sulphur and iron fertilization on yield attributes, yield and nutrient uptake of mungbean (*Vigna radiata*). Indian Journal of Agricultural Sciences 2013;83(4):472
2. Kavya P, Singh S, Hinduja N, Tiwari D, Sruthi S. Effect of foliar application of micronutrients on growth and yield of green gram (*Vigna radiata* L.) Legume Research - An International Journal 2023; 44(12):1460-1464.
3. Deshmukh MS, Zade SP, Gourkhede PH, Inamke PS. Effect of high-p-chelated micronutrient foliar spray on yield and quality of Chickpea. International Journal of Tropical Agriculture, 2017;35(3):707-712.
4. Kerkeb L, Connolly E. Iron transport and metabolism in plants. J. Gent. Eng. 2006; 27:119-140.
5. Katyal JC, Randhawa NS. Micronutrient. F.A.O. Fertilizer and plant nutrition. Bulletin No. 7. United Nation; 1983.
6. Yashona DS, Mishra US, Aher SB. Response of pulse crop to sole and combined mode of zinc application.

- Journal of Soils and Crops. 2018;28(2): 249–258
7. Kudi S, Swaroop N, David AA, Thomas T, Hasan A, Rao S. Effect of different levels of sulphur and zinc on soil health and yield of Greengram (*Vigna radiata* L.) Var. Patidar-111. *Journal of Pharmacognosy and Phytochemistry*. 2018;7(3):2271–2274.
  8. Patel MM, Patel IC, Patel RI, Acharya S. Effect of Zinc and Iron on yield and yield attributes of rainfed cowpea (*Vigna unguiculata* L. Walp). *Annals of Arid Zone*. 2011;50(1):17–19.
  9. Brown PH, Cakmak I, Zhang Q. Form and function of zinc in plants. Chap. 7, In A.D. Robson (Ed). *Zinc in Soils and Plants*, Kluwer Academic Publishers, Dordrecht. 1993;90-106.
  10. Alloway BJ. *Zinc in Soils and Crop Nutrition*. Publ. of International Zinc Association; 2004. Available:[http://www.iza.com/Documents/Communications/Publications/ALLOWAY\\_PRINT.pdf](http://www.iza.com/Documents/Communications/Publications/ALLOWAY_PRINT.pdf)
  11. Ali A, Ahmad B, Hussain I, Ali A, Shah FA. Effect of phosphorus and zinc on yield of lentil. *Pure and Applied Biology*. 2017;6(4): 1397–1402.
  12. Chandra G, Gambhir L, Upadhyay R. Effects of biofertilizer with and without molybdenum on growth and seed yield of Chickpea under Doon Valley of Uttarakhand. *Current Journal of Applied Science and Technology*. 2020;39(15): 133–139.
  13. Nasar J, Shah Z. Effect of iron and molybdenum on yield and nodulation of lentil. *ARPJ: Journal of Agriculture and Biological Sciences*. 2017;12(11): M332–339
  14. Williams RJP, Frausto da Silva JJR. The involvement of molybdenum in life. *Biochemical and Biophysical Research Communications*. 2002;292:293–299
  15. Roy RN, Finck A, Blair GJ, Tandon HLS. *Plant nutrition for food security. A guide for integrated nutrient management*. FAO Fertilizer and Plant Nutrition Bulletin 16. FAO, Rome, Italy. 2006;368.
  16. Togay Y, Togay N, Dogan Y. Research on the effect of phosphorus and molybdenum applications on the yield and yield parameters in lentil (*Lens culinaris* Medic.). *African Journal of Biotechnology*. 2008; 7(9):1256–1260.
  17. Alam MK, Uddin MM, Ahmed M, Latif MA, Rahman MM. Growth and green pod yield of garden pea varieties under different nutrient levels. *Journal of Agroforestry and Environment*. 2010;4(1):105–107.
  18. Ram S, Katiyar TPS. Effect of sulphur and zinc on the seed yield and protein content of summer mungbean under arid climate. *Int. J. Sci. Nature*. 2013;4(3):563-566.
  19. El Sayed Hameda EA, Amen El-Sh A, El-Morsy AH, Tolba MH. Effects of foliar spraying with microelements and different fertilizer sources on quality and yield of *Pisum sativum* L. plant. *International Research Journal of Agricultural Science and Soil Science*. 2012;21(1):17–24.
  20. Basavarajeswari. Expression profiling of zinc transporter genes in tomato grown under different concentrations of zinc. *International Journal of Microbiology Research*. 2018;10(6):1252-1255.
  21. Dube BK, Sharma CP, Chatterjee C. Response of Pigeonpea to Applied Zinc in Ustifluent Soils of Western Uttar Pradesh. *J. Indian Soc. Soil Sci*. 2001;49(3):471-475.
  22. Seifi Nadergholi M, Yarnia M, Rahimzade Khoei F. Effect of zinc and manganese and their application method on yield and yield components of common bean (*Phaseolus vulgaris* L. CV. Khomein). *Middle-East Journal of Science and Research*. 2011; 8(5):859-865. Retrieved:[https://www.idosi.org/mejsr/mejsr8\(5\)11/3.pdf](https://www.idosi.org/mejsr/mejsr8(5)11/3.pdf)
  23. Habbasha SF, Mohamed MH, Abd El-Lateef EM, Mekki BB, Ibrahim ME. Effect of combined zinc and nitrogen on yield, chemical constituents and nitrogen use efficiency of some chickpea cultivars under sandy soil conditions. *World J. Agric. Sci*. 2013;9(4):354-360.
  24. Rietra RP, Heinen M, Dimkpa CO, Bindraban PS. Effects of nutrient antagonism and synergism on yield and fertilizer use efficiency. *Communications in Soil Science and Plant Analysis*. 2017;48: 1895–1920. Available:<https://doi.org/10.1080/00103624.2017.1407429>.
  25. Khan MMH, Ahmed N, Naqvi SAH, Ahmad B, Dawar K, Rahie AA, Danish S. Synchronization of zinc and boron application methods and rates for improving the quality and yield attributes of *Mangifera indica* L. on sustainable basis. *Journal of King Saud University – Science*. 2022;34(8):102280.

- Available:<https://doi.org/10.1016/j.jksus.2022.102280>
26. Salehin F, Rahman S. Effect of zinc and nitrogen fertilizer and their application method on yield and yield components of *Phaseolus vulgaris* (L.) Agric. Sci. 2012; 3(1):9-13.
  27. Manjili MJ, Bidarigh S, Amiri E. Study the effect of foliar application of Nano chelate molybdenum fertilizer on the yield and yield components of peanut. Botany. 2014; 5(1):67-71.
  28. Hazra CR, Tripathi SB. Effect of secondary micronutrient on yield and quality of forages. Fertilizer News. 1998;43(12): 77-82.
  29. Sahu S, Lidder RS, Singh PK. Effect of micronutrients and biofertilizers on growth, yield and nutrient uptake by chickpea (*Cicer arietinum* L.) in Vertisols of Madhya Pradesh. Adv. Plant Sci. . 2008;21:501-503.
  30. Thapu U, Rai P, Suresh CP and Pal P. Effect of micronutrients on the growth and yield of pea in gangetic alluvial of West Bengal. Environ. Ecol. 2003;21:179-182.
  31. Meena KK, Meena RS, Kumawat SM. Effect of sulphur and iron fertilization on yield attributes, yield and nutrient uptake of mungbean (*Vigna radiata*). Indian Journal of Agricultural Sciences. 2012;83(4):472-6.
  32. Chauhan S, Titov A, Tomar DS. Effect of potassium, sulphur and zinc on growth, yield and oil content in soybean (*Glycine max* L.) in vertisols of central India. Indian J. Appl. Res. 2013;3(6):489-491.
  33. Badar R, Nisa Z, Ibrahim S. Supplementation of P with rhizobial inoculants to improve growth of Peanut plants. International Journal of Applied Research. 2015;1(4):9-23.
  34. Hirpara DV, Sakarvadia HL, Jadeja AS, Vekaria LC, Ponkia HP. Response of boron and molybdenum on groundnut (*Arachis hypogea* L.) under medium black calcareous soil. J Pharmacogn Phytochem. 2019;8(5):671-677.
  35. Brikics S, Milakovic Z, Kristek Z, Antunovic M. Pea yield and its quality depending on inoculation, nitrogen and molybdenum fertilization. Plant, Soil and Environ. 2004;50(1):39-45.
  36. Tantawy ELEM, Dalia ASN. Nodulation, growth, photosynthetic pigments and yield of broad bean plant (*Vicia faba* L.) as affected by nitrogen source, rhizobium inoculation and iron foliar application. J. Applied Sciences Research. 2013;9(1): 974-987.
  37. Piri Issa. Effect of phosphorus and micronutrients foliar application on sorghum yield. Annals of Bio. Res. 2012; 3(8):3998-4001.
  38. Balla H, Singh V, Tiwari D, Shaik MA. Effect of boron and molybdenum on growth rate and yield of groundnut (*Arachis hypogea* L.). J Pharmacogn Phytochem. 2020;9(6):1416-1419.
  39. Nandan B, Sharma BC, Chand G, Bazgalia K, Kumar R, Banotra M. Agronomic fortification of Zn and Fe in chickpea an emerging tool for nutritional security - A global perspective. Acta Scientific Nutritional Health. 2018;2(4):12-19.
  40. Usman M, Tahir M, Majeed MA. Effect of zinc sulphate as soil application and seed treatment on greengram (*Vigna radiata* L.). Pakistan Journal of Life and Social Sciences. 2014;12(2):87-91.
  41. Lokhande PB, Indulkar BS, Vaidya PH, Padghan AD, Wagh CB, Ingole AJ, Patil NM, Aundhkar A. Effect of phosphorus and zinc on yield and quality of greengram (*Vigna radiata* L.) in inceptisol. International Journal of Engineering Science and Computing. 2018;8(7):18647-18649.
  42. Singh G, Sekhon HS, Harpreet Kaur. Effect of farmyard manure, vermicompost and chemical nutrients on growth and yield of chickpea (*Cicer arietinum* L.). Int. J. Agric. Res. 2012;7(2):93-99.
  43. Khorgamy A, Farina A. Effect of phosphorus and zinc fertilisation on yield and yield components of chick pea cultivars. African Crop Science Conference Proceedings. 2009;9:205-208.
  44. Imsande J. Iron, sulphur and chlorophyll deficiencies: A need for an integrative approach in plant physiology. Plant Physiology. 1998;103:139-144.
  45. Oguz F. Research on the effects of different levels Molybdenum application on the yield and some yield components in Chickpea varieties on dry and irrigation conditions. M.Sc. Thesis, YuzuncuYil University Van, Turkey (unpublished); 2004.
  46. Roy PD, Narwal RP, Malik RS, Saha BN, Kumar S. Impact of zinc application methods on green gram (*Vigna radiata* L.) productivity and grain zinc fortification. J. Environ. Biol. 2014;35(5): 851-855.

47. Sharma A, Nakul HT, Jelgeri BR, Surwenshi A. Effect of micronutrients on growth, yield and yield components in pigeon pea (*Cajanus cajan* L.). Res. J. Agric. Sci. 2010;1:142-144.
48. Kumawat RN, Rathore PS, Talwar HS. Effect of sulphur and iron on crop growth attributes in summer green gram. Indian Journal of Plant Physiology. 2005;10(1):86-89.
49. Kumar R, Deka BC, Kumawat N, Ngachan SV. Effect of integrated nutrition, biofertilizers and zinc application on production potential and profitability of garden pea (*Pisum sativum* L.) in Eastern Himalaya, India. Legume Research. 2014; 37(6):614-620.
50. Valenciano JB, Miguélez-Frade MM, Marcelo V, Reinoso B. Response of irrigated common bean (*Phaseolus vulgaris*) yield to foliar zinc application in Spain. New Zealand Journal of Crop and Horticultural Science. 2010;35(3):325-330. Available:<https://doi.org/10.1080/01140670709510198>
51. Zeidan MS, Hozayn M, Abd-El-Salam MEE. Yield and quality of lentil as affected by micronutrient deficiencies in sandy soils. Journal of Applied Sciences Research. 2006;2(12):1342-1345. Retrieved:<http://www.aensiweb.com/old/jasr/jasr/1342-1345.pdf>
52. Shah KA, Gurjar R, Parmarand HC, Sonani VV. Effect of sulphur and zinc fertilization on yield and quality of pigeonpea in sandy loam soil. Green Farming. 2016;7(2):495-497.
53. Teixeira IR, Boren AB, Araujo GAA. Manganese and zinc leaf application on common bean grown on a cerrado soil. Scientia Agricola. 2004;61:77-81. Available:<https://doi.org/10.1590/S0103-90162004000100013>
54. Mady MA. Effect of foliar application with yeast extract and zinc on fruit setting and yield of Faba bean (*Vicia faba* L.). Journal of Biological, Chemical and Environmental Science. 2009;4(2):109-127. Retrieved:[https://www.bu.edu.eg/portal/uploads/Agriculture/Botany/1299/publications/Mohamed%20Ahmed%20Mohamed%20Mady\\_5.pdf](https://www.bu.edu.eg/portal/uploads/Agriculture/Botany/1299/publications/Mohamed%20Ahmed%20Mohamed%20Mady_5.pdf)
55. Rizk WM, Abdo FA. The response of two mungbean cultivars to zinc, manganese and boron II. Yield and chemical composition of seeds. Bulletin of Faculty of Agriculture, Cairo University. 2001;52(3):467-477.
56. Nasri M, Khalatbari M, Aliabadi Farahani H. Zn-foliar application influence on quality and quantity features in Phaseolus Vulgaris under different levels of N and K fertilizers. Advances in Environmental Biology. 2011;5(5):839-846. Retrieved:<http://www.aensiweb.com/old/aeb/2011/839-846.pdf>
57. Bozorgi HR, Faraji A, Danesh RK, Keshavarz A, Azarpour E, Tarighi F. Effect of plant density on yield and yield components of rice. World Applied Sciences Journal. 2011;12(11):2053-2057. Retrieved from:<https://citeseerx.ist.psu.edu/document?repid=rep1&type=pdf&doi=4db23100f9c40e0bf212d5ca1afc0453157aeee2>
58. Sarkar RK, Banik P. Response of green gram to nitrogen, phosphorus and molybdenum. Indian Journal of Hill Farming. 1991;5(1):75-76.
59. Ramakrishna A, Wani SP, Rao SC, Reddy SU. Effect of improved crop production technology on pigeonpea yield in resource poor rainfed areas. J. SAT Agric. Res. 2005;1(1):1-2.
60. Patil HM, Shete BT, Gaikwad CB. Effect of potash and zinc levels on growth and yield of pigeonpea. Ecology. 2006;12(2):341-344.
61. Ibrahim EA, Ramadanb WA. Cross effect of zinc foliar spray alone and combined with humic acid and chitosan on growth, nutrient elements content and yield of dry bean (*Phaseolus vulgaris* L.) plants sown at different dates. Scientia Horticulturae. 2015;184:101-105. Available:<https://doi.org/10.1016/j.scienta.2014.11.010>
62. Malik K, Kumar S, Arya KPS. Effect of zinc, molybdenum and urea on growth and yield of mungbean (*Vigna radiate* L. Wilczek). Adv. Res. J. of Crop Impro. 2015;6(1):58-62.
63. Naseri Rad H, Sayad V, Naseri Rad A. Effect of rhizobium bacteria (*Rhizobium leguminosarum*) and nano-Iron application on yield and yield components of different pinto beans genotypes. Agricultural Communication. 2014;2(2):22-27. Retrieved:[https://www.researchgate.net/publication/261885406\\_Effect\\_of\\_Rhizobium\\_Bacteria\\_Rhizobium\\_leguminosarum\\_and\\_Nano-Iron\\_Application\\_on\\_Yield\\_and](https://www.researchgate.net/publication/261885406_Effect_of_Rhizobium_Bacteria_Rhizobium_leguminosarum_and_Nano-Iron_Application_on_Yield_and)

- \_Yield\_ Components\_of\_Different\_Pinto\_Beans\_Genotypes
64. Heidarian AR, Kord H, Mostafavi K, Parviz Lak A, Amini Mashhadi F. Investigating Fe and Zn foliar application on yield and its components of soybean (*Glycine max* (L.) Merr.) at different growth stages. Journal of Agricultural Biotechnology and Sustainable Development, 20113(9):189-197. Retrieved from:[https://academicjournals.org/article/article1379436818\\_Heidarian%20et%20al.pdf](https://academicjournals.org/article/article1379436818_Heidarian%20et%20al.pdf)
65. Khan n, Tariq m, Ullah k, Muhammad d, Khan i, Rahatullah k, Ahmed n, Ahmed s. The effect of molybdenum and iron on nodulation, nitrogen fixation and yield of Chickpea Genotypes (*Cicer arietinum* L). Journal of Agriculture and Veterinary Science. 2014;7(1):63-79.
66. Mali GC, Sharma NN, Acharya HK, Gupta SK, Gupta PK. Response of pigeon pea to S and Zn fertilization Vertisols in South eastern plain of Rajasthan. Proceedings of National symposium on Arid Legumes for Food Nutrition Security and Promotion Trade, Hisar, India. 2003;15-16 May 2002:267-271.
67. Mali GC, Sharma NN, Acharya HK, Gupta SK, Gupta PK. Response of pigeon pea to S and Zn fertilization Vertisols in South eastern plain of Rajasthan. Proceedings of National symposium on Arid Legumes for Food Nutrition Security and Promotion Trade, Hisar, India. 2003;15-16 May 2002:267-271.
68. Salam PK, Rajput RS, Mishra PK, Anita, Shrivastava GK. Effect of micronutrients fertilization on productivity potential of urdbean. Ann. Agric. Res. New Series. 2004;25:329-332.
69. Kumawat RN, Rathore PS, Pareek N. Response of mung bean to sulphur and iron nutrition grown on calcareous soil of Western Rajasthan. Indian J. Pulse Res. 2006;19:228-230.
70. Kumar V, Dwivedi VN, Tiwari DD. Effect of phosphorus and iron on yield and mineral nutrition in chickpea. Ann. Plant Soil Res. 2009;11:16- 18.

© 2024 Pandey et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/111562>