



Line × Tester Method Used to Evaluate the Inbred Lines of Field Corn

M. M. Hoque¹, H. Z. Raihan¹, Tanjila Nasreen Trina² and Md. Razzab Ali^{3*}

¹*Plant Breeding Division, Bangladesh Agricultural Research Institute, Gazipur-1701, Bangladesh.*

²*Bangladesh Institute of Nuclear Agriculture, BINA Substation, Rangpur-5400, Bangladesh.*

³*Horticulture Research Centre, Bangladesh Agricultural Research Institute, Gazipur-1701, Bangladesh.*

Authors' contributions

This work was carried out in collaboration among all authors. Author MMH designed the study, performed the statistical analysis, wrote the protocol and wrote the first draft of the manuscript. Author HZR and MRA managed the analyses of the study. Author TNT managed the literature searches. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJAAR/2020/v14i430138

Editor(s):

(1) Dr. Daniele De Wrachien, University of Milan, Italy.

Reviewers:

(1) John C. Nwite, Federal College of Agriculture, Nigeria.

(2) Imene Rajhi, Centre of Biotechnology of Borj Cedria, Tunisia.

Complete Peer review History: <http://www.sdiarticle4.com/review-history/63436>

Original Research Article

Received 02 October 2020

Accepted 08 December 2020

Published 22 December 2020

ABSTRACT

Nineteen selected lines of field corn were crossed in a line × tester method with two testers to produce 38 hybrids during rabi 2018-2019. In the following year, all the hybrids were raised along with five commercial checks in an alpha lattice design with two replications. The lines E34, BML75, BML76, BML249, BIL106, CML465, CML481 and CML487 were better among the parents, showing GCA effects for yield and other traits could be used extensively in hybrid breeding program owing to increase yield. Furthermore, based on mean, SCA effects and standard heterosis of yield value, the crosses BML75 × BIL79, E34XBIL157, BML76 × BIL157 and BML249 × BIL157 were found to be better to increase the grain yield along with other traits and play pivotal role to development of commercial hybrid.

Keywords: Field corn; inbred lines; Line × tester; combining ability.

*Corresponding author: Email: razzab321@gmail.com;

1. INTRODUCTION

Maize (*Zea mays* L.) is a versatile crop with wider genetic variability and able to grow successfully all over the world covering all agro-climatic conditions viz. tropical, subtropical and temperate. Maize acreage and production have an increasing tendency, due to introduction of high yielding hybrids. Efforts are, therefore, required to develop hybrids with high yield potential in order to increase production of maize. Most resourceful use of nineteen inbreds would be possible only when reasonable information on the amount and type of genetic variation and combining ability effects in the materials is available. Heterosis and combining ability is prerequisite for developing a good economically viable hybrid maize variety [1]. Combining ability analysis is convenient to assess the potential inbred lines and also helps in identifying the nature of gene action involved in various quantitative traits. Combining ability is divided into two parts such as general combining ability (GCA) and specific combining ability (SCA). The GCA and SCA variances have been determined and related to the possible types of gene action involved. GCA is a good estimate of additive gene action, whereas SCA is a measure of non-additive gene action [2]. This information plays significant role in case of plant breeders for formulating hybrid breeding programs. A wide range of biometrical tools is available to breeders for characterizing genetic control of economically significant traits as a guide to decide upon a proper breeding methodology to encompass in hybrid breeding [3]. Line \times tester mating design developed by Kempthorne [4], which provides reliable information on the general and specific combining ability effects of parents and their hybrid combinations was used to generate the information. The design has been extensively used in maize by many researchers like, Joshi et al. [5] and Sharma et al. [6] and continues to be applied in quantitative genetic studies. The line \times tester analysis provides information on GCA of parents and specific combining ability (SCA) of hybrids which helps to identify good quality inbreds and hybrids, respectively [7,8]. Therefore, the present research work was undertaken to identify the nature and magnitude of gene action for yield and other important traits in maize.

2. MATERIALS AND METHODS

Nineteen inbred lines (as female parents) and two testers viz. BIL 79 and BIL 157 (as male

parents) were selected and crossed line \times Tester fashion in two isolation (Tuber Crop Research Sub Station, Bogra and Regional Horticultural Research Centre, Sibpur, Narshingdi) in 2018-19. The present investigation was carried out during rabi, 2019-20 at the experimental field of BARI, Gazipur. Experimental materials comprised of 38 F_1 's along with five checks viz. BARI hybrid maize-16, 981, Don111, Mohabir and Miracle. Seeds were sown at 1 December, 2019. Urea, Triple super phosphate (TSP), Muriate of potash (MoP), Gypsum, $ZnSO_4$ and boric acid were used as source of nitrogen, phosphorous, potassium, Sulphur, Zinc and Boron respectively. Cow dung (5 t ha^{-1}), urea (250 kg ha^{-1}), TSP (55 kg ha^{-1}), MoP (110 kg ha^{-1}) Gypsum (40 kg ha^{-1}), $ZnSO_4$ (5 kg ha^{-1}) and boric acid (1.5 kg ha^{-1}) were applied in the soil. The full amount of cow dung, one third of Urea, TSP and MoP, Gypsum, $ZnSO_4$ and boric acid was applied as basal dose at the time of final land preparation. Rest of the Urea was applied with 2 equal splits at 8 leaf stage and tassel emergence of plant. One healthy seedling was kept after proper thinning. Irrigation along with other inter cultural operations were done as and when necessary.

2.1 Design and Layout of the Experiment

The experiment was evaluated in Alpha lattice design with two replications having forty three plots consisted of single row of four meter lengths with row to row distance of 60 cm and plant to plant of 25 cm.

2.2 Data Collection

The data were taken on five randomly selected competitive plants from the rows of each plot for plant height (cm) and ear height (cm). Observations were recorded on the whole plot basis in respect of days to 50% tasseling, days to 50% silking, Days to maturity and measurement of grain yield, which data was converted to t ha^{-1} .

2.3 Statistical Analysis

The collected data were compiled and tabulated, which were subjected to Line \times Tester analyses following standard procedure given by AGRISTAT Package. Analysis of variance for combining ability and the mean sum of squares are presented in Table 1.

3. RESULTS AND DISCUSSION

The analysis result (Table 1) revealed that genotypes exhibited highly (1 percent) significant

differences among themselves for days to 50% tasseling, days to 50% silking, days to maturity, plant height, ear height and grain yield, demonstrating wide range of genetic variability among the genotypes. An experiment conducted by Talukder et al. [1], also observed genotypic difference for days to tasseling, days to silking, plant height, ear height and grain yield. The crosses showed significant differences, indicating varying performance of cross combinations. When the effects of crosses were partitioned into lines, testers and line \times tester effects. the interaction effects (line \times testers) were found to be significant for days to 50% tasseling, days to 50% silking, days to maturity, plant height, ear height and grain yield under study specifying the role of dominance and non-additive effects in all traits. The parents exhibited significant differences for days to 50% tasseling, days to 50% silking, days to maturity, plant height, ear height and grain yield, indicating greater diversity in the parental lines. Testers also showed significant variations among themselves except days to 50% tasseling and days to 50% silking. This result was agreement with the findings of Tucak et al. [9] on alfalfa for dry matter yield and Atif et al. [10] on maize for cob diameter, cob weight, number of kernels cob^{-1} , grain yield and harvest index, in case of testers, lines and line \times tester interaction in their study. Highly significant differences were also observed in checks for all characters except days to maturity. A comparison of magnitude of variance components due to gca and sca confirm the gene action in controlling the expression of traits. The ratio of gca and sca for all the traits were less than one except days to silking which indicated that these characters were governed by non-additive gene effects. Similar findings were reported by Kumar et al. [11] and Alam et al. [12] for grain yield, days to tasseling, days to silking, plant height, ear height and some other characters in maize in their study.

The proportional contribution of lines was higher for days to 50% tasseling, days to 50% silking, days to maturity, plant height, ear height and grain yield, representing their predominant maternal influence (Table 2). Raihan and Hoque [13] and Alam et al. [12] also obtained higher value for contribution to line than contribution to tester and line \times Tester for the same characters in maize. Motamedi et al. [14] was conducted an

experiment on maize and obtained less influence of testers regarding kernel yield.

3.1 General Combining Ability (GCA) Effects

The GCA effects of the parents are presented in Table 3. Negative estimates are considered desirable for days to tasseling, silking, maturity and plant height, ear height as those were observed to be associated with earliness dwarfness. Lines BIL 177 & BIL189 exhibited desirable negatively significant GCA effects for days to tasseling, silking and maturity. These two lines also showed negatively significant gca for plant height and ear height. Ahmed and Amiruzzaman [15] found similar results in some parents for the same characters in maize. However, their gca for yield were negative significant which is unexpected. Line E34, BML75, BML249 and CML481 showed positively significant gca for yield and negative significant gca for tasseling, silking, maturity, plant height and ear height. Ahmed and Amiruzzaman [15] also found positive significant gca for yield and negative significant gca for tasseling, silking, maturity, plant height and ear height in some parents in their study. Line BML76, BIL106, CML465 and CML487 showed positive and significant (1percent) gca for yield whereas most of the growth parameters along with maturity were negative significant. Similar result was also observed by Mia and Biswas [16]. So, Line E34, BML75, BML249, CML481 BML76, BIL106, CML465 and CML487 could be used in future breeding program.

3.2 Specific Combining Ability (SCA) Effects

High positive estimates of SCA in absolute values indicates that hybrid performance is relatively superior or inferior to parent lines general combining ability, showing the interactions of non-additive interactions resulting from the complementation degree among parent lines in relation frequency of alleles in loci in some dominance, while low estimates of sca in absolute value indicates that hybrids behave as expected in relation to gca of parent lines [17]. In the selection of parent lines used to produce hybrids, the effect of sca analyzed in an isolated way has a limiting value. Thus other parameters should be considered such as mean of hybrids and gca of the respective parent lines.

Table 1. Mean sum square and estimates of variance for yield and yield contributing traits of maize

Source	D.F	DT	DS	DM	PH	EH	YI
Genotype	42	14.62**	17.45**	38.37**	106.58**	134.76**	10.02**
Cross	37	15.63**	18.77**	43.30**	117.61**	145.86**	6.68**
Line	18	28.99*	34.15**	49.59**	157.56**	194.30**	7.73**
Tester	1	0.33	0.65	250.58**	180.12**	938.01**	43.25**
Lx T	18	3.11**	4.40**	25.50**	74.09**	53.40**	3.60**
Checks	4	8.65**	9.65**	2.35	9.85**	40.60**	4.32**
Check vs Cross	1	1.18	0.06	0.002	85.65**	100.74**	156.25**
Error	42	4.61	4.58	7.0537	37.37	16.88	0.43
Estimation of component of variance							
σ ² g(Line)		6.472	7.439	6.024	20.890	35.225	1.031
σ ² g(Tester)		0.352	-0.099	5.923	2.790	23.279	1.043
σ ² gca		0.222	0.255	0.315	0.770	1.637	0.055
σ ² sca		-0.796	-0.220	9.073	16.239	17.379	1.601
		-0.279	-1.159	0.035	0.047	0.094	0.034

* and ** Significant at 5 & 1 percent level, respectively; DT= Days to 50% Tasseling, DS= Days to 50% Silking, DM= Days to Maturity, PH= Plant height (cm), EH= Ear Height (cm), Yi= Grain Yield (t ha⁻¹)

Table 2. Propotional contribution (%) of lines, testers and their interactions to total variance in maize

Sources	DT	DS	DM	PH	EH	Yield
Contribution due to line	90.27	88.52	55.72	65.21	64.81	56.27
Contribution due to Tester	0.06	0.09	15.64	4.14	17.38	17.50
Contribution due to LxT	9.67	11.39	28.64	30.65	17.81	26.23

DT= Days to 50% Tasseling, DS= Days to 50% Silking, DM= Days to Maturity, PH= Plant height (cm), EH= Ear Height (cm), Yi= Grain Yield (t ha⁻¹)

Table 3. General combining ability (GCA) effects of yield and other characters

Line/Tester	DT	DS	DM	PH	EH	Yield(t ha ⁻¹)
Line						
1.Ki 21	1.57ns	1.88ns	0.14ns	-0.99ns	6.97**	-0.43ns
2.E34	-1.68ns	-2.37*	-5.11**	5.51ns	3.72ns	0.84*
3. BML76	1.32ns	1.13ns	-5.36**	-6.49*	7.72**	1.49**
4. E37	5.32 **	5.38 **	0.89 ns	2.26 ns	-4.03 ns	-0.46 ns
5. BIL65	-1.68 ns	-1.37 ns	4.14 **	10.74 **	18.78 **	-1.04 **
6. BIL 106	0.82 ns	0.13 ns	1.39 ns	-0.49 ns	6.22 **	0.90 **
7. BIL 182	0.57 ns	0.38 ns	-1.86 ns	-9.24 **	-0.28 ns	-2.02 **
8. BML 249	0.07 ns	0.38 ns	0.89 ns	2.76 ns	0.97 ns	2.53 **
9. 900M6	-0.43 ns	0.13 ns	4.64 **	8.01 *	0.47 ns	-0.15 ns
10. 900M10	0.32 ns	1.38 ns	2.89 *	3.76 ns	-1.28 ns	-0.80 *
11. BML75	-0.43 ns	-0.62 ns	-3.36 *	3.01 ns	-0.78 ns	2.18 **
12.PINA20	-0.93 ns	-1.12 ns	1.14 ns	9.01 **	2.22 ns	-1.47 **
13. CML 465	0.82 ns	0.63 ns	0.64 ns	10.01 **	9.47 **	0.97 **
14. CML 487	1.32 ns	0.88 ns	1.39 ns	-3.24 ns	10.22 **	0.70 *
15. CML480	1.57 ns	1.63 ns	3.14 *	2.26 ns	-0.03 ns	0.51 ns
16. CML 481	-0.18 ns	-0.37 ns	2.89 *	1.26 ns	-3.53 ns	0.84 *
17. CML 496	3.57 **	4.63 **	2.89 *	-0.24 ns	-5.53 *	-0.42 ns
18. BIL 177	-6.93 **	-7.37 **	-3.86 **	-7.74 *	-8.28 **	-2.45 **
19. BIL 189	-4.93 **	-5.37 **	-7.61 **	-8.74 **	-5.53 *	-1.71 **
SE(gi)	1.084	1.099	1.356	3.225	2.159	0.316
SE(gi-gj)	1.533	1.555	1.917	4.561	3.053	0.447
Tester						
1. BIL79	0.07	-0.09	-1.82	1.54	3.51	-0.75
2. BIL157	-0.07	0.09	1.82	-1.54	-3.51	0.75
SE(gj)	0.352	0.357	0.440	1.047	0.701	0.103
SE(gi-gj)	0.497	0.504	0.622	1.480	0.991	0.145

DT= Days to 50% Tasseling, DS=Days to 50% silking, DM= Days to Maturity, PH= Plant Height(cm), EH= Ear height

Table 4. Mean and specific combining ability (SCA) effects for different characters

Crosses	DT		DS		DM	
	Mean	SCA	Mean	SCA	Mean	SCA
1.Ki 21x Bil 79	97	-0.07 ns	101.5	-0.41 ns	143	-2.43 ns
2.Ki 21 x BIL 157	97	0.07 ns	102.5	0.41 ns	151.5	2.43 ns
3.E 34 x BIL 79	94	0.18 ns	98	0.34 ns	136.5	-3.68 ns
4. E 34 X BIL 157	93.5	-0.18 ns	97.5	-0.34 ns	147.5	3.68 ns
5. BML76 X BIL 79	97.5	0.68 ns	102.5	1.34 ns	138	-1.93 ns
6. BML76 X BIL 157	96	-0.68 ns	100	-1.34 ns	145.5	1.93 ns
7. E 37 X BIL 79	99	-1.82 ns	103	-2.41 ns	145.5	-0.68 ns
8. E 37 X BIL157	102.5	1.82 ns	108	2.41 ns	150.5	0.68 ns
9. BIL65 X BIL79	94	0.18 ns	99	0.34 ns	152.5	3.07 ns
10. BIL65 X BIL157	93.5	-0.18 ns	98.5	-0.34 ns	150	-3.07 ns
11. BIL106 X BIL79	96.5	0.18 ns	100	-0.16 ns	149.5	2.82 ns
12.BIL 106 X BIL157	96	-0.18 ns	100.5	0.16 ns	147.5	-2.82 ns
13.BIL 182 XBIL79	97.5	1.43 ns	101.5	1.09 ns	142.5	-0.93 ns
14. BIL1182 X BIL157	94.5	-1.43 ns	99.5	-1.09 ns	148	0.93 ns
15. BML249 X BIL79	95	-0.57 ns	99.5	-0.91 ns	144.5	-1.68 ns
16. BML249 X BIL157	96	0.57 ns	101.5	0.91 ns	151.5	1.68 ns
17. 900M6 X BIL79	95	-0.07 ns	101	0.84 ns	148.5	-1.43 ns
18 900M6 X BIL157	95	0.07 ns	99.5	-0.84 ns	155	1.43 ns
19. 900M10 X BIL79	95.5	-0.32 ns	100	-1.41 ns	150	1.82 ns
20. 900M10 X BIL157	96	0.32 ns	103	1.41 ns	150	-1.82 ns
21. BML75 X BIL79	95	-0.07 ns	99.5	0.09 ns	142.5	0.57 ns
22. BML75 X BIL157	95	0.07 ns	99.5	-0.09 ns	145	-0.57 ns
23. PINA 20 X BIL79	95	0.43 ns	99	0.09 ns	147.5	1.07 ns
24. PINA20 X BIL157	94	-0.43 ns	99	-0.09 ns	149	-1.07 ns
25.CML465 X BIL79	96.5	0.18 ns	101	0.34 ns	147	1.07 ns
26. CML465 X BIL157	96	-0.18 ns	100.5	-0.34 ns	148.5	-1.07 ns
27. CML480 X BIL79	96.5	-0.32 ns	100	-0.91 ns	147	0.32 ns
28. CML480 X BIL157	97	0.32 ns	102	0.91 ns	150	-0.32 ns
29. CML481 X BIL79	97.5	0.43 ns	103	1.34 ns	149	0.57 ns
30. CML481 X BIL157	96.5	-0.43 ns	100.5	-1.34 ns	151.5	-0.57 ns
31. CML487 X BIL79	93	-2.32 ns	98	-1.66 ns	150	1.82 ns
32. CML487 X BIL157	97.5	2.32 ns	101.5	1.66 ns	150	-1.82 ns
33. CML496 X BIL79	99	-0.07 ns	105	0.34 ns	147.5	-0.68 ns
34. CML496 X BIL157	99	0.07 ns	104.5	-0.34 ns	152.5	0.68 ns
35. BIL177 X BIL79	89.5	0.93 ns	93.5	0.84 ns	147	5.57 **
36. BIL177 X BIL157	87.5	-0.93 ns	92	-0.84 ns	139.5	-5.57 **
37. BIL189 X BIL79	91.5	0.93 ns	95.5	0.84 ns	132.5	-5.18 **
38. BIL189 X BIL157	89.50	-0.93ns	94.00	-0.84ns	146.50	5.18**
SE(Sij)	1.533		1.555		1.917	
SE(Sij-Skl)	2.168		2.199		2.711	

Table 4. (Cont'd)

Crosses	PH		EH		Yield (t ha ⁻¹)	
	Mean	SCA	Mean	SCA	Mean	SCA
1.Ki 21x Bil 79	187.5	-2.04 ns	106	-1.26 ns	8.92	0.64 ns
2.Ki 21 x BIL 157	188.5	2.04 ns	101.5	1.26 ns	9.16	-0.64 ns
3.E 34 x BIL 79	196	-0.04 ns	104	-0.01 ns	8.2	-1.35 **
4. E 34 X BIL 157	193	0.04 ns	97	0.01 ns	12.41	1.35 **
5. BML76 X BIL 79	177.5	-6.54 ns	105.5	-2.51 ns	9.17	-1.03 *
6. BML76 X BIL 157	187.5	6.54 ns	103.5	2.51 ns	12.74	1.03 *
7. E 37 X BIL 79	199	6.21 ns	100.5	4.24 ns	9.17	0.92 *
8. E 37 X BIL157	183.5	-6.21 ns	85	-4.24 ns	8.84	-0.92 *
9. BIL65 X BIL79	184	4.21 ns	82.5	0.99 ns	8.28	0.60 ns
10. BIL65 X BIL157	172.5	-4.21 ns	73.5	-0.99 ns	8.59	-0.60 ns
11. BIL106 X BIL79	187.5	-2.54 ns	110.5	3.99 ns	9.99	0.38 ns
12.BIL 106 X BIL157	189.5	2.54 ns	95.5	-3.99 ns	10.74	-0.38 ns
13.BIL 182 XBIL79	185.5	4.21 ns	99.5	-0.51 ns	6.33	-0.36 ns

Crosses	PH		EH		Yield (t ha ⁻¹)	
	Mean	SCA	Mean	SCA	Mean	SCA
14. BIL1182 X BIL157	174	-4.21 ns	93.5	0.51 ns	8.57	0.36 ns
15. BML249 X BIL79	186	-7.29 ns	92.5	-8.76 **	8.97	-2.27 **
16. BML249 X BIL157	197.5	7.29 ns	103	8.76 **	15.02	2.27 **
17. 900M6 X BIL79	204.5	5.96 ns	105.5	4.74 ns	9.03	0.46 ns
18 900M6 X BIL157	189.5	-5.96 ns	89	-4.74 ns	9.62	-0.46 ns
19. 900M10 X BIL79	191	-3.29 ns	96	-3.01 ns	7.99	0.07 ns
20. 900M10 X BIL157	194.5	3.29 ns	95	3.01 ns	9.35	-0.07 ns
21. BML75 X BIL79	193.5	-0.04 ns	94	-5.51 ns	12.41	1.51 **
22. BML75 X BIL157	190.5	0.04 ns	98	5.51 ns	10.9	-1.51 **
23. PINA 20 X BIL79	205.5	5.96 ns	102	-0.51 ns	7.15	-0.09 ns
24. PINA20 X BIL157	190.5	-5.96 ns	96	0.51 ns	8.84	0.09 ns
25.CML465 X BIL79	199	-1.54 ns	110.5	0.74 ns	9.32	-0.37 ns
26. CML465 X BIL157	199	1.54 ns	102	-0.74 ns	11.57	0.37 ns
27. CML480 X BIL79	186.5	-0.79 ns	111.5	0.99 ns	8.87	-0.55 ns
28. CML480 X BIL157	185	0.79 ns	102.5	-0.99 ns	11.48	0.55 ns
29. CML481 X BIL79	191.5	-1.29 ns	104.5	4.24 ns	8.95	-0.28 ns
30. CML481 X BIL157	191	1.29 ns	89	-4.24 ns	11.02	0.28 ns
31. CML487 X BIL79	188.5	-3.29 ns	94	-2.76 ns	9.62	0.06 ns
32. CML487 X BIL157	192	3.29 ns	92.5	2.76 ns	11	-0.06 ns
33. CML496 X BIL79	188	-2.29 ns	92.5	-2.26 ns	7.94	-0.35 ns
34. CML496 X BIL157	189.5	2.29 ns	90	2.26 ns	10.16	0.35 ns
35. BIL177 X BIL79	189.5	6.71 ns	95.5	3.49 ns	8.02	1.75 **
36. BIL177 X BIL157	173	-6.71 ns	81.5	-3.49 ns	6.02	-1.75 **
37. BIL189 X BIL79	179.5	-2.29 ns	98.5	3.74 ns	7.27	0.27 ns
38. BIL189 X BIL157	181	2.29 ns	84	-3.74 ns	8.25	-0.27 ns
SE(Sij)		4.561		3.053		0.472
SE(Sij-Ski)		6.451		4.318		0.632

DT= Days to 50% Tasseling, DS= Days to 50% Silking, DM= Days to Maturity, PH= Plant height (cm), EH= Ear Height (cm), Yi= Grain Yield (t ha⁻¹)

Table 5. Heterosis (%) standard heterosis over check BHM16 for different characters

Crosses	DT	DS	DM	PH	EH	YIELD
1.Ki 21x Bil 79	3.19 ns	3.05 ns	-1.72 ns	-1.32 ns	10.99 *	-21.75 **
2.Ki 21 x BIL 157	3.19 ns	4.06 ns	4.12 *	-0.79 ns	6.28 ns	-19.69 **
3.E 34 x BIL 79	0.00 ns	-0.51 ns	-6.19 **	3.16 ns	8.90 *	-28.07 **
4. E 34 X BIL 157	-0.53 ns	-1.02 ns	1.37 ns	1.58 ns	1.57 ns	8.90 ns
5. BML76 X BIL 79	3.72 ns	4.06 ns	-5.15 **	-6.58 *	10.47 *	-19.56 **
6. BML76 X BIL 157	2.13 ns	1.52 ns	0.00 ns	-1.32 ns	8.38 ns	11.80 *
7. E 37 X BIL 79	5.32 *	4.57 *	0.00 ns	4.74 ns	5.24 ns	-19.56 **
8. E 37 X BIL157	9.04 **	9.64 **	3.44 ns	-3.42 ns	10.99 *	-22.41 **
9. BIL65 X BIL79	0.00 ns	0.51 ns	4.81 *	-3.16 ns	13.61 **	-27.37 **
10. BIL65 X BIL157	-0.53 ns	0.00 ns	3.09 ns	-9.21 **	23.04 **	-24.69 **
11. BIL106 X BIL79	2.66 ns	1.52 ns	2.75 ns	-1.32 ns	15.71 **	-12.37 *
12.BIL 106 X BIL157	2.13 ns	2.03 ns	1.37 ns	-0.26 ns	0.00 ns	-5.75 ns
13.BIL 182 XBIL79	3.72 ns	3.05 ns	-2.06 ns	-2.37 ns	4.19 ns	-44.47 **
14. BIL1182 X BIL157	0.53 ns	1.02 ns	1.72 ns	-8.42 *	-2.09 ns	-24.87 **
15. BML249 X BIL79	1.06 ns	1.02 ns	-0.69 ns	-2.11 ns	-3.14 ns	-21.27 **
16. BML249 X BIL157	2.13 ns	3.05 ns	4.12 *	3.95 ns	7.85 ns	31.75 **
17. 900M6 X BIL79	1.06 ns	2.54 ns	2.06 ns	7.63 *	10.47 *	-20.75 **
18 900M6 X BIL157	1.06 ns	1.02 ns	6.53 **	-0.26 ns	-6.81 ns	-15.66 **
19. 900M10 X BIL79	1.60 ns	1.52 ns	3.09 ns	0.53 ns	0.52 ns	-29.91 **
20. 900M10 X BIL157	2.13 ns	4.57 *	3.09 ns	2.37 ns	-0.52 ns	-17.98 **
21. BML75 X BIL79	1.06 ns	1.02 ns	-2.06 ns	1.84 ns	-1.57 ns	8.82 ns
22. BML75 X BIL157	1.06 ns	1.02 ns	-0.34 ns	0.26 ns	2.62 ns	-4.39 ns
23. PINA 20 X BIL79	1.06 ns	0.51 ns	1.37 ns	8.16 *	6.81 ns	-37.24 **
24. PINA20 X BIL157	0.00 ns	0.51 ns	2.41 ns	0.26 ns	0.52 ns	-22.46 **
25.CML465 X BIL79	2.66 ns	2.54 ns	1.03 ns	4.74 ns	15.71 **	-18.25 **
26. CML465 X BIL157	2.13 ns	2.03 ns	2.06 ns	4.74 ns	6.81 ns	1.49 ns
27. CML480 X BIL79	2.66 ns	1.52 ns	1.03 ns	-1.84 ns	16.75 **	-22.19 **
28. CML480 X BIL157	3.19 ns	3.55 ns	3.09 ns	-2.63 ns	7.33 ns	0.70 ns

Crosses	DT	DS	DM	PH	EH	YIELD
29. CML481 X BIL79	3.72 ns	4.57 *	2.41 ns	0.79 ns	9.42 *	-21.49 **
30. CML481 X BIL157	2.66 ns	2.03 ns	4.12 *	0.53 ns	-6.81 ns	-3.33 ns
31. CML487 X BIL79	-1.06 ns	-0.51 ns	3.09 ns	-0.79 ns	-1.57 ns	-15.61 **
32. CML487 X BIL157	3.72 ns	3.05 ns	3.09 ns	1.05 ns	-3.14 ns	-3.46 ns
33. CML496 X BIL79	5.32 *	6.60 **	1.37 ns	-1.05 ns	-3.14 ns	-30.31 **
34. CML496 X BIL157	5.32 *	6.09 **	4.81 *	-0.26 ns	-5.76 ns	-10.92 ns
35. BIL177 X BIL79	-4.79 *	-5.08 *	1.03 ns	-0.26 ns	0.00 ns	-29.65 **
36. BIL177 X BIL157	-6.91 **	-6.60 **	-4.12 *	-8.95 **	14.66 **	-47.15 **
37. BIL189 X BIL79	-2.66 ns	-3.05 ns	-8.93 **	-5.53 ns	3.14 ns	-36.18 **
38. BIL189 X BIL157	-4.79 *	-4.57 *	0.69 ns	-4.74 ns	12.04 **	-27.63 **
SE	1.518	1.514	1.878	4.322	2.905	0.464
CD(0.05)	4.315	4.303	5.338	12.287	8.259	1.320
CD(0.01)	5.753	5.737	7.118	16.382	11.012	1.760

DT= Days to 50% Tasseling, DS= Days to 50% Silking, DM= Days to Maturity, PH= Plant height (cm), EH= Ear Height (cm), Yi= Grain Yield (t ha⁻¹)

Therefore, superior hybrid combinations, which are important for breeding are involved at least. The mean value and sca effects of the hybrids are presented in the Table 4. In respect of days to tasseling and days to silking, no cross combination was recorded for significant and negative effects. Two cross combinations (BIL177x BIL79 and BIL177x BIL157) were recorded for significant and negative effects for early maturing plant. Similar findings were also reported by Alam et al. [12] and Bhavana et al. [18]. In case of maize, negative and significant value is expected for plant and ear height to develop short statured plant. No significant and negative effects were found for plant height. The lowest plant height (173 cm) was observed in BIL65 x BIL157 and BIL177x BIL157. One cross (BML249 x BIL79) showed significant and negative sca effect for ear height. Amin et al. [19] also observed significant and negative sca for ear height in maize. Six crosses (E34x BIL157, BML76x BIL157, E37x BIL79, BML249x BIL157, BML75x BIL79 and BIL177x BIL79) showed significant and positive sca effect for yield. Similar findings were also observed by Raihan and Hoque [13], Ahmed and Amiruzzaman [15] and Amin et al. [19].

3.3 HETEROSIS

The standard heterosis expressed by the F₁ hybrids over standard check BHM16 for different characters are presented in Table 5. Significant and positive heterosis is expected for yield and negative heterosis for other characters. Significant negative heterosis were observed in 3, 3, 4 and 4 crosses for days to tasseling, days to silking, plant height and ear height respectively. Talukder et al. [1] also observed significant negative heterosis in some crosses of mentioned characters of maize in their studies.

one parental line which has the most favorable effects of gca. Significant positive standard heterosis is expected for grain yield. Significant positive standard heterosis was observed in BML76x BIL157 (11.80%) and BML249x BIL157 (31.75%) for grain yield. On the other hand, non-significant and positive heterosis was observed in E34x BIL157 (8.90%) and BML75x BIL79 (8.82%). Akhi et al. [20] and Ahmed and Amiruzzaman [15] also observed positive and significant standard heterosis in some crosses for yield as well as negative significant heterosis in some crosses for maturity, growth parameter in maize in their studies.

4. CONCLUSION

The parental line E34, BML75, BML76, BML249, BIL106, CMI 465, CML 481 and CML487 were found promising and could be used extensively in hybrid breeding program owing to increase yield. Considering heterosis, sca value, mean value and gca of their parent the crosses E34 X BIL157, BML75 X BIL79, BML76 X BIL157, BML249 X BIL157 were found promising and could be useful towards increasing maize yield through developing hybrid.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Talukder MZA, Karim ANMS, Ahmed S, Amiruzzaman M. Combining ability and heterosis on yield and its component traits in maize (*Zea mays* L.). Bangladesh J. Agril. Res. 2016;41(3):565-577.

2. Sharief AE, El-Kalla SE, Gado HE, Yousef H. Heterosis in yellow maize. Australian J. Crop Science. 2009;3:146-154.
3. Gowda Rk, Kage U, Loththaswa HC, Shekara BG, Shobha D. Combining ability studies in maize (*Zea mays* L.). Molecular Plant Breeding. 2013;3:(14):116-127. DOI: 10.5376/ijh.2013.03.0014
4. Kempthorne O. An introduction to genetic statistics. New York, John Wiley and Sons, Inc. London: Chapman and Hall Limited;1957.
5. Joshi V, Dubey RB, Marker S. Combining ability for polygenic traits in early maturity hybrids of maize (*Zea mays* L.). Indian J. Genet.PI. Breed. 2002;62:312-315.
6. Sharma S, Narwal R, Kumar MS, Dass S. Line x tester analysis in maize (*Zea mays* L.). Forage Res. 2004;30:28-30.
7. Silva VQR, AmraJu AT, Gonçalves LSA, Freitas JSP, Candido LS, Vittorazzi C. Combining ability of tropical and temperate inbred lines of popcorn. Genet.Mole.Res. 2010;9(3):1742-1750.
8. Moterle LA, Braccini AL, Scapim CA, Pinto RJB, Goncalves LSA, Rodrigues R. Combining ability of popcorn lines for seed quality and agronomic traits. Euphytica. 2011;185(3):337-347.
9. Tucak M, Popovic S, Cupic T, Spanic V, Simic B, Meglic V. Combining abilities and heterosis for dry matter yield in alfalfa diallel crosses. NARDI FUNDULEA, Romanian Agril. Res. 2012;29:71-77.
10. Atif I, Awadalla A, Mutasim M. Combining ability and heterosis for yield and yield components in maize (*Zea mays* L.). Aust. J. Basic and Appl. Sci. 2012;6(10):36-41.
11. Kumar GP, Narsimha Reddy V, Sudheerkumar S, VenkateshwaraRao P, Combining ability studies in newly developed inbred lines in maize (*Zea mays* L.). Intl. J. of Pl., Ani.andEnv. Sci. 2014;4 (4):229-234.
12. Alam SS, Begum S, Amiruzzaman M. Evaluations of inbred lines of maize through line x Tester method (Set III). Annual Research Report: Maize, Barley, Millet and Sorghum. Plant Breeding Division, BARI, Gazipur. 2016;76-80.
13. Raihan H, Hoque. MM. Evaluations of inbred lines of maize through line x tester method. Annual Research Report: Maize, Barley, Millet and Sorghum. Plant Breeding Division, BARI, Gazipur. 2019;28-32.
14. Motamedi M, Choukan R, Hervan Em, Bihamta MR, Kajouri FD. Investigation of genetic control for yield and related traits in maize (*Zea mays* L.) lines derived from temperate and sub-tropical germplasm. Int. J. Biosciences. 2014;5(12):123-129.
15. Ahmed A, Amiruzzaman M, Evaluations of inbred lines of maize through line x tester method (Set VI).Annual Research Report: Maize, Barley, Millet and Sorghum. Plant Breeding Division, BARI, Gazipur. 2017;80-89.
16. Mia MA, Biswas A. Evaluations of inbred lines through through line x tester method (Set III). Annual Research Report: Maize, Barley, Millet and Sorghum. Plant Breeding Division, BARI, Gazipur. 201753-61.
17. Dhasarathan M, Babu C, Iyanar K. Combining ability and gene action studies for yield and quality traits in baby corn (*Zea mays* L.). SABRAO J. Breed. and Gene. 2015;47(1):60-69.
18. Bhavana P, Singh RP, Gadag RN. Gene action and heterosis for yield and yield components in maize (*Zea mays*). Indian J. of Agril. Sci. 2011;81(2):163-166.
19. Amin MN, Amiruzzaman M, Ahmed A, Ali MR. Evaluation of Inbred Lines of Maize (*Zea mays* L.) through Line x Tester Method. Bangladesh J. Agril.Res. 2014;39(4):675-683.
20. Akhi AH, Karim ANM, Ahmed S, Amiruzzaman M. Evaluations of inbred lines through line x Tester method (Set II). Annual Research Report: Maize, Barley, Millet and Sorghum. Plant Breeding Division, BARI, Gazipur. 2017;45-52.

© 2020 Hoque et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<http://www.sdiarticle4.com/review-history/63436>