

RESEARCH ARTICLE

A combined computational and experimental investigation of the filtration function of splenic macrophages in sickle cell disease

Guansheng Li¹, Yuhao Qiang², He Li^{3*}, Xuejin Li⁴, Pierre A. Buffet^{5*}, Ming Dao^{2*}, George Em Karniadakis^{1*}

1 Division of Applied Mathematics, Brown University, Providence, Rhode Island, United States of America, **2** Department of Materials Science and Engineering, Massachusetts Institute of Technology, Cambridge, Massachusetts, United States of America, **3** School of Chemical, Materials and Biomedical Engineering, University of Georgia, Athens, Georgia, United States of America, **4** Department of Engineering Mechanics and Center for X-Mechanics, Zhejiang University, Hangzhou, Zhejiang, China, **5** Université Paris Cité and Université des Antilles, Inserm, Biologie Intégrée du Globule Rouge, Paris, France

✉ These authors contributed equally to this work.

* he.li3@uga.edu (HL); pabuffet@gmail.com (PAB); mingdao@mit.edu (MD); george_karniadakis@brown.edu (GEK)



OPEN ACCESS

Citation: Li G, Qiang Y, Li H, Li X, Buffet PA, Dao M, et al. (2023) A combined computational and experimental investigation of the filtration function of splenic macrophages in sickle cell disease. *PLoS Comput Biol* 19(12): e1011223. <https://doi.org/10.1371/journal.pcbi.1011223>

Editor: Jeffrey J. Saucerman, University of Virginia, UNITED STATES

Received: May 30, 2023

Accepted: November 3, 2023

Published: December 13, 2023

Copyright: © 2023 Li et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All data are in the manuscript and/or [Supporting information](#) files.

Funding: National Institute of Health under grant number R01HL154150(G. E. K.). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: The authors have declared that no competing interests exist.

Abstract

Being the largest lymphatic organ in the body, the spleen also constantly controls the quality of red blood cells (RBCs) in circulation through its two major filtration components, namely interendothelial slits (IES) and red pulp macrophages. In contrast to the extensive studies in understanding the filtration function of IES, fewer works investigate how the splenic macrophages retain the aged and diseased RBCs, i.e., RBCs in sickle cell disease (SCD). Herein, we perform a computational study informed by companion experiments to quantify the dynamics of RBCs captured and retained by the macrophages. We first calibrate the parameters in the computational model based on microfluidic experimental measurements for sickle RBCs under normoxia and hypoxia, as those parameters are not available in the literature. Next, we quantify the impact of key factors expected to dictate the RBC retention by the macrophages in the spleen, namely, blood flow conditions, RBC aggregation, hematocrit, RBC morphology, and oxygen levels. Our simulation results show that hypoxic conditions could enhance the adhesion between the sickle RBCs and macrophages. This, in turn, increases the retention of RBCs by as much as four-fold, which could be a possible cause of RBC congestion in the spleen of patients with SCD. Our study on the impact of RBC aggregation illustrates a ‘clustering effect’, where multiple RBCs in one aggregate can make contact and adhere to the macrophages, leading to a higher retention rate than that resulting from RBC-macrophage pair interactions. Our simulations of sickle RBCs flowing past macrophages for a range of blood flow velocities indicate that the increased blood velocity could quickly attenuate the function of the red pulp macrophages on detaining aged or diseased RBCs, thereby providing a possible rationale for the slow blood flow in the open circulation of the spleen. Furthermore, we quantify the impact of RBC morphology on their tendency to be retained by the macrophages. We find that the sickle and granular-shaped RBCs are more likely to be filtered by macrophages in the spleen. This finding is consistent with the

observation of low percentages of these two forms of sickle RBCs in the blood smear of SCD patients. Taken together, our experimental and simulation results aid in our quantitative understanding of the function of splenic macrophages in retaining the diseased RBCs and provide an opportunity to combine such knowledge with the current knowledge of the interaction between IES and traversing RBCs to apprehend the complete filtration function of the spleen in SCD.

Author summary

The spleen, the largest organ of the lymphatic system within the abdominal region, is positioned adjacent to the greater curvature of the stomach and is ensconced within the omentum. It constantly controls the quality of red blood cells (RBCs) in circulation.

Erythrophagocytosis is one of the critical processes to remove senescent and diseased red blood cells (RBCs) from circulation in the spleen. In this study, we aim to quantify the influence of key factors that are expected to govern the retention of RBCs by macrophages in the spleen. These factors include blood flow conditions, RBC aggregation, hematocrit levels, RBC morphology, and oxygen levels. Our simulation results show that hypoxic conditions could enhance the adhesion between the sickle RBCs and macrophages. Taken together, our experimental and simulation results aid in our quantitative understanding of the function of splenic macrophages in retaining the diseased RBCs and provide an opportunity to combine such knowledge with the current knowledge of the interaction between IES and traversing RBCs to apprehend the complete filtration function of the spleen in SCD.

Introduction

The spleen, a dark red to blue-black elongated organ, is located in the abdomen and adjacent to the greater curvature of the stomach and within the omentum [1]. As the most significant secondary immune organ in the human body, the spleen prevents pathogenic microorganisms from remaining and multiplying in the bloodstream by initiating immune reactions [2, 3]. Another essential role of the spleen is to clear red blood cells (RBCs) from circulation when their biochemical or biomechanical properties hinder their circulation functionality. The two physiological functions of the spleen are accomplished within its two functionally and morphologically distinct compartments, namely white pulp and red pulp [4]. The white pulp comprises three subcompartments: the periarteriolar lymphoid sheath (PALS), the follicles, and the marginal zone (MZ). While the PALS contains central arterioles surrounded predominantly by the T cells, the follicles and the MZ are occupied mainly by B cells [2, 5–7]. After passing through the white pulp, blood flows into the medium-sized central artery from the splenic artery and then engages in either fast or slow microcirculation. As much as 90% of the splenic blood travels from the perifollicular zone to the venous sinus lumen through the fast microcirculation pathway [8], whereas only 10% of the blood flows through the slow microcirculation, where the RBCs need to traverse the red pulp, a three-dimensional meshwork of splenic cords and venous sinuses, which works like a blood filter that removes damaged and aged RBCs [2, 9, 10]. Before returning to the vascular beds, RBCs have to squeeze through the narrow slits between endothelial cells (IES) in the wall of sinuses, the most stringent challenge for the deformability of RBCs in microcirculation, resulting in the retention of less deformable RBCs or the removal of intraerythrocytic bodies [11–13].

In the red pulp, RBCs also make close contact with abundant splenic macrophages, which are capable of detecting the deleterious changes in the RBC membrane and engulfing the aged or damaged RBCs through erythrophagocytosis [2, 14, 15]. The key factors that determine the potential targets of phagocytosis have been considered to be the biochemical signaling triggered by the interaction between the ligands on the RBCs and the receptors on the macrophages, such as binding of antibodies (NAb)s to the band-3 proteins [16, 17], increased exposure of PS [18, 19], decreased expression of CD47 [20] and conformational changes in CD47 [21]. It is thought that this ligand-receptor interaction could generate so-called “eat me” signals and provoke RBC clearance by macrophages in the spleen. Recently, the impact of biophysical properties of the targeted cells and the macrophages are also investigated. An *in vitro* experimental investigation [22] suggested that the impact of RBC rigidity could override that of CD47 in the process of phagocytosis, indicating that less deformable RBCs are more likely to be engulfed by splenic macrophages. This finding was echoed by a subsequent study [23] showing a high propensity of macrophages to recognize and phagocytose the lysed and less deformable RBCs over intact ones.

Although the function of the spleen in RBC physiology has been well studied, its role in RBC disorders is not yet fully understood. For example, in sickle cell disease (SCD), the mutant sickle hemoglobin (HbS) could polymerize into stiff fiber bundles under hypoxia, causing the RBCs to sickle [24, 25]. Sickling of RBCs induces drastic alterations of the RBC rigidity and leads to their distorted heterogeneous shapes, which are categorized into elongated, granular, oval, holly-leaf, and crescent (classic sickle) shapes [26–28]. In addition to their increased stiffness, sickle RBCs are featured with enhanced cell adhesion, contributing to the initiation and propagation of vaso-occlusion events, a hallmark of SCD [29–31]. The spleen is one of the most common early organs to be affected in SCD. In SCD infants with still functional spleen, these sickle RBCs are amenable to mechanical retention by the spleen, a process that likely contributes to hemolytic anemia [32] and triggers acute splenic sequestration crises (ASSC), a life-threatening complication of SCD [33–40]. Children with SCD between five months and two years of age have a higher risk for ASSC, manifested as an abrupt fall in the hemoglobin level and splenomegaly [37, 41–45]. SCD patients with multiple episodes of ASSC require surgical splenectomy, a supportive transfusion program, or both. They are at higher risk of complications [46, 47]. To date, the mechanisms causing ASSC remain elusive. An improved understanding of the filtration function of the spleen in SCD could advance our knowledge of this early complication of SCD.

Significant efforts have been accumulated to understand the filtration function of splenic IES in blood diseases. Several *ex vivo* [48–53] experiments were conducted through perfusing malaria-infected RBCs and biochemically treated RBCs that resemble the diseased RBCs through the donated human spleen and reported that RBCs with decreased deformability could be mechanically retained at IES. These findings were confirmed by a number of *in vitro* investigations [52, 54–56], where microfluidic chips were devised to mimic the filtration function of IES in blood diseases such as SCD, malaria, and spherocytosis. On the other hand, numerous computational studies have been performed to simulate the dynamics of RBCs passing through the IES and quantify the critical factors that dictate the passage of the RBCs, such as the surface-to-volume ratio of RBCs, the shear and bending modulus of the cells, RBC membrane viscosity, and the size of the IES [57–61]. These simulations have complimented the *ex vivo* and *in vitro* experiments by providing the analysis of the deformation of RBCs, the shear strain and stress on the RBC membrane as well as the potential vesiculation and lysis of RBCs during their traversal through IES, which could not be directly observed from the experiments.

In contrast to the extensive literature on exploring the function of IES in clearing the aged and diseased RBCs in the spleen, there is a lack of studies on quantifying the filtration function

of the splenic macrophages. A quantitative understanding of the role of macrophages in removing the RBCs will open the way to integrate the two major filtration components in the spleen that constantly control the quality of circulating RBCs and advance our knowledge of the underlying mechanism causing the splenic complications in SCD. Thus, in this work, we perform an integrated experimental and computational study to explore macrophage clearance of RBC suspension under physiologically relevant flow conditions in the spleen. We will quantify the influence of the velocity, hematocrit, and various shapes of sickle RBCs on the clearance efficiency under splenic conditions, which will be combined with our current understanding of the filtration function of IES to postulate the complete filtration function of the spleen in SCD.

Materials and methods

Ethics statement

Healthy blood samples were acquired from the local blood bank, while sickle blood samples were obtained from homozygous (HbSS) sickle cell disease (SCD) patients at Massachusetts General Hospital. The collection of sickle blood samples followed the guidelines of an Excess Human Material Protocol, which was approved by the Partners Healthcare Institutional Review Board (IRB) with a waiver of consent. Additional HbSS blood samples were drawn from SCD patients at the University of Pittsburgh, following the University of Pittsburgh IRB protocol PRO08110422. In vitro microfluidic experiments were conducted under an approved exempt protocol (Massachusetts Institute of Technology IRB protocol E-1523).

Computational methods and models

DPD method and cellular level blood cell models. Several computational models of RBCs have been developed in the last two decades to simulate the dynamics of normal and diseased RBCs. Based on their level of complexity, these RBC models can be categorized into the protein-level RBC models [62–69], which are widely used in simulating the pathological alterations of RBC membrane structure in blood disorders, and cellular-level RBC models [70–75], which are mostly used in modeling blood cell suspensions or blood flow. Due to the high computational cost of the protein-level RBC models, we employ a cellular-level model [76] developed based on dissipative particle dynamics (DPD) [77] to simulate the normal and sickle RBCs as well as macrophages (more details on the cellular-level models and the values of model parameters can be found in [S1 Text](#)). The DPD method is a mesoscopic particle-based simulation technique, where each DPD particle represents a lump of molecules and interacts with other particles through soft pairwise forces [78]. DPD can provide the correct hydrodynamic behavior of fluids at the mesoscale, and it has been successfully applied to study complex fluids [79–82]. The formulation of the DPD method can be found in [S1 Text](#).

Stochastic adhesion model. To characterize the initial adhesion between RBCs and macrophages, we employ the stochastic bond formation/dissociation model proposed by Hammer and Apte [83, 84], where the adhesive bond can be formed and dissociated with predefined possibilities P_{on} and P_{off} within a critical length d_{on} and d_{off} , respectively [83] and they can be calculated by

$$P_{on} = 1 - \exp(-k_{on}\Delta t), \quad (1)$$

$$P_{off} = 1 - \exp(-k_{off}\Delta t), \quad (2)$$

where Δt is the timestep of the simulation; k_{on} and k_{off} are the formation and dissociation rates

Table 1. Model parameters for adhesion dynamics between RBCs and macrophages.

Parameters	Simulation	Physical	Reported Range
K_s	50 ~ 150	(2.4 ~ 7.2) $\mu\text{N}/\text{m}$	(1 ~ 10 ⁴) $\mu\text{N}/\text{m}$ [36, 83, 85, 86]
l_0	0.3	0.3 μm	(0.0 ~ 0.2) μm [80, 87]
d_{on}	0.75	0.75 μm	(0.0 ~ 1.0) μm [80, 86, 87]
d_{off}	0.75	0.75 μm	(0.0 ~ 1.0) μm [80, 86, 88]
σ_{on}	10.58	0.51 $\mu\text{N}/\text{m}$	(0.5 ~ 0.6) $\mu\text{N}/\text{m}$ [87]
σ_{off}	3.49	0.17 $\mu\text{N}/\text{m}$	(0.05 ~ 0.75) $\mu\text{N}/\text{m}$ [87]
k_{on}^0	100 ~ 150	(5.42 ~ 8.14) * 10 ⁴ s ⁻¹	(0.001 ~ 1.2) * 10 ⁶ s ⁻¹ [86, 89]
k_{off}^0	0.001 ~ 0.01	0.54 ~ 5.42s ⁻¹	(0.5 ~ 300)s ⁻¹ [90]

<https://doi.org/10.1371/journal.pcbi.1011223.t001>

defined as:

$$k_{on} = k_{on}^0 \exp\left(\frac{-\sigma_{on}(l - l_0)^2}{2K_B T}\right), \quad (3)$$

$$k_{off} = k_{off}^0 \exp\left(\frac{-\sigma_{off}(l - l_0)^2}{2K_B T}\right), \quad (4)$$

where T is the temperature, and k_{on}^0 , k_{off}^0 are the formation and dissociation rates at the equilibrium distance l_0 . σ_{on} and σ_{off} are the effective formation and rupture strengths within a given reactive distance d_{on} and d_{off} , respectively. For a given bond, an adhesion potential energy is given by $U_{adh} = 0.5 \sum_{m=1}^{N_b} K_s (l - l_0)^2$, and the adhesive force between cells and macrophages is calculated by $F_{adh} = -\frac{\partial U_{adh}}{\partial X}$, where K_s is the adhesion strength, N_b is the number of the formed bonds, l is the bond length. The following two conditions are used to determine the formation of a bond

$$l < d_{on} \text{ and } P_{on} > \lambda_1. \quad (5)$$

On the other hand, the following two conditions lead to the dissociation of a bond,

$$l < d_{off} \text{ and } P_{off} > \lambda_2, \quad (6)$$

where the λ_1 and λ_2 are random numbers drawn from a uniform distribution between [0, 1]. The values of the parameters used in the adhesion models and the associated references are summarised in [Table 1](#).

Experimental and simulation setups

Experimental setup. [Fig 1A](#) illustrates the experimental setup and the observation of cell-macrophage adhesion. All experiments were conducted within five days of blood collection. Before the microfluidics experiments, the red blood cells were gently washed twice with phosphate-buffered saline (PBS) solution at 2,000 rpm for 2 min at room temperature (20°C). The hematocrit of the RBC suspensions used within the microfluidic channels is estimated to be 1.0–2%. Before the adhesion experiments, RBCs were resuspended in PBS (2%, v/v) and were opsonized by incubation with 0.5 μM IgG (Rockland, Limerick, PA, USA) at room temperature (20°C). THP-1 macrophages (ATCC, Manassas, VA, USA) were cultured in RPMI media (StemCell Technologies, Cambridge, MA, USA) supplemented with 10% FBS (Sigma-Aldrich, St. Louis, MO, USA), and were differentiated for two days using 100 ng/ml phorbol myristate




Fig 1. Simulation setup for modeling RBC retention by macrophages under Couette flow. A The experiment setup and an observation from the experiment. B The snapshots of the cells retained by macrophages in the experiment (left) and simulation (right). C The geometry of the simulation setup, where the RBC suspension flows past the macrophages, adhered firmly to the bottom of the chip. D Flow streamlines and contours of the velocity field in the simulation. The velocity of the top wall drives the flow, while the bottom wall has zero velocity.

<https://doi.org/10.1371/journal.pcbi.1011223.g001>

acetate (PMA) (StemCell Technologies, Cambridge, MA, USA). To study the adhesion dynamics of sickle RBCs under shear flow and hypoxia, we conducted experiments using a specially developed hypoxic microfluidic device [91]. This device consists of a dual-layer microchannel construction with a gas-permeable polydimethylsiloxane (PDMS) membrane (150 μm thick) and two microchannels: a ‘flow microchannel’ for the RBC suspension and a ‘gas microchannel’ for delivering the desired gas mixture of carbon dioxide (CO_2), oxygen (O_2), and nitrogen (N_2). Deoxygenation and oxygenation conditions of the cells were controlled by switching between two different gas mixtures: 20% O_2 , 5% CO_2 with the balance of N_2 , and an oxygen-poor gas mixture: 2% O_2 , 5% CO_2 with the balance of N_2 , respectively. The flow microchannel, which was 5 mm wide, was connected to a syringe pump (Harvard Apparatus) to control the flow rate (Q) within the microchannel. In this study, the flow rate (Q) ranged from 0 to 20 ml/hr to generate different shear stresses for attaching and detaching the RBCs as required. Imaging of the microfluidic device was carried out using Olympus IX71 inverted microscopes (Olympus, Tokyo, Japan) equipped with Olympus DP72 cameras for image acquisition. All testing was performed at 37°C using a heating incubator (ibidi heating system; ibidi USA,

Fitchburg, WI). The microfluidic device was fabricated by permanent covalent bonding of PDMS channels and the glass substrate following air plasma treatment for 1 minute in a plasma cleaner. The PDMS channels were fabricated by casting customized SU-8/Si master molds with a degassed PDMS mixture of base and curing agent (10:1, w/w) using the soft-lithography technique. The macrophages were loaded into the microfluidic device by gently pipetting the cell suspension into the microfluidic channel. The macrophage viability was confirmed by its morphological evidence of no cytotoxicity or apoptosis (for more details, see [S2 Text](#)).

It is well-known that macrophages display distinct properties when exposed to hypoxic conditions, such as morphology, migration, chemotaxis, adherence to endothelial cells, bacterial killing, differentiation/polarization, and protumorigenic activity [92–95]. In phagocytosis, a prior experimental investigation [96] has shown that hypoxia can enhance the capacity of macrophages to engulf microbeads. Specifically, the phagocytic index is increased by 2, 2.5, and more than 3 folds when the macrophages are treated under hypoxia for 15 mins, 60 mins, and 120 mins before phagocytosis experiments, respectively. Since the current study focuses on investigating the impact of hypoxia on sickle RBCs and their adhesion to macrophages, all the adhesion experiments under hypoxic conditions were conducted within a very short time duration (< 10 minutes) to minimize the impact of hypoxia to macrophages.

Simulation setup. Based on the setup of a typical microfluidic experiment in the left of [Fig 1B](#), we construct a companion microchannel *in silico* as shown in the right of [Fig 1B](#), where the macrophages are placed on the bottom of a microchannel. The geometry of the microchannel in our simulation is constructed as illustrated in [Fig 1C](#), which shows a typical example of simulating the filtration of sickle RBC suspension by a group of macrophages. The height of the microchannel in our simulation is $h = 20\mu\text{m}$, and the width and length are determined by the distribution of macrophages and the region of interest recorded in the experiments. We place the macrophages initially on the bottom of the microchannel. Strong adhesion between the macrophages and the channel bottom wall is assumed to ensure no apparent movement of macrophages in our simulation, consistent with the observations from the microfluidic experiments. Since the focus of the simulation is on the adhesion of RBCs to the macrophages, we only simulate the flow around the macrophages (flow field within $20\mu\text{m}$ from the bottom of the microfluidic channel wall, beyond which these simulated macrophages have no impact on the flowing RBCs or blood flow), instead of the entire channel, to reduce the computational cost. In this case, the height of the simulation domain ($20\mu\text{m}$) is only 4% of the height of the microfluidic channel. As a result, we approximate the flow field in the simulation domain as a Couette flow. [Fig 1D](#) shows the velocity field and streamlines of the flow in the simulation domain. The flow direction in the simulation domain is from left to right, and the top wall's velocity varies to match the experimental conditions. In this work, we consider the adhesion of sickle RBCs under both normoxia and hypoxia. The shear and bending moduli of sickle RBCs under these conditions are summarized in [Table 2](#). The shear modulus and bending modulus of normal RBC are chosen to be $E_{s0} = 4.792\mu\text{N}/\text{m}$ and $E_{b0} = 2.9 * 10^{-19}\text{J}$, following our prior work in [76]. In addition, as observed from the blood samples of SCD patients, four kinds of shaped cells are considered [28], including biconcave, granular,

Table 2. Summary of the typical parameters of RBCs under normoxia and hypoxia used in the simulations.

Conditions	Type	$E_s(\text{N}/\text{m})$	$E_b(\text{J})$	$k_{on}^0(\text{s}^{-1})$	$k_{off}^0(\text{s}^{-1})$	$Ks(\mu\text{N}/\text{m})$
Normoxia	Sickle	$5E_{s0}$	$5E_{b0}$	$5.42 * 10^4$	2.71	2.9
Hypoxia	Sickle	$50E_{s0}$	$50E_{b0}$	$8.14 * 10^4$	0.54	6.7

<https://doi.org/10.1371/journal.pcbi.1011223.t002>

elongated, and sickle shapes. The normal and sickle RBC models consist of 500 vertices, 996 triangles, and 1494 edges, with a surface area $S_R = 132.87\mu\text{m}^2$ and a volume $V_R = 92.46\mu\text{m}^3$. The diameter of macrophages ranges from $16 \sim 22\mu\text{m}$, following the measurements from the experiment, and the height of the macrophages is selected to be $h_M = 11.00\mu\text{m}$. Periodic boundary conditions are imposed in the X and Y directions in the simulation, and bounce-back conditions are applied on the channel wall to prevent fluid particles from penetrating. To align with the experiment's timeframe, we employ an accelerated time scale in simulation, and the detail can be found in [S5 Text](#).

All simulations are computed using the extended version of code developed based on the LAMMPS platform. Each simulation runs for ~ 3000000 time steps. A typical simulation requires 2304 CPU core hours running on the computational resources (Intel Xeon E5–2670 2.6 GHz 24-core processors) at the Center for Computation and Visualization at Brown University.

Calibration of the adhesion model based on microfluidic experiments

First, we calibrate the parameters in the adhesion model by matching our simulation results with the experimental observations under normoxia and hypoxia, as the model parameters are not distinguished based on the oxygenation levels in the existing literature. Based on our observations and measurements from microfluidic experiments, we find both normal and sickle RBCs can adhere to macrophages under flow conditions comparable to those in the red pulp of the spleen ($\sim 150\mu\text{m/s}$). However, the sickle RBCs under hypoxia detach from macrophages at higher flow velocity than normal and sickle RBCs under normoxia, suggesting a strong adhesive force is triggered for hypoxic sickle cells. The critical detachment velocities, measured at the top of the macrophages, are determined based on the various flow conditions. Typically, the normal RBCs and sickle RBCs under normoxia detach from the macrophages at velocities of approximately $500\mu\text{m/s}$ (Experiment [S1 Video](#), Simulation [S2 Video](#)) and $1000\mu\text{m/s}$ (Experiment [S3 Video](#), Simulation [S4 Video](#)), respectively. In comparison, a sickle cell under hypoxia has a detaching velocity greater than $2500\mu\text{m/s}$ (Experiment [S5 Video](#), Simulation [S6 Video](#)).

By matching the simulation results with experimental measurements, we calibrate the key parameters of the stochastic adhesion model, including cell adhesion strength K_s , effective formation strength σ_{on} , rupture strength σ_{off} , bond formation rate k_{on}^0 and bond dissociation rate k_{off}^0 . We note that the effective formation strength σ_{on} and rupture strength σ_{off} have nominal influences on cell adhesion dynamics, so we follow the values used in our previous simulations of sickle RBC adherence to the substrate [85]. The experimental results revealed that, under hypoxic and normoxic conditions, the average number of RBCs retained per macrophage stood at approximately 2.7 and 0.75, respectively. To match these observations, we tune the parameters k_{on}^0 and k_{off}^0 , inspired by our previous study in [85] and find that a higher bond formation rate k_{on}^0 and lower bond break rate k_{off}^0 promote the formation of the bonds between RBCs and macrophages. The parameters k_{on}^0 and k_{off}^0 are chosen from the range of $5.42 * 10^4 - 8.14 * 10^4\text{s}^{-1}$ and $0.54-2.71\text{s}^{-1}$, respectively, following the values reported in the literature. Eventually, we choose unique parameters from the two sets corresponding to various conditions, including $k_{on}^0 = 5.42*10^4\text{s}^{-1}$ and $k_{off}^0 = 2.71\text{s}^{-1}$ for normoxia condition, $k_{on}^0 = 8.14*10^4\text{s}^{-1}$ and $k_{off}^0 = 0.54\text{s}^{-1}$ for hypoxia, the details about the calibration process for k_{on}^0 and k_{off}^0 is described in [S4 Text](#). Then, we investigate the adhesion dynamics of normal and sickle RBCs under both normoxia and hypoxia, through which we infer the cell adhesion strength K_s ,

by increasing the flow velocity until the onset of cell detachment. Specifically, we determine the K_s value by fitting the experimental flow velocity values required for cell detachment. A greater K_s will contribute to a stronger cell adhesion strength, resulting in a firm adhesion and a high flow velocity for inducing RBC detachment from macrophages. In comparison, the adhered RBCs can easily detach from the macrophages with a smaller K_s . The details of how we identify the adhesive strength K_s are described in [S3 Text](#). All the calibrated parameters are summarized in [Table 2](#). The sensitivity analysis of the parameters K_s , k_{on}^0 , and k_{off}^0 under hypoxic conditions is also illustrated in [S4 Text](#). This analysis involved keeping two of the three parameters constant while increasing one to observe its impact on the number of RBCs retained on macrophages. The results reveal that the sensitivity ranking among these parameters was $K_s > k_{on}^0 > k_{off}^0$.

Results

In this section, we integrate computational modeling with microfluidic experiments to quantify the filtrated function of macrophages in microchannels. First, the aggregation of RBCs is introduced to examine the impact of the RBC aggregation force on the macrophage filtration function. Additionally, we investigate the influence of the hematocrit of RBC suspension and blood flow velocity patterns, including constant velocity and step-wise velocities, on the adhesion of sickle RBCs to the macrophages. Finally, we present the effects of the morphology of sickle cells on the filtration by macrophages.

Sickle RBC adhesion to macrophages under normoxia state

[Fig 2A](#) shows three snapshots of sickle cell suspension adhering to the macrophages under normoxia. The top row displays the experimental observations([S7 Video](#)), whereas the bottom row depicts the corresponding simulation results([S8 Video](#)). The configuration of the macrophages in the simulation is derived from the experiments, and the radius range varies from 9 to 11 μm . The velocity around the macrophages is approximately 150 $\mu\text{m}/\text{s}$, and the hematocrit is about 2% in the microchannel. Here, t_1 marks the initial stage with no cell flowing through the macrophages, t_2 represents the snapshot when the first RBC adheres to the macrophages, and t_3 is a snapshot when the number of adhered RBCs reaches a plateau. [Fig 2B](#) shows the geometry of the microchannel, with the blue cells representing the adhered cells to macrophages. We also count the number of cells attached to the macrophages over time in the experiment and the corresponding simulation as illustrated in [Fig 2C](#), which shows that only three cells are detained by the macrophages from the suspension both in the simulation and experiment within the same period.

Sickle RBC adhesion to macrophages under hypoxia state

[Fig 3A](#) shows three snapshots of sickle cell suspension filtrated by the macrophages under hypoxia. The experimental observations([S9 Video](#)) and simulation results([S10 Video](#)) are illustrated at the top and bottom rows, respectively. The configuration of the macrophages in the simulation is reproduced based on the experimental observations, and the radius varies from 8 to 11 μm . The velocity around the macrophages is selected to be 150 $\mu\text{m}/\text{s}$, close to the physiological blood flow velocities in the red pulp of the spleen, and the hematocrit is about 2% in the microchannel. Again, t_1 marks the initial stage with no cell flow through the macrophages, t_2 represents the snapshot when the first RBC adheres to the macrophages, and t_3 is a snapshot when the number of adhered RBCs reaches a plateau. [Fig 3B](#) describes the geometry of the microchannel, with the blue-colored cells highlighting the RBCs attached to the macrophages.




Fig 2. Modeling the sickle RBC retention by macrophages under normoxia. A $t_1 \sim t_3$ shows the three sequential snapshots of RBCs adhering to macrophages from microfluidic experiments (top) and the simulation (bottom). B The microfluidic chip's geometry and the simulation's final snapshot, with the blue-colored cell adhering to the macrophages. C The number of sickle cells retained by the macrophages is recorded over time for microfluidic experiments and simulations.

<https://doi.org/10.1371/journal.pcbi.1011223.g002>

We also count the number of cells adhered to the macrophages over time in the simulations and experiments as illustrated in Fig 3C, where we note that the number of cells adhered to the macrophages increases over time. The number of cells detained by the macrophages under hypoxia is approximately four times higher than that in the case of normoxia.

Aggregation of RBCs promotes their retention by macrophages

RBC aggregation has been observed in low-shear blood flow where a linear stacking of RBCs can form a chain-like structure, or “rouleau” [97]. Under pathological conditions such as type 2 diabetes and SCD, the enhanced RBC adhesion due to increased levels of plasma factors like fibrinogen could lead to more pronounced rouleau formation in patient's blood [98–101], causing blood hyperviscosity in capillaries [102, 103]. Several experimental investigations have been conducted to explore the RBC-RBC aggregation strength using different *in vitro* experimental techniques as well as numerical simulations [104–106]. Despite these aforementioned studies, many important aspects of the influence of cell aggregation on the RBC adhesion to macrophages are still poorly understood. Herein, we apply a Morse potential to sickle RBCs in proximity to generate intercellular adhesive force for simulating the aggregation of sickle




Fig 3. Retention of sickle RBCs by macrophages under hypoxia. A $t_1 \sim t_3$ depicts the snapshots of RBCs adhering to macrophages in microfluidic experiments (top) and simulation (bottom), including the initial (t_1), intermediate (t_2), and final stage (t_3). B The microfluidic chip's geometry and the simulation's final snapshot, with the blue-colored cells highlighting the RBCs firmly adhered to the macrophages. C The number of sickle cells retained by macrophages is recorded over time in microfluidic experiments and simulations.

<https://doi.org/10.1371/journal.pcbi.1011223.g003>

RBCs. This strategy has been used in many previous works for modeling the formation of RBC aggregates [98, 107]. Fig 4(A, top row) depicts a process of sickle RBCs detached by macrophages under hypoxia in the microfluidic channel. The first row shows the three snapshots of the experiment (EXP) (S11 Video). In contrast, the middle and bottom show the simulation results for sickle RBCs with (SIM1) (S12 Video) and without (SIM2) (S13 Video) considering RBC aggregation. In the simulation, the velocity around the macrophages is about $150 \mu\text{m/s}$, and the radius of macrophages is approximately $11 \mu\text{m}$. Here, $t_1 - t_3$ depict the snapshots of the entire computational and experimental process, with t_1 and t_3 representing the snapshot of the first cell retained by the macrophages and the final snapshot when the number of adhered RBCs reaches a plateau. The number of RBCs adhered to the macrophages measured from EXP, SIM1, and SIM2 is plotted in Fig 4B, where we observe that the number of RBCs retained by the macrophages computed from SIM1 is comparable to the results of EXP, and twice that




Fig 4. Investigation of the impact of RBC aggregation on RBC retention by macrophages under hypoxia. A Snapshot of RBCs adhering to macrophages from experiments and simulations. SIM1 represents the simulations considering the aggregation between sickle RBCs, while SIM2 does not. B The plot depicts the number of RBCs adhered to the macrophages over time for experimental and simulation studies. (C-D) The plots describe the two possible adhesion mechanisms that regulate the RBC adhesion to the macrophages (C, cluster filtration) and (D, transitive filtration).

<https://doi.org/10.1371/journal.pcbi.1011223.g004>

of SIM2, indicating the promoting impact of RBCs aggregation to the retention of sickle RBCs by macrophages. Next, we dissect the mechanism of how cell aggregation improves RBC retention. Fig 4C and 4D illustrate two potential mechanisms that distinguish the filtration results in SIM1 and SIM2, namely cluster filtration (Fig 4C) and transitive filtration (Fig 4D). Cluster filtration occurs when an RBC aggregate approaches and adheres to the macrophage. Multiple RBCs in the aggregate make contact and adhere to the macrophages, leading to a higher retention rate. Under the transitive filtration mechanism, the RBCs adhere to the macrophages individually; thus, the resulting RBC retention rate is lower than the cluster filtration. As observed from our microfluidic experiments, both mechanisms may occur in the process of RBC filtration by splenic macrophages, depending on the extent of the RBC-RBC adhesion.

Increased Hct of RBC suspension enhances RBC retention by macrophages

In this section, we will examine the effect of Hct of RBC suspension on macrophage retention. We select the Hct to be $Hct = 1.5\%$, 2.8% , 3.6% , and 5.0% in our simulations, consistent with the Hct we used in the microfluidic experiments. The variation of Hct is achieved by changing the microchannel length, while the number of RBCs in the simulations remains constant. The radius of the macrophages is $11\mu m$, and the velocity is $150\mu m/s$ around the macrophage. Fig 5A shows the snapshots for these four examined Hct when the number of adhered RBCs in the




Fig 5. Investigation of the impact of hematocrit on sickle RBC retention by macrophages under hypoxia. A Snapshots of RBC suspension with various hematocrits (1.5%, 2.8%, 3.6%, 5%) flowing past a macrophage. B The number of RBCs retained by the macrophages is recorded over time under different hematocrits.

<https://doi.org/10.1371/journal.pcbi.1011223.g005>

simulations reaches a plateau. We record the number of RBCs adhered to macrophages over time under different Hct levels in Fig 5B, where we observe that although the number of adhered RBCs in all four cases increases with time, simulations with a greater H_{ct} takes less time to reach the plateau, as a greater H_{ct} could increase the chance of RBCs making contact with macrophages. Additionally, we note that the macrophages retain fewer cells at $H_{ct} = 1.5\%$, likely because a lower H_{ct} diminishes the extent of the RBC interaction that could push RBCs toward macrophages and thus promote RBC adhesion.

Impact of flow velocities on RBC retention by macrophages

Next, we investigate the impact of blood flow velocities on the retention of RBCs by macrophages. Fig 6A depicts the snapshots of RBCs adhering to macrophages at various velocities ranging from $100\text{--}500\mu\text{m/s}$. The radius of the macrophages in the simulations is $11\mu\text{m}$ and the hematocrit is 1.5%. Fig 6B plots the change of average velocity of the RBCs suspension over time. The results illustrate that the average velocity decreases over time due to the retention of RBCs by the macrophages and eventually reaches a saturated baseline. Fig 6C summarizes the number of cells adhered to the macrophages over time. We note that the number of adhered RBCs rapidly approaches the plateau when a greater velocity is applied in the simulation, consistent with the velocity variations in Fig 6B. When the velocity equals $100\mu\text{m/s}$, $200\mu\text{m/s}$, and $300\mu\text{m/s}$, respectively, the total number of adhered cells is the same when the simulations reach equilibrium. In contrast, the total number of adhered RBCs at equilibrium decreases as the velocity is increased to $400\mu\text{m/s}$ and $500\mu\text{m/s}$. These results suggest that the increased blood velocity could attenuate the adhesion of sickle RBCs to the macrophages.

To further explore the effect of velocity on the RBC retention, we implement a setup that integrates simulation (S15 Video) and experiment (S14 Video) to investigate the critical velocity for the RBC adhesion to macrophages using blood velocities regulated in a step-wise manner. The examined velocities are decreased from $1000\mu\text{m/s}$ to $100\mu\text{m/s}$. Fig 7A shows the consecutive snapshots at different velocities. t_1 is the snapshot when the velocity is $v = 1000\mu\text{m/s}$ and no RBCs are retained by the macrophage. When the velocity is decreased to




Fig 6. Investigation of the impact of blood velocities on sickle RBC retention by macrophages under hypoxia. A The snapshots of sickle RBCs suspension flowing past a macrophage under various velocities. B The average velocity of the RBC suspension changes over time during the retaining process. C The number of RBCs retained by the macrophages is recorded over time under various flow velocities.

<https://doi.org/10.1371/journal.pcbi.1011223.g006>

about $v = 500 \mu\text{m/s}$, an RBC is captured by the macrophage at t_2 . As the velocity continuously decreases to around t_3 ($250 \mu\text{m/s}$) and t_4 ($100 \mu\text{m/s}$), an increased number of RBCs adhere to the macrophage. Fig 7B describes the velocity changes over time in the simulation and experiments, as well as the front view of snapshots of the simulation at various velocities. Following the cell suspension simulations, we perform the single-cell adhesive dynamics study under the velocity $v = 150, 500, 750 \mu\text{m/s}$ in Fig 7C to explain the mechanism causing the difference in the RBC retention at different flow velocities. Our results show that when the velocity is greater than $500 \mu\text{m/s}$, the flowing RBC cannot make contact with the macrophage, and they thus cannot be captured and retained by the macrophage. In contrast, a reduced flow velocity will enable contact between the traveling RBCs and the macrophage.

Impact of the shapes of sickle RBCs on their retention by macrophages

Finally, we investigate the effect of the morphology of the sickle RBCs on their retention by macrophages under hypoxia. Following our prior work in [28], four RBC shapes are




Fig 7. The dynamics of sickle RBCs interacting with a macrophage under a blood flow with step-wise changes. A Sequential snapshots of RBCs adhering to macrophages under velocities that are decreased step-wise. Top: microfluidic experiments. Bottom: the companion simulations. B The step-wise velocity changes over time, and the front view of the simulation snapshots captured at velocities of $v = 150, 250, 500, 1000 \mu\text{m/s}$. C The impact of the flow velocity on the trajectories of single RBCs flowing past a macrophage.

<https://doi.org/10.1371/journal.pcbi.1011223.g007>

considered, including biconcave, elongated, granular, and sickle shapes. Guided by the experimental observation in Fig 8A, the radius of the macrophages in the simulation is selected from 8 to $11 \mu\text{m}$. The *Hct* is set to be 2%. Fig 8A shows the snapshots of experiments and simulations with four morphologically altered RBCs. Fig 8B illustrates the changes in the average RBC velocities over time when the four types of RBC suspension flow past the macrophages at a velocity of $150 \mu\text{m/s}$, where we observe continuous decreases of RBC velocities due to the gradual adhesion of RBCs to the macrophage. Fig 8C records the number of RBCs retained by macrophages for five groups of RBC suspension during the simulation, implying a potential ranking of the likelihood of sickle RBC adhesion to the macrophages: sickle > granular > mix > biconcave > elongated. These results suggest that there may be a ranked preference for RBC shapes when macrophages clear sickle RBCs in the spleen.

Discussion and summary

In this work, the retention of sickle RBCs by macrophages by integrating simulations and microfluidic experiments is quantified, aiming to improve our quantitative understanding of the filtration function of macrophages in the red pulp of the spleen. Based on the microfluidic experiments, we calibrate the stochastic adhesion model employed to represent the adhesion between the RBCs and macrophages. We focus on identifying the adhesive forces between the




Fig 8. Investigation of the impact of sickle RBC morphology on their retention by macrophages under hypoxia. A Snapshots of retention of RBCs with various shapes by macrophages. B The average velocity of sickle cells during the retaining process under a bulk flow velocity of $v = 150 \mu\text{m/s}$. C The number of RBCs adhered to the macrophages is recorded over time for sickle RBCs with different shapes.

<https://doi.org/10.1371/journal.pcbi.1011223.g008>

sickle RBCs and macrophages under normoxia and hypoxia, which are missing in the literature. By matching our simulation results with experimental observations, we discover that the adhesive strength between sickle RBCs and macrophages is much more robust under hypoxia than under normoxia. As a result, the number of sickle RBCs retained by the macrophages under hypoxia is approximately four times higher than that under normoxia. Notably, this enhanced adhesion strength results in a firm contact of sickle RBCs to macrophages, which can sustain flow velocities more than ten times higher than the flow velocities in the red pulp of the spleen. Although the underlying molecular mechanism causing the increase in the adhesive strength under hypoxia is not yet clear, our results indicate that the filtration function of the splenic macrophages is upregulated under a more hypoxic environment.

Sickle RBCs under hypoxia are featured with high RBC adhesion that facilitates their adhesion to endothelium cells [108, 109], but also to other sickle RBCs [100, 101], both of which could contribute to the initiation and propagation of vaso-occlusion events in SCD. The microfluidic experiments demonstrate that the formation of RBC aggregates promotes the retention of sickle RBCs by macrophages. The companion simulations we performed further dissect the adhesion process and show that the RBC aggregate adhered to the macrophages

could either directly capture the RBCs in the flow or slow down their motion such that the macrophages could retain them. These findings support the role of RBC aggregation in promoting RBC retention by macrophages. Additionally, we examine the impact of Hct on RBC retention by macrophages. The results show that an increased Hct could boost the chance of RBCs interacting with macrophages, thereby facilitating RBC retention. Considering that the Hct in the red pulp could be as high as 70%, the Hct-induced enhancement of RBC retention could become more pronounced *in vivo*. Although the RBC aggregation and increased Hct promote the retention of RBCs by the macrophages, it is unclear whether the macrophages could efficiently engulf these adhered RBCs. Phagocytosis of RBCs has been primarily studied *in vitro*, and the phagocytic time could vary from minutes to hours [22, 23, 110, 111]. Failure to a timely elimination of these retained RBCs could cause congestion and even ASSC in the red pulp, which has been evidenced by pathological studies of the spleen of SCD patients [42, 58].

We also quantify the impact of blood flow velocities on the retention of RBCs by macrophages by adjusting step-wise the blood flow velocities in the microchannel. The simulation results show that when the flow velocity is varied between 100–300 $\mu\text{m/s}$, which are comparable to the blood velocities in the red pulp, no notable changes occur to the total number of retained RBCs. However, as velocity continues to increase, the number of the retained cells will decrease rapidly, implying that the increased blood velocity could attenuate the function of the red pulp macrophages on detaining aged or diseased RBCs. This finding provides a possible rationale for the slow blood flow in the open circulation of the spleen in facilitating the filtering of RBCs by red pulp macrophages. Our simulations of sickle RBCs with various shapes show shape preference when sickle RBCs are susceptible to retention by macrophages, and the preferential shapes rank in the following order: sickle>granular>mix>biconcave >elongated. Our results in the current work suggest that the sickle and granular-shaped RBCs are more likely to be retained in the spleen. This finding is consistent with the observation in the literature that the sickle and granular-shaped RBCs are less frequently observed than the discoid or elongated form of RBCs in the blood smear of SCD patients [28].

We note some limitations in our experimental results and simulations. In the microfluidic experiments, diluted blood samples from SCD patients (Hct \leq 5%) were used to observe the adhesion dynamics of sickle RBCs to the macrophages to quantify the number of adhered RBCs and reduce the risk of clogging the microfluidic channels. This value of Hct used is much smaller than the physiological value of \sim 70% in the red pulp of the spleen. This could lead to an underestimation of the function of macrophages in retaining the sickle RBCs. Moreover, we did not consider the macrophages' engulfment of the adhered RBCs, as the phagocytic time is much longer than the simulation time required for forming RBC-macrophage adhesion. We note that modeling the phagocytosis process requires more detailed modeling of intracellular components, such as the polymerization of actin to filaments and the interaction between the actin filaments and macrophage membrane, which is beyond the capability of the models employed in the current study. A general kinetic cell adhesion model proposed by Bell and Dembo [112, 113] is used to mimic the collective effect of ligand-receptor binding between the sickle RBCs and macrophages. This model cannot distinguish the contribution from various ligands, such as the band-3 proteins [16, 17], exposure of PS [18, 19], decreased expression of CD47 [20, 21] and conformational changes in CD47 [21].

In summary, we have employed mesoscopic computational models that are validated by microfluidic experiments to simulate the adhesion of sickle RBCs to macrophages, aiming to infer the filtration function of the red pulp macrophages. This important spleen function has not been well-studied. We quantified the impact of several key factors that could dictate RBC retention by splenic macrophages, such as blood flow conditions, RBC aggregation, Hct, RBC

morphology, and particularly oxygen levels, which have not been systematically explored. Our experimental and simulation results improve our quantitative understanding of the filtration function of splenic macrophages and provide an opportunity to combine the new findings with the current knowledge of the filtration function of IES to apprehend the complete filtration function of the spleen in SCD.

Supporting information

S1 Text. Hydrodynamics and particle based blood cell models.

(PDF)

S2 Text. Details of experiments.

(PDF)

S3 Text. Calibrate the adhesion strength K_s based on microfluidic experiments.

(PDF)

S4 Text. Calibrate the bond formation and dissociation k_{on}^0, k_{off}^0 and the sensitivity analysis of the parameters.

(PDF)

S5 Text. Accelerated time sale.

(PDF)

S1 Video. Experimental observation of detachment dynamics of single normal cell under the normoxia.

(MP4)

S2 Video. Simulation observation of detachment dynamics of single normal cell under the normoxia.

(MP4)

S3 Video. Experimental observation of detachment dynamics of paired sickle cells under the normoxia.

(MP4)

S4 Video. Simulation observation of detachment dynamics of paired sickle cells under the normoxia.

(MP4)

S5 Video. Experimental observation of detachment dynamics of single sickle cell under the hypoxia.

(MP4)

S6 Video. Simulation observation of detachment dynamics of single sickle cell under the hypoxia.

(MP4)

S7 Video. Experimental observations of The sickle RBCs suspension filtrated by macrophages under normoxia.

(MP4)

S8 Video. Computational observations of The sickle RBCs suspension filtrated by macrophages under normoxia.

(MP4)

S9 Video. The experimental observations of The sickle RBCs suspension filtrated by macrophages under hypoxia.

(MP4)

S10 Video. The computational observations of The sickle RBCs suspension filtrated by macrophages under hypoxia.

(MP4)

S11 Video. The experimental observations of the impact of cell-cell interaction on the filtration of sickle cell suspension by macrophages with cell-cell interaction under hypoxia.

(MP4)

S12 Video. The computational observations of the impact of cell-cell interaction on the filtration of sickle cell suspension by macrophages with cell-cell interaction under hypoxia.

Simulation video without cell-cell interaction.

(MP4)

S13 Video. The computational observations of the impact of cell-cell interaction on the filtration of sickle cell suspension by macrophages with cell-cell interaction under hypoxia.

(MP4)

S14 Video. The experimental observations of the filtration of sickle cell by macrophages at continuously varying velocity under hypoxia.

(MP4)

S15 Video. The computational observations of the filtration of sickle cell by macrophages at continuously varying velocity under hypoxia.

(MP4)

Acknowledgments

High Performance Computing resources were provided by the Center for Computation and Visualization at Brown University.

Author Contributions

Conceptualization: Guansheng Li, He Li, George Em Karniadakis.

Data curation: Guansheng Li, Yuhao Qiang.

Formal analysis: Guansheng Li, Yuhao Qiang, He Li, Xuejin Li, Pierre A. Buffet, Ming Dao, George Em Karniadakis.

Funding acquisition: George Em Karniadakis.

Investigation: Guansheng Li, Yuhao Qiang, He Li.

Methodology: Guansheng Li.

Project administration: George Em Karniadakis.

Software: Guansheng Li, He Li, Xuejin Li, George Em Karniadakis.

Supervision: George Em Karniadakis.

Validation: Guansheng Li.

Visualization: Guansheng Li.

Writing – original draft: Guansheng Li.

Writing – review & editing: Guansheng Li, Yuhao Qiang, He Li, Xuejin Li, Pierre A. Buffet, Ming Dao, George Em Karniadakis.

References

1. Kraal G. Cells in the marginal zone of the spleen. *International Review of Cytology* 1992; 132:31–74. [https://doi.org/10.1016/S0074-7696\(08\)62453-5](https://doi.org/10.1016/S0074-7696(08)62453-5) PMID: 1555921
2. Mebius RE, Kraal G. Structure and function of the spleen. *Nature Reviews Immunology* 2005; 5:606–616. <https://doi.org/10.1038/nri1669> PMID: 16056254
3. Li H, Liu ZL, Lu L, Buffet P, Karniadakis GE. How the spleen reshapes and retains young and old red blood cells: A computational investigation. *PLoS Computational Biology* 2021; 17:e1009516. <https://doi.org/10.1371/journal.pcbi.1009516> PMID: 34723962
4. Groom AC. Microcirculation of the spleen: new concepts, new challenges. *Microvascular Research* 1987; 34:269–289. [https://doi.org/10.1016/0026-2862\(87\)90061-6](https://doi.org/10.1016/0026-2862(87)90061-6) PMID: 3323854
5. Matsuno K, Ezaki T, Kotani M. Splenic outer periarterial lymphoid sheath (PALS): an immunoproliferative microenvironment constituted by antigen-laden marginal metallophilic and ED2-positive macrophages in the rat. *Cell and Tissue Research* 1989; 257:459–470. PMID: 2790931
6. Förster R, Schubel A, Breitfeld D, Kremmer E, Renner-Müller I, Wolf E et al. CCR7 Coordinates the Primary Immune Response by Establishing Functional Microenvironments in Secondary Lymphoid Organs. *Cell* 1997; 99:23–33.
7. Ansel KM, Ngo VN, Hyman PL, Luther SA, Förster R, Sedgwick JD et al. A chemokine-driven positive feedback loop organizes lymphoid follicles. *Nature*. 2000; 406:309–314. <https://doi.org/10.1038/35018581> PMID: 10917533
8. Schmidt EE, MacDonald IC, Groom AC. Comparative aspects of splenic microcirculatory pathways in mammals: the region bordering the white pulp. *Scanning Microscopy*. 1993; 7:613–628. PMID: 8108677
9. Saito H, Yokoi Y, Watanabe S, Tajima J, Kuroda H, Namihisa T. Reticular meshwork of the spleen in rats studied by electron microscopy. *American Journal of Anatomy*. 1998; 181:235–252. <https://doi.org/10.1002/aja.1001810303>
10. Groom AC, Schmidt EE, MacDonald IC. Microcirculatory pathways and blood flow in spleen: new insights from washout kinetics, corrosion casts, and quantitative intravital videomicroscopy. *Scanning Microscopy*. 1991; 5:159–174. PMID: 2052921
11. Chen L, Weiss L. The role of the sinus wall in the passage of erythrocytes through the spleen. *Blood*. 1973; 41:529–537. <https://doi.org/10.1182/blood.V41.4.529.529> PMID: 4688868
12. Chen L, Weiss L. Electron microscopy of the red pulp of human spleen. *American Journal of Anatomy*. 1972; 134:425–457. <https://doi.org/10.1002/aja.1001340404> PMID: 5047597
13. Weiss L. A scanning electron microscopic study of the spleen. *Blood*. 1974; 43:665–691. <https://doi.org/10.1182/blood.V43.5.665.665> PMID: 4821401
14. Duez J, Holleran JP, Ndour PA, Pionneau C, Diakité S, Roussel C et al. Mechanical clearance of red blood cells by the human spleen: Potential therapeutic applications of a biomimetic RBC filtration method. *Transfusion Clinique et Biologique*. 2015; 22:151–157 <https://doi.org/10.1016/j.tracli.2015.05.004> PMID: 26138907
15. Mohanty JG, Nagababu E, Rifkind JM. Red blood cell oxidative stress impairs oxygen delivery and induces red blood cell aging. *Frontiers in Physiology* 2014; 5:84. <https://doi.org/10.3389/fphys.2014.00084> PMID: 24616707
16. Arese P, Turrini F, Schwarzer E. Band 3/complement-mediated recognition and removal of normally senescent and pathological human erythrocytes. *Cellular Physiology and Biochemistry* 2005; 16:133–146. <https://doi.org/10.1159/000089839> PMID: 16301814
17. Lutz H. Innate immune and non-immune mediators of erythrocyte clearance. *Cellular and Molecular Biology* 2004; 50:107–116. PMID: 15095782
18. Boas FE, Forman L, Beutler E. Phosphatidylserine exposure and red cell viability in red cell aging and in hemolytic anemia. *Proceedings of the National Academy of Sciences* 1998; 95:3077–3081. <https://doi.org/10.1073/pnas.95.6.3077> PMID: 9501218
19. Connor J, Pak CC, Schroit AJ. Exposure of phosphatidylserine in the outer leaflet of human red blood cells. Relationship to cell density, cell age, and clearance by mononuclear cells. *Journal of Biological Chemistry* 1994; 269:2399–2404. [https://doi.org/10.1016/S0021-9258\(17\)41959-4](https://doi.org/10.1016/S0021-9258(17)41959-4) PMID: 8300565

20. Lv Z, Bian Z, Shi L, Niu S, Ha B, Tremblay A et al. Loss of cell surface CD47 clustering formation and binding avidity to SIRP α facilitate apoptotic cell clearance by macrophages. *The Journal of Immunology* 2015; 195:661–671. <https://doi.org/10.4049/jimmunol.1401719> PMID: 26085683
21. Burger P, Hilarius-Stokman P, Korte D, Berg TK., Bruggen BV. CD47 functions as a molecular switch for erythrocyte phagocytosis. *Blood* 2012; 119:5512–5521. <https://doi.org/10.1182/blood-2011-10-386805> PMID: 22427202
22. Sosale NG, Rouhiparkouhi T, Bradshaw AM, Dimova R, Lipowsky R, Discher DE. Cell rigidity and shape override CD47's "self"-signaling in phagocytosis by hyperactivating myosin-II. *Blood* 2015; 125:542–552. <https://doi.org/10.1182/blood-2014-06-585299> PMID: 25411427
23. Klei TRL, Dalimot J, Nota B, Veldhuis M, Mul FPJ, Rademakers T et al. Hemolysis in the spleen drives erythrocyte turnover. *Blood* 2020; 136:1579–1589. PMID: 32777816
24. Pauling L, Itano HA, Singer SJ, Wells IC. Sickle cell anemia. *Science* 1949; 110:543–8. <https://doi.org/10.1126/science.110.2865.543> PMID: 15395398
25. Ingram VM. Gene mutations in human haemoglobin: the chemical difference between normal and sickle cell haemoglobin. *Nature* 1957; 180(4581):326–328. <https://doi.org/10.1038/180326a0> PMID: 13464827
26. Noguchi CT, Schechter AN. The intracellular polymerization of sickle hemoglobin and its relevance to sickle cell disease. *Blood* 1981; 58:1057–1068. <https://doi.org/10.1182/blood.V58.6.1057> PMID: 7030432
27. Eaton WA, Hofrichter J. Sickle cell hemoglobin polymerization. *Advances in Protein Chemistry* 1990; 40:63–279. [https://doi.org/10.1016/S0065-3233\(08\)60287-9](https://doi.org/10.1016/S0065-3233(08)60287-9) PMID: 2195851
28. Xu M, Papageorgiou DP, Abidi SZ, Dao M, Zhao H, Karniadakis GE. A deep convolutional neural network for classification of red blood cells in sickle cell anemia. *PLoS Computational Biology* 2017; 13(10):e1005746. <https://doi.org/10.1371/journal.pcbi.1005746> PMID: 29049291
29. Kaul DK, Fabry ME, Nagel RL. Microvascular sites and characteristics of sickle cell adhesion to vascular endothelium in shear flow conditions: pathophysiological implications. *Proceedings of the National Academy of Sciences* 1989; 86(9):3356–3360. <https://doi.org/10.1073/pnas.86.9.3356> PMID: 2497465
30. Manwani D, Bieker JJ. The erythroblastic island. *Current Topics in Developmental Biology* 2008; 82:23–53. [https://doi.org/10.1016/S0070-2153\(07\)00002-6](https://doi.org/10.1016/S0070-2153(07)00002-6) PMID: 18282516
31. Papageorgiou DP, Abidi SZ, Chang H, Li X, Kato GJ, Karniadakis GE et al. Simultaneous polymerization and adhesion under hypoxia in sickle cell disease. *Proc. Natl. Acad. Sci.* 2018; 115:9473–9478. <https://doi.org/10.1073/pnas.1807405115> PMID: 30190429
32. Pizzi M, Fuligni F, Santoro L, Sabattini E, Ichino M, De Vito R et al. Spleen histology in children with sickle cell disease and hereditary spherocytosis: hints on the disease pathophysiology. *Human Pathology* 2017; 60:95–103. <https://doi.org/10.1016/j.humpath.2016.09.028> PMID: 27771375
33. Topley JM, Rogers DW, Stevens MC, Serjeant GR. Acute splenic sequestration and hypersplenism in the first five years in homozygous sickle cell disease. *Archives of Disease in Childhood* 1981; 56:765–769. <https://doi.org/10.1136/adc.56.10.765> PMID: 7305414
34. Emond AM, Collis R, Darvill D, Higgs DR, Maude GH, Serjeant GR. Acute splenic sequestration in homozygous sickle cell disease: natural history and management. *The Journal of Pediatrics* 1985; 107:201–206. [https://doi.org/10.1016/S0022-3476\(85\)80125-6](https://doi.org/10.1016/S0022-3476(85)80125-6) PMID: 4020541
35. Warkentin TE, Barr RD, Ali MA, Mohandas N. Recurrent acute splenic sequestration crisis due to interacting genetic defects: hemoglobin SC disease and hereditary spherocytosis. *Blood* 1990; 75:266–270. <https://doi.org/10.1182/blood.V75.1.266> PMID: 2294990
36. Ye T, Zhang X, Li G, Wang S. Biomechanics in thrombus formation from direct cellular simulations. *Phys. Rev. E.* 2020; 102:042410. <https://doi.org/10.1103/PhysRevE.102.042410> PMID: 33212741
37. Brousse V, Elie C, Benkerrou M, Odièvre M, Lesprit E, Bernaudin F et al. Acute splenic sequestration crisis in sickle cell disease: cohort study of 190 paediatric patients. *British Journal of Haematology* 2012; 156:643–648. <https://doi.org/10.1111/j.1365-2141.2011.08999.x> PMID: 22224796
38. Hoss SE, Cochet S, Marin M, Lapoumériou C, Dussiot M, Bouazza N et al. Insights into determinants of spleen injury in sickle cell anemia. *Blood Advances* 2019; 3:2328–2336. <https://doi.org/10.1182/bloodadvances.2019000106> PMID: 31391165
39. Al-Salem AH. Indications and complications of splenectomy for children with sickle cell disease. *Journal of Pediatric Surgery* 2006; 41(11):1909–1915. <https://doi.org/10.1016/j.jpedsurg.2006.06.020> PMID: 17101369
40. Hoss SE, Brousse V. Considering the spleen in sickle cell disease. *Expert Review of Hematology* 2019; 12:563–573. <https://doi.org/10.1080/17474086.2019.1627192> PMID: 31195851
41. Howard JO, Telfer P. *The Spleen. Sickle Cell Disease in Clinical Practice*, 167–172, 2015. Springer.

42. Brousse V, Buffet P, Rees D. The spleen and sickle cell disease: the sick (led) spleen. *British Journal of Haematology* 2014; 166(2):165–176. Wiley Online Library. <https://doi.org/10.1111/bjh.12950> PMID: 24862308
43. Serjeant GR. Management of sickle cell disease: challenges and risks of transfusion. *International Journal of Clinical Transfusion Medicine* 2016; 4:109–119. Dove Press. <https://doi.org/10.2147/IJCTM.S96918>
44. Larsson F, Åhlin A, Marshall MH, Abrahamsson J. Acute splenic sequestration in children with sickle cell disease—an overview. *Lakartidningen* 2016; 113. PMID: 28026839
45. Ibrahimova A, Bernstein B, Abid R, Apollonsky N. The Effect of Splenectomy on Development of Cerebral Vasculopathy in Children with Sickle Cell Disease. *Blood* 2016; 18:3647. Am Soc Hematology. <https://doi.org/10.1182/blood.V128.22.3647.3647>
46. Ware RE, de Montalembert M, Tshilolo L, Abboud MR. Sickle cell disease. *The Lancet* 2017; 390(10091):311–323. Elsevier. [https://doi.org/10.1016/S0140-6736\(17\)30193-9](https://doi.org/10.1016/S0140-6736(17)30193-9)
47. Owusu-Ofori S, Remington T. Splenectomy versus conservative management for acute sequestration crises in people with sickle cell disease. *The Cochrane Library* 2017.
48. Buffet PA, Milon G, Brousse V, Correas J, Dousset B, Couvelard A et al. Ex vivo perfusion of human spleens maintains clearing and processing functions. *Blood* 2006; 107:3745–3752. <https://doi.org/10.1182/blood-2005-10-4094> PMID: 16384927
49. Safeukui I, Correas J, Brousse V, Hiirt D, Deplaine G, Mulé S et al. Retention of Plasmodium falciparum ring-infected erythrocytes in the slow, open microcirculation of the human spleen. *Blood* 2008; 112:2520–2528. <https://doi.org/10.1182/blood-2008-03-146779> PMID: 18579796
50. Safeukui I, Buffet PA, Deplaine G, Perrot S, Brousse V, Ndour A et al. Quantitative assessment of sensing and sequestration of spherocytic erythrocytes by the human spleen. *Blood* 2012; 120:424–430. <https://doi.org/10.1182/blood-2012-01-404103> PMID: 22510876
51. Deplaine G, Safeukui I, Jeddi F, Lacoste F, Brousse V, Perrot S et al. The sensing of poorly deformable red blood cells by the human spleen can be mimicked in vitro. *Blood* 2011; 117:e88–e95. Am Soc Hematology. <https://doi.org/10.1182/blood-2010-10-312801> PMID: 21163923
52. Safeukui I, Buffet PA, Deplaine G, Perrot S, Brousse V, Sauvanet A et al. Sensing of red blood cells with decreased membrane deformability by the human spleen. *Blood Advances*, 2, 2581–2587, 2018, American Society of Hematology Washington, DC.
53. Li G, Qiang Y, Li H, Li X, Dao M, Karniadakis GE. In silico and in vitro study of the adhesion dynamics of erythrophagocytosis in sickle cell disease. *Biophysical Journal*. 2023; 122:P2590–2604. <https://doi.org/10.1016/j.bpj.2023.05.022> PMID: 37231647
54. Picot J, Ndour PA, Lefevre SD, El NW, Tawfik H, Galimand J et al. A biomimetic microfluidic chip to study the circulation and mechanical retention of red blood cells in the spleen. *American Journal of Hematology*, 90, 339–345, 2015. <https://doi.org/10.1002/ajh.23941> PMID: 25641515
55. Gambhire P, Atwell S, Iss C, Bedu F, Ozerov I, Badens C et al. High Aspect Ratio Sub-Micrometer Channels Using Wet Etching: Application to the Dynamics of Red Blood Cell Transiting through Biomimetic Splenic Slits. *PLoS Comput. Biol.*, 13, 1700967, 2017.
56. Du E, Diez-Silva M, Kato GJ, Dao M, Suresh S. Kinetics of sickle cell biorheology and implications for painful vasoocclusive crisis. *Proceedings of the National Academy of Sciences*, 112(5), 1422–1427, 2015. <https://doi.org/10.1073/pnas.1424111112> PMID: 25605910
57. Qi X, Wang S, Ma S, Han K, Li X. Quantitative prediction of flow dynamics and mechanical retention of surface-altered red blood cells through a splenic slit. *Physics of Fluids*, 33, 051902, 2021. <https://doi.org/10.1063/5.0050747>
58. Qiang Y, Sissoko A, Liu ZL, Dong T, Zheng F, Kong F et al. Theoretical model of efficient phagocytosis driven by curved membrane proteins and active cytoskeleton forces. *Proc. Natl. Acad. Sci.*, 120, e2217607120, 2022. <https://doi.org/10.1073/pnas.2217607120>
59. Lu H, Peng Z. Boundary integral simulations of a red blood cell squeezing through a submicron slit under prescribed inlet and outlet pressures. *Physics of Fluids*, 31, 031902, 2019. <https://doi.org/10.1063/1.5081057>
60. Salehyar S, Zhu Q. Deformation and internal stress in a red blood cell as it is driven through a slit by an incoming flow. *Soft Matter*, 12, 3156–3164, 2016. <https://doi.org/10.1039/C5SM02933C> PMID: 26865054
61. Pivkin IV, Peng Z, Karniadakis GE, Buffet PA, Dao M, Suresh S. Biomechanics of red blood cells in human spleen and consequences for physiology and disease. *Proceedings of the National Academy of Sciences*, 113, 7804–7809, 2016. <https://doi.org/10.1073/pnas.1606751113> PMID: 27354532
62. Zhang P, Zhang L, Slepian MJ, Deng Y, Bluestein D. A multiscale biomechanical model of platelets: Correlating with in-vitro results. *Journal of Biomechanics*. 2017; 50:26–33. <https://doi.org/10.1016/j.jbiomech.2016.11.019> PMID: 27894676

63. Li H, Yang J, Chu TT, Naidu R, Lu L, Chandramohanadas R et al. Cytoskeleton remodeling induces membrane stiffness and stability changes of maturing reticulocytes. *Biophysical Journal*. 2018; 114:2014–2023. <https://doi.org/10.1016/j.bpj.2018.03.004> PMID: 29694877
64. Chang H, Li X, Li H, Karniadakis GE. MD/DPD multiscale framework for predicting morphology and stresses of red blood cells in health and disease. *PLoS Computational Biology*. 2016; 12:e1005173. <https://doi.org/10.1371/journal.pcbi.1005173> PMID: 27792725
65. Dearnley M, Chu T, Zhang Y, Looker O, Huang C, Klonis N et al. Reversible host cell remodeling underpins deformability changes in malaria parasite sexual blood stages. *Proceedings of the National Academy of Sciences*. 2016; 113:4800–4805. <https://doi.org/10.1073/pnas.1520194113> PMID: 27071094
66. Li G, Ye T, Li X. Parallel modeling of cell suspension flow in complex micro-networks with inflow/outflow boundary conditions. *Journal of Computational Physics*. 2020; 401:109031. <https://doi.org/10.1016/j.jcp.2019.109031>
67. Li G, Ye T, Wang S, Li X, Haq UR. Numerical design of a highly efficient microfluidic chip for blood plasma separation. *Physics of Fluids*. 2020; 32:031903. <https://doi.org/10.1063/1.5144436>
68. Li G, Ye T, Yang B, Wang S, Li X. Temporal-spatial heterogeneity of hematocrit in microvascular networks. *Physics of Fluids*. 2023; 35:021906. <https://doi.org/10.1063/5.0139641>
69. Dorken-Gallastegi A, Lee Y, Li G, Li H, Naar L, Li X et al. Circulating Cellular Clusters Are Associated with Thrombotic Complications and Clinical Outcomes in COVID-19. *iScience*. 2023; 26:107202. <https://doi.org/10.1016/j.isci.2023.107202> PMID: 37485375
70. Geekiyanage NM, Balanant MA, Sauret E, Saha S, Flowe R, Lim CT et al. A coarse-grained red blood cell membrane model to study stomatocyte-discocyte-echinocyte morphologies. *PLoS One*. 2019; 14:e0215447. <https://doi.org/10.1371/journal.pone.0215447> PMID: 31002688
71. Geekiyanage N, Sauret E, Saha S, Flower R, Gu Y. Modelling of red blood cell morphological and deformability changes during in-vitro storage. *Applied Sciences*. 2020; 10:3209. <https://doi.org/10.3390/app10093209>
72. Yazdani A, Deng Y, Li H, Javadi E, Li Z, Jamali S et al. Integrating blood cell mechanics, platelet adhesive dynamics and coagulation cascade for modelling thrombus formation in normal and diabetic blood. *Journal of the Royal Society Interface*. 2021; 18:20200834. <https://doi.org/10.1098/rsif.2020.0834> PMID: 33530862
73. Czaja B, Bouter JD, Heisler M, Závodszy G, Karst S, Sarunic M et al. The effect of stiffened diabetic red blood cells on wall shear stress in a reconstructed 3D microaneurysm. *Computer Methods in Biomechanics and Biomedical Engineering*. 2022:1–19.
74. Závodszy G, Rooij BV, Azizi V, Hoekstra A. Cellular level in-silico modeling of blood rheology with an improved material model for red blood cells. *Frontiers in Physiology*. 2017; 8:563. <https://doi.org/10.3389/fphys.2017.00563> PMID: 28824458
75. Ma S, Wang S, Qi X, Han K, Jin X, Li Z et al. Multiscale computational framework for predicting viscoelasticity of red blood cells in aging and mechanical fatigue. *Computer Methods in Applied Mechanics and Engineering*. 2022; 391:114535. <https://doi.org/10.1016/j.cma.2021.114535>
76. Fedosov D A, Caswell B, Karniadakis GE. A multiscale red blood cell model with accurate mechanics, rheology, and dynamics. *Biophysical Journal*. 2010; 98:2215–2225. <https://doi.org/10.1016/j.bpj.2010.02.002> PMID: 20483330
77. Español P. Dissipative particle dynamics with energy conservation. *Europhys. Lett*. 1997; 40:631.
78. Groot RD, Warren PB. Dissipative particle dynamics: Bridging the gap between atomistic and mesoscopic simulation. *The Journal of Chemical Physics*. 1997; 107:4423–4435. <https://doi.org/10.1063/1.474784>
79. Li G, Ye T, Xia Z, Wang S, Zhu Z. Analysis and prediction of hematocrit in microvascular networks. *International Journal of Engineering Science*. 2023; 191:103901. <https://doi.org/10.1016/j.ijengsci.2023.103901>
80. Lei H, Karniadakis GE. Probing vasoocclusion phenomena in sickle cell anemia via mesoscopic simulations. *Proc. Natl. Acad. Sci*. 2013; 110:11326–11330. <https://doi.org/10.1073/pnas.1221297110> PMID: 23798393
81. Chang H, Yazdani A, Li X, Douglas K, Mantzoros CS, Karniadakis GE. Quantifying platelet margination in diabetic blood flow. *Biophysical Journal*. 2018; 115:1371–1382. <https://doi.org/10.1016/j.bpj.2018.08.031> PMID: 30224049
82. Han K, Ma S, Sun J, Xu M, Qi X, Wang S et al. In silico modeling of patient-specific blood rheology in type 2 diabetes mellitus. *Biophysical Journal*. 2023; 122(8):1445–1458. <https://doi.org/10.1016/j.bpj.2023.03.010> PMID: 36905122

83. Hammer DA, Apte SM. Simulation of cell rolling and adhesion on surfaces in shear flow: general results and analysis of selectin-mediated neutrophil adhesion. *Biophys. J.* 1992; 63:35–57. [https://doi.org/10.1016/S0006-3495\(92\)81577-1](https://doi.org/10.1016/S0006-3495(92)81577-1) PMID: 1384734
84. Fedosov DA, Caswell B, Karniadakis GE. Wall shear stress-based model for adhesive dynamics of red blood cells in malaria. *Biophys. J.* 2011; 9:2084–2093. <https://doi.org/10.1016/j.bpj.2011.03.027> PMID: 21539775
85. Deng Y, Papageorgiou DP, Chang H, Abidi SZ, Li X, Dao M et al. Quantifying Shear-Induced Deformation and Detachment of Individual Adherent Sickle Red Blood Cells. *Biophys. J.* 2019; 116:360–371. <https://doi.org/10.1016/j.bpj.2018.12.008> PMID: 30612714
86. Wang S, Ye T, Li G, Zhang X, Shi H. Margination and adhesion dynamics of tumor cells in a real microvascular network. *PLoS Comput. Biol.* 2021; 17:e1008746. <https://doi.org/10.1371/journal.pcbi.1008746> PMID: 33606686
87. Fedosov DA, Caswell B, Suresh S, Karniadakis GE. Quantifying the biophysical characteristics of Plasmodium-falciparum-parasitized red blood cells in microcirculation. *Proc. Natl. Acad. Sci.* 2011; 108:35–39. <https://doi.org/10.1073/pnas.1009492108> PMID: 21173269
88. Marshall BT, Sarangapani KK, Wu J, Lawrence MB, McEver RP, Zhu C. Measuring Molecular Elasticity by Atomic Force Microscope Cantilever Fluctuations. *Biophys. J.* 2006; 90:681–692. <https://doi.org/10.1529/biophysj.105.061010> PMID: 16258054
89. Ulrich SS, Ronen A. L-selectin-mediated leukocyte tethering in shear flow is controlled by multiple contacts and cytoskeletal anchorage facilitating fast rebinding events. *Proc. Natl. Acad. Sci.* 2004; 101:6940–6945. <https://doi.org/10.1073/pnas.0305822101>
90. Ronen A, Chen S, Puri KD, Finger EB, Springer TA. The kinetics of L-selectin tethers and the mechanics of selectin-mediated rolling. *J. Cell Biol.* 1997; 138:1169–1180. <https://doi.org/10.1083/jcb.138.5.1169>
91. Qiang Y, Liu J, Dao M, Du E. In vitro assay for single-cell characterization of impaired deformability in red blood cells under recurrent episodes of hypoxia. *Lab on a Chip.* 2021; 21:3458–3470. <https://doi.org/10.1039/d1lc00598g> PMID: 34378625
92. Egners A, Erdem M, Cramer T. The response of macrophages and neutrophils to hypoxia in the context of cancer and other inflammatory diseases. *Mediators of Inflammation.* 2016; 2016. <https://doi.org/10.1155/2016/2053646> PMID: 27034586
93. Silva VL. Exploiting the cancer niche: tumor-associated macrophages and hypoxia as promising synergistic targets for nano-based therapy. *Journal of Controlled Release.* 2017; 253:82–96. <https://doi.org/10.1016/j.jconrel.2017.03.013> PMID: 28285930
94. Acosta-Iborra B, Elorza A, Olazabal I. Macrophage oxygen sensing modulates antigen presentation and phagocytic functions involving IFN- γ production through the HIF-1 α transcription factor. *The Journal of Immunology.* 2009; 182(5):3155–3164. <https://doi.org/10.4049/jimmunol.0801710> PMID: 19234213
95. Bai R, Li Y, Jian L. The hypoxia-driven crosstalk between tumor and tumor-associated macrophages: mechanisms and clinical treatment strategies. *Molecular Cancer.* 2022; 21(1):177. <https://doi.org/10.1186/s12943-022-01645-2> PMID: 36071472
96. Anand RJ, Gripar SC, Li J. Hypoxia causes an increase in phagocytosis by macrophages in a HIF-1 α -dependent manner. *Journal of Leukocyte Biology.* 2007; 82(5):1257–1265. <https://doi.org/10.1189/jlb.0307195> PMID: 17675562
97. Bäuml H, Neu B, Donath E, Kiesewetter H. Basic phenomena of red blood cell rouleaux formation. *Biorheology.* 1999; 36:439–442. PMID: 10818642
98. Deng Y, Papageorgiou DP, Li X. Quantifying fibrinogen-dependent aggregation of red blood cells in type 2 diabetes mellitus. *Biophysical Journal.* 2020; 119:900–912. <https://doi.org/10.1016/j.bpj.2020.07.026> PMID: 32814061
99. Cloutier G, Zimmer A, Yu FT, Chiasson J. Increased shear rate resistance and fastest kinetics of erythrocyte aggregation in diabetes measured with ultrasound. *Diabetes Care.* 2008; 31(7):1400–1402. <https://doi.org/10.2337/dc07-1802> PMID: 18375419
100. Obiefuna PC, Photiades DP. Sick cell discocytes form more rouleaux in vitro than normal erythrocytes. *The Journal of Tropical Medicine and Hygiene.* 1990; 93(3):210–214. PMID: 2348499
101. Obiefuna PC. Rouleaux formation in sickle cell traits. *The Journal of Tropical Medicine and Hygiene.* 1991; 94(1):42–44. PMID: 1995933
102. Johnson CS. Arterial blood pressure and hyperviscosity in sickle cell disease. *Hematology/Oncology Clinics.* 2005; 19:827–837. <https://doi.org/10.1016/j.hoc.2005.08.006> PMID: 16214646
103. Cho YI, Mooney MP, Cho DJ. Hemorheological disorders in diabetes mellitus. *Journal of Diabetes Science and Technology.* 2008; 2:1130–1138. <https://doi.org/10.1177/193229680800200622> PMID: 19885302

104. Shin S, Nam J, Hou J, Suh J. A transient, microfluidic approach to the investigation of erythrocyte aggregation: the threshold shear-stress for erythrocyte disaggregation. *Clinical Hemorheology and Microcirculation*. 2009; 42:117–125. <https://doi.org/10.3233/CH-2009-1191> PMID: 19433885
105. Lee B, Ko J, Lim H, Nam J, Shin S. Investigation of critical shear stress with simultaneous measurement of electrical impedance, capacitance and light backscattering. *Clinical Hemorheology and Microcirculation*. 2012; 51:203–212. <https://doi.org/10.3233/CH-2011-1526> PMID: 22240385
106. Flormann D, Aouane O, Kaestner L, Ruloff C, Misbah C, Podgorski T et al. The buckling instability of aggregating red blood cells. *Scientific Reports*. 2017; 7:1–10. <https://doi.org/10.1038/s41598-017-07634-6> PMID: 28801570
107. Ye T, Peng L, Li G. Red blood cell distribution in a microvascular network with successive bifurcations. *Biomechanics and Modeling in Mechanobiology*. 2019; 18:1821–1835. <https://doi.org/10.1007/s10237-019-01179-5> PMID: 31161352
108. Hoover R, Rubin R, Wise G, Warren R. Adhesion of normal and sickle erythrocytes to endothelial monolayer cultures. *Blood*. 1979; 54(4):872–876. <https://doi.org/10.1182/blood.V54.4.872.872> PMID: 476304
109. Hebbel RP, Boogaerts MAB, Eaton JW, Steinberg MH. Erythrocyte adherence to endothelium in sickle-cell anemia: a possible determinant of disease severity. *New England Journal of Medicine*. 1980; 302(18):992–995. <https://doi.org/10.1056/NEJM198005013021803> PMID: 7366623
110. Celada A, Cruchaud A, Perrin LH. Opsonic activity of human immune serum on in vitro phagocytosis of *Plasmodium falciparum* infected red blood cells by monocytes. *Clinical and Experimental Immunology*. 1982; 47(3):635. PMID: 7044626
111. Ishikawa-Sekigami T, Kaneko Y, Saito Y, Murata Y, Okazawa H, Ohnishi H et al. Enhanced phagocytosis of CD47-deficient red blood cells by splenic macrophages requires SHPS-1. *Biochem. Biophys. Res. Co*. 2006; 343:1197–1200. <https://doi.org/10.1016/j.bbrc.2006.03.094> PMID: 16580635
112. Bell GI. Models for the specific adhesion of cells to cells: a theoretical framework for adhesion mediated by reversible bonds between cell surface molecules. *Science*. 1978; 200:618–627. <https://doi.org/10.1126/science.347575> PMID: 347575
113. Dembo M. On peeling an adherent cell from a surface. *Lectures on Mathematics in the Life Sciences, Some Mathematical Problems in Biology*. 1994; 24:51–77.