



# **A Case Report on Aneurysm of the Atypical Right Subclavian Artery: A Rare Embryological Aberration**

**Suresh Rao <sup>a++\*</sup> and Syam Prasad Darla <sup>b#</sup>**

<sup>a</sup> *Department of Anatomy, Subbaiah Institute of Medical Sciences, Shivamogga, India.*

<sup>b</sup> *Department of Radiology, EWMSC, Trinidad.*

## **Authors' contributions**

*This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.*

## **Article Information**

### **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/125740>

**Case Study**

**Received: 26/08/2024**

**Accepted: 29/10/2024**

**Published: 04/11/2024**

## **ABSTRACT**

Atypical or an aberrant right subclavian artery, is the most prevalent vascular anomaly of the aortic arch but a rare embryological aberration. It is typically discovered by accident during imaging tests or autopsy and is usually asymptomatic. Dysphagia and dyspnea are the most common symptoms, which arise when the abnormal artery compresses the surrounding structures. It could be linked to further cardiac or vascular anomalies. We are presenting an 86-year-old woman who has an aneurysm connected to her atypical right subclavian artery. In order to prevent surgical complications, clinicians who undertake thoracic diagnostic and surgical procedures in thorax should pay special attention to the unusual or aberrant right subclavian artery.

<sup>++</sup> Professor;

<sup>#</sup> DM Resident;

<sup>\*</sup>Corresponding author: E-mail: [s4chavan@yahoo.co.in](mailto:s4chavan@yahoo.co.in), [s4chavan@yahoo.in](mailto:s4chavan@yahoo.in);

**Cite as:** Rao, Suresh, and Syam Prasad Darla. 2024. "A Case Report on Aneurysm of the Atypical Right Subclavian Artery: A Rare Embryological Aberration". *Asian Journal of Case Reports in Medicine and Health* 7 (1):156-60. <https://journalajcrmh.com/index.php/AJCRMH/article/view/204>.

**Keywords:** Aberrant subclavian artery; aneurysm; dysphagia; dyspnea.

## 1. INTRODUCTION

“Subclavian artery is the principal artery of the upper limb, but it also supplies a considerable part of the neck and brain through its branches. On the right side it is a branch of the brachiocephalic artery posterior to the sternoclavicular joint. On the left side it is a branch of the arch of aorta from where it ascends on the mediastinal pleura grooving the left lung, and enters neck posterior to the left sternoclavicular joint. Both arteries pursue a similar course in the neck. Each artery arches laterally from the sternoclavicular joint to the outer border of the first rib where it ends by becoming continuous as axillary artery. However in 0.4% to 1.8% of the general population, right subclavian artery can originate directly from the aortic arch, which is located distal to the left subclavian artery. Such a course of the right subclavian artery is conventionally named the ‘lusory artery’. It is also called aberrant right subclavian artery (ARSA) or atypical right subclavian artery” [1-3]. “This anomaly might be an isolated defect or concurrent with other congenital heart defects. As an isolated defect, it is mostly asymptomatic and thus recognized incidentally during investigations or autopsy. The lusory artery forms an incomplete vascular ring and therefore, it seldom causes pressure and dislocation of the esophagus or trachea” [4-7]. The anomaly may be linked to certain clinical signs as chest discomfort, stridor, dysphagia or both (dysphagia lusoria) and dyspnea [8]. Even in cases when this aberration is asymptomatic, one should be aware of the clinical implications of this vascular anomaly especially in patients with dysphagia, dyspnea and who undergo surgical procedures in thorax, such as esophagectomy or vascular surgery and endovascular procedures involving the aortic arch and its branches.

“Arteria lusoria is another name for the anomalous right subclavian artery. With an incidence of between 0.5% and 2.5% in the general population, it is one of the most prevalent aortic arch abnormalities. In 80% of instances, the right subclavian artery begins as the terminal branch of the aortic arch and travels down the right arm after passing through the midline of the body, between the oesophagus and vertebral column and, less frequently, (15% of cases), between the trachea and oesophagus” [9]. “While some people never have any symptoms, others

may experience coughing, pyrosis, dysphagia, dyspnea, acute limb ischaemia, retrosternal discomfort, and even Homer's syndrome” [10]. “The right subclavian artery, which runs posteriorly, and the right common carotid artery, which is positioned anteriorly, compress the oesophagus, resulting in the clinical signs of arteria lusoria” [11].

Several techniques are proposed for the preoperative diagnosis of this anomaly: Particularly in patients with symptoms, a barium swallow may indicate the abnormality. CT and magnetic resonance imaging are two noninvasive angiography techniques that can be excellent confirming tools. This abnormality can be accurately diagnosed with multislice CT.

## 2. CASE REPORT

We report that an 86-year-old woman with persistent and steadily increasing abdominal pain for two weeks was seen at EWMSC, Mount Hope Hospital. She also reported having three months of continuous, progressively worsening chest pain. She feels as though her chest is heavy, and her dysphagia is getting worse. Her dysphagia began with solid food and progressed to liquids over time. Before the presentation, she was unable to swallow any food or liquids and would frequently regurgitate. The patient has been diagnosed with diabetes for 15 years and is currently taking 500 mg of metformin once daily as part of their normal prescription regimen, no noteworthy personal or family history [12,13].

The patient's CT scan revealed an incidental finding of a vascular anomaly, that the right subclavian artery originating from the arch of aorta as the last branch with aneurysmal dilation was confirmed. The aneurysm displaced the trachea and the surrounding upper portion of the esophagus anteriorly, compressing them both significantly (Fig. 1-3).



## 3. DISCUSSION


Normally during embryogenesis, regression of the distal right dorsal aorta occurs leading to the right 4<sup>th</sup> aortic branch and 7<sup>th</sup> intersegmental artery becoming the right subclavian artery. On the left side the subclavian artery is derived entirely from the seventh cervical intersegmental artery, which arises from the dorsal aorta opposite the attachment of the fourth arch artery.

Aberrant right subclavian artery is caused by the involution of the right 4<sup>th</sup> aortic branch and proximal right dorsal aorta with a persistent distal right dorsal aorta and 7<sup>th</sup> intersegmental artery forming the right subclavian artery. This causes the right subclavian artery to originate left of the midline occasionally causing compression. The retroesophageal course of the right subclavian artery behind the esophagus, although usually asymptomatic, may cause compression and a type of dysphagia known as dysphagia lursoria [14-16].

The lusoria artery occasionally emerges from an aortic arch diverticulum at the proximal descending aorta, which Kommerell initially identified as the Kommerell diverticulum in 1936 [17,18]. Kommerell's diverticulum can be found in anywhere between 15% and 60% of Lusoria artery instances [19]. Aneurysms are almost always found at or near the origin of the artery, and are suspected to result from degeneration of a diverticulum of Kommerell. Because these aneurysms can rupture, thrombose, or embolize even if they are asymptomatic, they should always be treated [20].

The radiographic results of 25 cases of aberrant right subclavian artery that were first shown on CT were assessed by Hara et al. [21]. In 95% of the patients with right regtrotracheal subclavian artery, he found that the retrotracheal course of the aberrant artery could result in a posterior tracheal impression, which manifests as a vascular retrotracheal opacity [8]. A case of right regtrotracheal subclavian artery producing asthma was reported by Parker et al. [9]. The trachea's anterior displacement from right regtrotracheal subclavian artery may be the cause of the asthma. When an angiographer approaches the ascending thoracic aorta via the right axillary, brachial, or radial approaches, right regtrotracheal subclavian artery is equally crucial from a clinical standpoint [10,11]. When ascending aorta catheterization becomes challenging, the anomalous right subclavian artery may be the cause [17]. The brachiocephalic trunk, the first branch of the aortic arch, allows direct access to the ascending aorta, making it simple to reach it by the right radial artery. However, things change and angiography becomes a difficult task when an aberrant right subclavian artery is present [18].





**Fig. 1-3. Vascular anomaly of right subclavian artery**

It is well recognized that chromosomal abnormalities, particularly trisomy 21, are linked to aberrant right subclavian artery. According to a recent ultrasonography research, 37.5% of fetuses with Down's syndrome had an aberrant right subclavian artery, when they were between 13 and 26 weeks gestation [22,23]. Apart from the fact that aberrant right subclavian artery is linked to a higher frequency of intra-cardiac abnormalities, Borenstein et al. report that aberrant right subclavian artery is relatively more common in fetuses with chromosomal disorders, specifically trisomy 21, compared to euploid fetuses. They conclude by saying that right subclavian artery position evaluation is probably going to become a standard ultrasonography diagnostic for chromosomal abnormalities in the second trimester of pregnancy [19,20].

#### **4. CONCLUSION**

A proper understanding of the anatomic and morphologic variations of the aortic arch is imperative for clinicians performing diagnostic and surgical procedures in the thorax, particularly with regard to the atypical or aberrant right subclavian artery, given the increasing number of imaging studies and catheter-based treatments performed these days.

#### **DISCLAIMER (ARTIFICIAL INTELLIGENCE)**

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

#### **CONSENT**

As per international standards or university standards, patient(s) written consent has been collected and preserved by the author(s).

#### **ETHICAL APPROVAL**

As per international standards or university standards written ethical approval has been collected and preserved by the author(s).

#### **COMPETING INTERESTS**

Authors have declared that no competing interests exist.

#### **REFERENCES**

1. Dandelooy J, Coveliers JP, Van Schil PE, Anguille S. Dysphagia lusoria. *CMAJ* 2009;181:498.
2. Hart PA, Karnath PS. Dysphagia lusoria. *Mayo Clin Proc.* 2012;87:e17.
3. Pramesh CS, Saklani AP, Parmar V, Acharya S, Badwe RA. Aberrant subclavian artery causing difficulty in transhiatal esophageal dissection. *Dis Esophagus.* 2003;16:173-176.
4. Skalski JH, Religa Z: *Kardiochirurgia dziecięca.* Wydawnictwo Śląsk. 2003;2:361–372.
5. Saito T, Tamatsukuri Y, Hitosugi T et al: Three cases of retroesophageal right subclavian artery. *J Nippon Med Sch.* 2005;72:375–382.
6. Epstein DA, Debord JR: Abnormalities associated with aberrant right subclavian arteries-a case report. *Vasc Endovascular Surg.* 2002;36:297–303.
7. Czerwinski F, Mierzwa A, Krzanowski K et al: Rare cases of arteries branching directly from aortic arch in the man. *Folia Morphol.* 1993;52:223–231.
8. Abraham V, Mathew A, Cherian V, Chandran S, Mathew G. Aberrant subclavian artery: Anatomical curiosity or clinical entity. *Int J Surg.* 2009;7:106-109.

9. Amore D Casazza D. Casalino A. et al. Symptomatic aberrant right subclavian artery advantages of a lessinvasive surgical approach. Ann. Thorac Cardivasc Surg. 2020;26:104-107.
10. Coles M Madray VM Mareddy C et al. Dysphagialusoria a vascular etiology? JGH Open. 2020; 4:1238-1239.
11. Loukas M, LOuis RG, Gaspard J, Fudalej M, Tubbs RS, Merds W. A retrotracheal right subclavian art+ry in asso<ciation with a verteLral artery and thyroidea ima. Folia Morphol. 2006;65(3):236-241.
12. Zalel Y, Achiron R, Yagel S, Kivilevitch Z. Fetal aberrant right subclavian artery in normal fetuses ancl dc>wn syndrome. Ultrasound Obstet Gy/1eco/. 2008;1:25-29.
13. Borenstein M, Minekawa R, Zidere V, Nicolaidis K, Allan LD. Aberrant right subclavian artery at 16 to 23+6 weeks of gestation: a marker for chromosomal abnormality. Ultrasound Obstet Gynecol. 2010;36(5):548-552.
14. Abhaikzhand RK, LoLivard Y, Gobeil OF, Loubeyre C, Lefevre T, Morice MC. The problem of arteria lusoria in radial transradial coronary angiography and angioplasty. Catheter Cardiovasc Interv. 2001;54:196-201.
15. Campeau L. Percutaneous radial approach for coronary angiography. Cathet Cardiovasc Diagn. 1999;16:3-7.
16. Jung Ho Huh, Kook-Jin Chun, Moo Hyun Kim. Overcome the tortuosity from radial artery to subclavian artery-Tips and Tricks, TRI manual. Available:<http://www.w-tri.org/images/emanual-0107.PDF>.
17. Ocaya A Retrooesophageal right subclavian artery a case report and review of the literature. Afr Health Sci. 2015; 15:1034-1037.
18. Kommerell BF. Verlagerung des Oesophagus durch eine abnorm verlaufende Arteria subclavia dextra (*Arteria lusoria*). Forstsch. Geb. Roetgenstr. 1936;54:590-595.
19. Brown DL, Chapman WC, Edwards WH, Coltharp WH, Stoney WS. Dysphagia lusoria: aberrant right subclavian artery with aKommerell's diverticulum. Am. Surg. 1993;59(9):582-586.
20. Polguy M, Chrzanowski L, Kasprzak JD, Stefanczyk L, Topol M, Majos A. The aberrant right subclavian artery (Arteria lusoria): the morphological and clinical aspects of one of the most important variations--a systematic study of 141 reports. ScientificWorldJournal. 2014;94:1010-1011.
21. Hara M, Satake M, Itoh M, Ogino H, Shiraki N, Taniguschi H, Shibamoto Y. Radiographic findings of aberrant right subclavian artery initially depictedon CT. Radiat Med. 2003;21:161-65.
22. Kieffer E, Bahnini A, Koskas F. Aberrant subclavian artery: surgical treatment in thirty-three adult patients. J. Vasc. Surg., 1994;19(1):100-109.
23. Parker JM, Gary-Freitas B, Berg BW. Symptomatic vascular rings in adulthood: an uncommon mimic of asthma. J Asthma. 2000;37:275-280.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/125740>